

NOTA PALEONTOLOGICA

Parrots (Aves, Psittaciformes) in the Pleistocene of Uruguay



Claudia P. TAMBUSSI¹, Carolina ACOSTA HOSPITALECHE¹, Andrés RINDERKNECHT² and Martín UBILLA³

Introduction

Cyanoliseus Bonaparte, 1854 (Psittaciformes, Psittacidae, Arini) is the most frequent parrot in the Pleistocene record of Argentina. It comprises three species: the extinct *C. ensenadensis* (Cattoi, 1957) Tonni, 1972 and *C. patagonopsis* Acosta Hospitaleche and Tambussi, 2006, and the extant *C. patagonus* (Vieillot, 1817). A complete list of the Pleistocene Psittacidae known at the present is given by Acosta Hospitaleche and Tambussi (2006) and Tambussi *et al.* (2007). Nowadays, the burrowing parakeet *Cyanoliseus patagonus* (Psittacidae, Arini) is found in Argentina and the center of Chile, occasionally reaching Uruguay in winter (Collar, 1997; Bucher and Rodriguez, 1986) (figure 1). This paper reports the presence of *Cyanoliseus* in the Pleistocene of Uruguay, the first fossil parrot recorded in this country; comments about the environmental are also reported.

Materials and methods

The material under study was compared with all living species inhabitant Uruguay and representatives of all genera of parrots living in Argentina (Appendix 1). Osteological terminology from Baumel and Witmer (1993) was used for the descriptions and their English equivalents through the discussion. Measurements are in millimeters with 0.1 mm accuracy.

The specimens studied are housed at the Museo Nacional de Historia Natural y Antropología, Montevideo, Uruguay (MNHN); Paleontología Vertebrados of the Facultad de Ciencias,

Universidad de la República, Uruguay (FC-DPV); Museo de La Plata, La Plata, Argentina (MLP); Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina (MACN).

Systematic paleontology

Order PSITTACIFORMES Wagler, 1830

Family PSITTACIDAE Illiger, 1811

Tribe ARINI Smith, 1975

Genus *Cyanoliseus* Bonaparte, 1854

Type species. *Cyanoliseus patagonus* (Vieillot, 1817).

Remarks. The presence of the following associated characters is suitable to identify both humeri as *Cyanoliseus*: (1) deep and wide *incisura capitidis*, (2) presence of a small tubercle in the center of the distal edge of the humerus head, (3) well developed crista deltopectoralis and (4) a deep *incisura intercondylaris*.

Comparative description. The distal extension of the crista deltopectoralis is larger than that of the crista bicipitalis, while in *Brotogeris*, *Pionus*, *Pyrrhura*, *Forpus* and *Pionopsitta*, both cristae are similar in their extension. As with other species of *Cyanoliseus*, the small tubercle in the center of the internal distal edge of the humerus head is present. The *sulcus ligamentorum transversus* is deeper and wider than in *Amazona*, *Ara*, *Nandayus*, *Aratinga*, *Myiopsitta*, *Brotogeris*, *Pionus*, *Pyrrhura*, *Pionopsitta*, *Anodorhynchus* and *Forpus*. The *fossa musculus brachialis* is deeper than in *Ara*, *Aratinga*, *Amazona*, *Pionus*, *Pionopsitta*, *Pyrrhura*, *Myiopsitta*, *Forpus*, *Anodorhynchus* and *Brotogeris*. The *condylus dorsalis* is elongated, while in *Amazona*, *Ara*, *Aratinga*, *Myiopsitta*, *Brotogeris*, *Pionus*, *Pyrrhura*, *Pionopsitta*, *Anodorhynchus* and *Forpus*, it is rounded. The *condylus ventralis* is oval and asymmetric, whereas in *Amazona*, *Ara*, *Brotogeris*, *Pionus*, *Pyrrhura*, *Pionopsitta*, and *Forpus* is symmetric.

The *incisura intercondylaris* is wider than in

AMGHB2-0002-7014/09\$00.00+.50

¹CONICET. División Paleontología Vertebrados, Museo de La Plata, Paseo del Bosque s/n, B1900FWA La Plata, Argentina. tambussi@museo.fcnym.unlp.edu.ar; acostacaro@museo.fcnym.unlp.edu.ar

²Museo Nacional de Historia Natural y Antropología, Casilla de Correo 399, 11.000 Montevideo, Uruguay. rinderk@adinet.com.uy

³Facultad de Ciencias, Universidad de la República, Iguá 4225, 11400 Montevideo, Uruguay. ubilla@fcien.edu.uy




Figure 1. Location map showing the fossiliferous locality in Uruguay (A: *Cyanoliseus ensenadensis*; B: *C. patagonus*). Shaded area indicates the current area of occurrence of *Cyanoliseus patagonus* in Uruguay / mapa de Uruguay señalando la ubicación de las localidades donde fueron exhumados los fósiles (A: *Cyanoliseus ensenadensis*; B: *C. patagonus*). El área sombreada indica el sector actual de distribución de *Cyanoliseus patagonus*.

Amazona, *Ara*, *Nandayus*, *Aratinga*, *Myiopsitta*, *Brotogeris*, *Pionus*, *Pyrrhura*, *Pionopsitta*, *Anodorhynchus* and *Forpus*, and the *incisura capitidis* is wider and deeper than in *Amazona*, *Ara*, *Nandayus*, *Aratinga*, *Brotogeris*, *Pionus*, *Pyrrhura*, *Pionopsitta*, *Anodorhynchus* and *Forpus*.

Cyanoliseus patagonus (Vieillot, 1817)

Figure 2

Material. Complete right humerus (MNHN 1716).

Geographic locality. Department of Canelones, Balneario San Luis (Interbalnearia Route, km 64.5), Uruguay; coastal cliffs near the Río de la Plata estuary (34° 47' S; 55° 37' W) (figure 1).

Stratigraphic provenance. Libertad Formation (Pleistocene).

The fossil comes from a stratigraphic level composed by quartz-feldspar sands (reworked loess) at 90 cm under a horizon of calcretes (paleosoil). In this place the sediments -referred to the Libertad Formation- are composed of loess, reworked loess, brownish siltstone, paleosoils, and fluvial deposits (Loureiro *et al.*, 2002). Further details about the age of these sediments are not possible at this stage. However, on the basis of its stratigraphic relationships, some authors have considered to be lower to middle Pleistocene in age (Bossi and Navarro, 1991; Panario and Gutiérrez, 1999).

Comments. According to the compared material, the

humerus length of living *C. patagonus* ranges from 47 to 51 mm (n=8). The length of the humerus of the MNHN-1716 specimen is included into the confidence limits of the mean of *C. patagonus* which indicates a non significant statistical difference between the fossil remains and the available sample for this character. Its size is intermediate between *C. patagonopsis* and *C. ensenadensis*.

Some differences allow us to differentiate it from *C. patagonopsis*: the *tuberculum dorsale* is proportionally bigger, the *condylus dorsalis* is more rounded and its edges are less acute, the *condylus ventralis* is more asymmetric, the *tuberculum supracondylare ventrale* is less robust, and the *sulcus ligamentorum transversus* is not so expanded.

Cyanoliseus ensenadensis (Cattoi, 1957)

Figure 3

Material. Right humerus without distal portion (FC-DPV-1417).

Geographic locality. Sopas creek, Salto Department (31°15'S 57°00'W), Uruguay (figure 1).

Stratigraphic provenance. Brownish mudstones of the Sopas Formation (late Pleistocene). This unit is widespread in northern Uruguay and is in general characterized by massive brownish mudstones, sandstones and conglomerates formed mainly in a fluvial environment, but also including paleosoils (Ubilla *et al.*, 2004).

Comments. *C. ensenadensis* is the smallest species of




Figure 2. Humerus of *Cyanoliseus patagonus* MNHN 1716. 1, cranial view; 2, caudal view / húmero de *Cyanoliseus patagonus* MNHN 1716. 1, vista craneal; 2, vista caudal. cb, crista bicipitale; cd, condylus dorsalis; cdp, crista deltopectoralis; cv, condylus ventralis; ic, incisura capitidis; sl, sulcus ligamentorum transversus; tb, tuberculum dorsale; tsv, tuberculum supracondylare ventrale.

the genus *Cyanoliseus*. The studied humerus is smaller than *C. patagonus*. The holotype, until now the only known specimen, measures 44.7 mm length. Although the *sulcus ligamentorum transversus* is small in this genus, it is even less expanded in *C. patagonopsis* and *C. patagonus*.

Discussion and conclusions

The Pampean Region of *Cyanoliseus* indicates that the three species lived during the Pleistocene, but only the two smallest species have been found in Uruguay: *C. ensenadensis* and *C. patagonus*.

Coincidentally, all fossil species of *Cyanoliseus* are diagnosed using humeri, which are useful bones in the identification of these taxa. The humeral characters are: 1- a distal extension of the deltopectoral crest bigger than the bicipital crest, 2- a small tubercle in the center of the internal distal edge of the humerus head, 3- a deep and wide transverse ligamentary groove, 4- a deep brachial fossa, 5- an oval and asymmetric ventral condyle, 6- an enlarged dorsal condyle with its proximal end more acute, 7- a wide and deep capital groove, and 8- a broad dorsal tubercle.

The MNHN 1716 is morphologically indistinguishable from that of modern species. *Cyanoliseus patagonus* is a winter migrant whose presence in Uruguay was confirmed at the end of the 70's (Cuello, 1985; Azpiroz, 2001), and is recorded today in southwest of Uruguay, but only sporadically (Claramunt and Cuello, 2004) (figure 1).

The associated paleofauna with *Cyanoliseus patagonus* recovered from Libertad Formation at San Luis locality includes passerine birds, colubrids, and anurans (Rinderknecht, 1998, Claramunt and Rinderknecht, 2005), a woodpecker of the genus *Colaptes* Vigors, 1826 Rinderknecht and Claramunt, 2000), rodents of the genera *Akodon* Meyen, 1833 and *Galea* Meyen, 1831, and an assemblage of typical Pleistocene mammals (Ubilla and Rinderknecht, 2001). The co-occurrence of the likely grazer *Glyptodon* Owen, 1839, the grass mouse *Akodon*, the cui *Galea* and this parrot supports, the idea of the presence of open or semiopen environments, considering the environmental adaptations inferred or known for these taxa (Rood, 1972; Walker, 1975; Scillato-Yané et al., 1995; Fariña and Vizcaíno, 2001). Living burrowing parrots almost always nest in holes (Masello and Quillfeldt, 2002), in the banks of rivers and other steep. An analysis related to environmental variables that could constrain the distribution of living representatives of *C. patagonus*, indicates subhumid to semiarid climates associated with 600-800 mm of annual precipitation (Tambussi et al., 2007).

The other fossil parrot described here belongs to



Figure 3. Humerus of *Cyanoliseus ensenadensis* FC-DPV-1417. 1, cranial view; 2, caudal view / húmero de *Cyanoliseus ensenadensis* FC-DPV-1417. 1, vista craneal; 2, vista caudal.

C. ensenadensis, characterized by its size and the transversal ligament groove which is less expanded than that of *C. patagonopsis*. *Cyanoliseus ensenadensis* is the smallest species of the genus, being a third smaller than the living species *C. patagonus* and two thirds smaller than the extinct *C. patagonopsis* (Acosta Hospitaleche and Tambussi, 2006).

Fauna associated with the humerus FC-DPV-1417 attributed to *C. ensenadensis* includes a variety of fossil vertebrates such as glyptodonts, horses like *Equus neogeus* and *Hippidion principale*, *Machaeraneria patachonica*, *Toxodon platensis*, deer like *Ozotoceros*, mostly related to open or semiopen environments (Ubilla, 2004; Ubilla et al., 2004) with a TL/OSL age of 43.5 +/- 3.6 ka BP (Ubilla, 2004). Earthworm aestivation chambers described from this layer indicates a seasonal climate (Verde et al., 2007). In the Sopas Formation birds were found from different outcrops: aff. *Rhea* sp. Brisson, 1760, *Chloephaga picta* (Gmelin, 1789) and *Cariama cristata* (Linnaeus, 1766) (Ubilla et al., 2004; Tambussi et al., 2005). Open to semiopen environment under seasonal climatic conditions are suggested by paleoecological evidence (Verde et al., 2007) and the record of seriemas and magellan geese. According to mammalian biostratigraphic information, the faunal assemblage is correlated to the Lujanian Age/Stage of the Buenos Aires province, and ¹⁴C ages are >43 ky BP (Ubilla et al., 2004), which is older than the fauna of the Guerrero Member of the Lujan Formation (between 21ka and 10 ka BP, Cione and Tonni, 1999; Tonni et al., 2003) of the Pampean Region.

In Argentina, *Cyanoliseus ensenadensis* has been recorded at Olivos (Ensenadan Formation) instead assignable to the Ensenadan stage (early to middle Pleistocene). Biostratigraphically, they are defined by

the presence of fossil mammals such as *Mesotherium cristatum* Serrés, 1867, *Arctotherium angustidens* Gervais and Ameghino, 1880, *Theriodictis platensis* Mercerat, 1891, *Glyptodon munizi* Ameghino, 1881, *Neosclerocalyptus pseudornatus* (Ameghino, 1889) Paula Couto, 1957 and *Eutatus seguini* Gervais, 1867. Taking into account biostratigraphic and absolute ages available (Ubilla, 2004), the record of *C. ense-nadensis* in Uruguay belongs to later sediments than those of Argentina.

Until now, the fossil record of *Cyanoliseus* was limited to the Pleistocene of Argentina. Consequently, the material reported here not only expands the geographic range of *Cyanoliseus* during the Pleistocene, but is also the first time that any fossil Psittacidae has been reported from Uruguay. Therefore, the record of these parrots increases our knowledge of the Uruguayan Pleistocene avifauna, where Psittaciformes have never previously been found.

Acknowledgments

To Santiago Claramunt who help us to study the materials from Uruguay. P. Tubaro (MACN) who facilitated comparisons with the extant parrots of the MACN under his care. D. Waterhouse and an anonymous reviewer provided helpful comments. It is a contribution to PICT 32617 (C. Tambussi), CSIC Project (M. Ubilla) and IGCP-518 (Bridgland-Latrubesse).

References

- Acosta Hospitaleche, C. and Tambussi, C. 2006. *Cyanoliseus patagonopsis* nov. sp (Aves, Psittaciformes) del Pleistoceno Medio de Punta Hermengo, provincia de Buenos Aires. *Ameghiniana* 43: 249-253.
- Ameghino, F. 1881. *La antigüedad del hombre en el Plata*. Masson-Igon Hermanos 2: 557.
- Ameghino, F. 1889. Contribución al conocimiento de los mamíferos fósiles de la República Argentina. *Actas de la Academia Nacional de Ciencias* (Córdoba) 6: 1-1027.
- Azpiroz, A.B. 2001. *Aves del Uruguay. Lista e introducción a su biología y conservación*. Aves Uruguay-GUPECA. Graphis Editorial Montevideo, 1-104 pp.
- Baumel, J.J. and Witmer, L.M. 1993. Osteology. In: J. Baumel, A. King, J. Breazile, H. Evans and J. Vanden Berge (eds.), *Handbook of Avian Anatomy*. Publications of the Nuttall Ornithological Club, Harvard University. Cambridge, Massachusetts: 45-132.
- Bonaparte, C. 1854. Tableau des Perroquets. *Revue et Magasin de Zoologie Pure et Appliquée*. Series 2 vol. 1 1854 (Meetings of Nov. 13, 20, 27, Dec. 4, 11, 18, 25): 145-158.
- Bossi, J. and Navarro, R. 1991. Geología del Uruguay. UDELAR-Dpto. *Publicaciones*. Montevideo 2: 463-966.
- Bucher, E.H. and Rodriguez, E.N. 1986. Sobre la presencia del loro barranquero (*Cyanoliseus patagonus*) en el Uruguay. *El Hornero* 12: 303-304.
- Cattoi, N. 1957. Contribución al conocimiento de la avifauna extinguida del Pleistoceno de la República Argentina. *Ameghiniana* 1: 17-24.
- Cione, A. and Tonni, E. 1999. Biostratigraphy and chronological scale of upper-most Cenozoic in the Pampean Area, Argentina. In: M. Salemme (ed.), *Quaternary of South America and Antarctic Peninsula*. Balkema, Rotterdam, Brookfield, pp. 23-51.
- Claramunt, S. and Cuello, J.P. 2004. Diversidad de la biota Uruguaya: Aves. *Anales del Museo Nacional de Historia Natural y Antropología* 2ª Serie 10: 1-76.
- Claramunt, S. and Rinderknecht, A. 2005. A new fossil furnariid from the Pleistocene of Uruguay, with remarks on nasal type, cranial kinetics, and relationships of the extinct genus *Pseudoseisuropsis*. *The Condor* 107: 114 - 127.
- Collar, N.J. 1997. Family Psittacidae (Parrots). In: J. del Hoyo, A.E. Elliot and J. Sargatal (eds.), *Handbook of the Birds of the World*, vol. 4. Sandgrouse to Coockos. Lynx Ediciones, Barcelona, pp. 280-477.
- Cuello, J. 1985. Lista de referencia y bibliografía de las aves uruguayas. *Museo Dámaso Antonio Larrañaga*, Serie de Divulgación, Montevideo 1: 1-116.
- Gervais, P. 1867. Sur une nouvelle collection d'ossements fossiles de Mammifères recueillié par M. Fr. Seguin dans la Confédération Argentine. *Comptes rendus des séances de l' Académie des Sciences* 65: 279-282, Paris.
- Gervais, H. and Ameghino, F. 1880. *Los mamíferos fósiles de la América del Sur*. Sabih e Igon, París y Buenos Aires: 225.
- Fariña, R. and Vizcaíno, S.F. 2001. Carved teeth and strange jaws: How glyptodonts masticated. *Acta Palaeontologica Polonica* 46: 219-234.
- Loureiro, J., Rinderknecht, A. and Sánchez, I. 2002. Estudio de magnetismo de rocas en sucesiones cenozoicas de Uruguay. *Revista de la Sociedad Uruguaya de Geología* III época 9: 29-41.
- Masello, J. and Quillfeldt, P. 2002. Chick growth and breeding success of the burrowing parrot. *The Condor* 104: 574-586.
- Mercerat, A. 1891. Caracteres diagnósticos de algunas especies de Creodonta conservadas en el Museo de La Plata. *Revista del Museo de La Plata* 2: 51-52.
- Owen, R. 1839. Fossil Mammalia, (2). In: C. Darwin (ed.), *The Zoology of the voyage of H.M.S. Beagle under the command of Captain Fitzroy, N. R., during the years 1832 to 1836*, Smith, Elder and Co., London. 1: 41-64.
- Panario, D. and Gutiérrez, O. 1999. The continental uruguayan Cenozoic: an overview. *Quaternary International* 62: 75-84.
- Paula Couto, J.C. 1957. Sobre um gliptodonte do Brasil. *Boletim Divisão de Geologia e Mineralogía Rio de Janeiro* 165: 1-37.
- Rinderknecht, A. 1998. Nuevos microvertebrados fósiles para el Pleistoceno superior del Uruguay (Amphibia, Reptilia, Aves). *Comunicaciones Paleontológicas del Museo de Historia Natural de Montevideo* 2: 133-144.
- Rinderknecht, A. and Claramunt, S. 2000. Primer registro de *Colaptes Vigors*, 1826, para el Pleistoceno del Uruguay (Aves: Piciformes: Picidae). *Comunicaciones Paleontológicas del Museo de Historia Natural de Montevideo* 2: 157-160.
- Rood, J. 1972. Ecological and behavioural comparisons of three genera of argentine cavies. In: J. Cullen and C. Beer (eds.), *Animal Behavior Monographs*, Part 1 London. Baillière Tindall, pp. 1-83.
- Scillato-Yané, G.J., Carlini, A., Vizcaíno, S.F. and Ortiz Jaureguizar, E. 1995. Los Xenarthros. In: M.T. Alberdi, G. Leone and E.P. Tonni (eds.), *Evolución biológica y climática de la Región Pampeana durante los últimos cinco millones de años. Un ensayo de correlación con el Mediterráneo Occidental*, Monografías del Museo Nacional de Ciencias Naturales 12, CSIC (España): pp. 181-211.
- Serres, M. 1867. De l'ostéographie du *Mesotherium* et ses affinités zoologiques. *Comptes Rendu de l' Académie des Sciences* 65: 6-848.
- Tambussi, C., Ubilla, M., Acosta Hospitaleche, C. and Perea, D. 2005. Fossil records and palaeoenvironmental implications of *Chloephaga picta* (Gmelin, 1789) (Magellan Goose) and *Cariama cristata* (Linnaeus, 1766) (Seriema) from the Late Pleistocene of Uruguay. *Neues Jahrbuch Geologie und Paläontologie Mh.* 5: 257-268.

- Tambussi, C., Acosta Hospitaleche, C. and Horlent, N. 2007. La avifauna del cuaternario de Argentina: inferencias paleoambientales a partir del registro de los Psittacidae. In: G. Pons and D. Vicens (eds), *Geomorfologi Litoral i Quaternari*, Monografías Societat D'Historie Natural De Balears, pp. 69-80.
- Tonni, E.P. 1972. *Cyanoliseus ensenadensis* (Cattoi) nov. comb. (Aves, Psittaciformes) del Pleistoceno medio de la provincia de Buenos Aires. *Ameghiniana* 9: 145-148.
- Tonni, E., Huarte, R., Carbonari, J. and Figini, A., 2003. New radiocarbon chronology for the Guerrero Member of the Luján Formation (Buenos Aires): paleoclimatic significance. *Quaternary International* 109-110: 45-48.
- Ubilla, M. 2004. Mammalian biostratigraphy of Pleistocene fluvial deposits in northern Uruguay, South America. *Proceedings of the Geologists' Association London* 115: 347-357.
- Ubilla, M. and Rinderknecht, A. 2001. El género *Galea* Meyen, 1831 (Rodentia, Caviidae) en el Pleistoceno de Uruguay, primeros registros y descripción de una nueva especie extinguida. *Boletín Real Sociedad Española Historia Natural Serie Geológica* 96: 111-122.
- Ubilla, M., Perea, D., Goso, C. and Lorenzo, N. 2004. Late Pleistocene vertebrates from northern Uruguay: tools for biostratigraphic, climatic and environmental reconstruction. *Quaternary International* 114: 129-142.
- Verde, M., Ubilla, M., Jiménez, J. and Genisse, J. 2007. A new earthworm trace fossil from palaeosols: aestivation chambers from the late pleistocene sopas formation of Uruguay. *Palaeogeography, Palaeoclimatology, Palaeoecology* 243: 339-347.
- Vieillot, L. 1817. *Nouveau Dictionnaire d'Histoire Naturelle Appliquée aux Arts*. First ed. 1803-4, Nouv Ed. 1816-19: p.367.
- Walker, E. 1975. *Mammals of the World*. John Hopkins University Press Baltimore 2, pp. 647-1500.

Recibido: 24 de junio de 2008.

Aceptado: 25 de marzo de 2009.

Appendix I. List of extant species and materials used for comparisons. Fossil specimens are indicated with # / lista de especies y materiales utilizados para las comparaciones. Los especímenes fósiles están indicados con #.

Amazona aestiva (Linnaeus, 1758) MLP 621, MACN 27741, 17308; *Ara chloropterus* Gray, 1859 MACN 53522, 20986; *Ara militaris* (Linnaeus, 1766) MLP 408; *Aratinga acuticaudata* (Vieillot, 1818) MLP 68; *Aratinga aurea* (Gmelin, 1788) MLP 342, MACN 54878; *Aratinga leucophthalma* (Statius Muller, 1776) MLP 407, MACN 20404; *Anodorhynchus hyacinthinus* (Latham, 1790) MACN 23591, 23590; *Brotogeris versicolurus* MLP 348, MACN 18238; *Cyanoliseus patagonus* (Vieillot, 1817) MLP 65, MLP 260 a-g (n=8) and #MLP 81-VII-20-21, *C. ensenadensis* #MACN 17716, *C. patagonopsis* #MLP 81-VII-20-20, *Forpus passerinus* (Linnaeus, 1758) MACN 54856, *Myiopsitta monachus* Boddaert, 1783 MLP 66, 262, *Nandayus nenday* (Vieillot, 1823) MLP 257, 405, 406, *Pionopsitta pileata* (Scopoli, 1769) MLP 67, *Pionus maximiliani* (Kuhl, 1820) MLP 343, MACN 17309, *Pyrrhura frontalis* (Vieillot, 1818) MLP 351, MACN 17307.