

## First record of *Gymnogeophagus lipokarenos* Malabarba, Malabarba & Reis, 2015 (Teleostei: Cichliformes) from Argentina.

Jorge Casciotta<sup>1,2,\*</sup>, Adriana Almirón<sup>1</sup>, Štěpánka Řičanová<sup>3</sup>, Klára Dragová<sup>3</sup>, Lubomir Piálek<sup>3</sup>, Felipe Alonso<sup>4</sup> & Oldřich Řičan<sup>3</sup>

<sup>1</sup> UNLP, Facultad de Ciencias Naturales y Museo, División Zoología Vertebrados, Paseo del Bosque, 1900 La Plata, Buenos Aires, Argentina

<sup>2</sup> CIC, Comisión de Investigaciones Científicas de la Provincia de Buenos Aires, Argentina

<sup>3</sup> University of South Bohemia, Faculty of Science, Department of Zoology, Branišovská 31, 37005 České Budějovice, Czech Republic

<sup>4</sup> CONICET - Instituto de Bio y Geociencias del NOA (IBIGEO), CCT-Salta, 9 de Julio 14, 4405 Rosario de Lerma, Salta, Argentina

\* jcasciotta@gmail.com

### Abstract

*Gymnogeophagus lipokarenos* is registered for the first time from Argentina in the Province of Misiones. The species was originally described as endemic to the Upper Uruguay freshwater ecoregion in Brazil. Our findings extend its distribution into the Lower Uruguay ecoregion throughout the province of Misiones.

### Introduction

The genus *Gymnogeophagus* inhabits the La Plata basin, Laguna dos Patos system, and rio Tramandaí basin, plus a single record of *G. balzanii* is from the rio Guaporé in the Amazon basin (Malabarba et al., 2015; Loureiro et al., 2016). *Gymnogeophagus* is currently diagnosed by the presence of a forward directed spine on the top of the first dorsal pterygiophore and the loss of bony supraneurals (Reis & Malabarba, 1988). The species of this genus show two reproductive strategies: substrate brooding and mouth brooding. The later one is present in species which are informally known as the "*G. gymnogenys* species group". Two years ago, Malabarba et al. (2015) described five new species belonging to this species group from the freshwaters of Brazil and Uruguay, among them *G. lipokarenos*. The aim of this contribution is to register this species for the first time for Argentina, from the río Uruguay basin in Misiones.

### Examined material

*Gymnogeophagus lipokarenos* (figs. 1-4, 7): all lots from Argentina, Misiones province, Uruguay river basin: MLP 11251, 2 exs., 82,7-86,2 mm SL, arroyo Paraiso (27°2'52,49"S 54°5'54,97"W) 22.11.2016 (fig. 1-2). MLP 11252, 3 exs., 116,3-127,0 mm SL, lower arroyo Paraiso (27°9'26,58"S 54°4'0,46"W) 24.11.2016. MLP 11253, 4 exs., 72,8-127,4 mm SL, arroyo Guerrero (arroyo López) (27°45'57,4"S 55°09'33,7"W), 03.12.2007 (fig. 3). MLP 11254, 2 exs., 80,0-87,3 mm SL, arroyo Fortaleza (26°45'56,6"S 54°10'57,4"W), 01.12.2007. MLP 11255, 5 exs., 61,8-87,2 mm SL, arroyo Shangai (27°28'13,8"S 54°41'24,5"W). MACN-Ict 12306, 6 exs., 61,6-94,5 mm SL, arroyo El Saltito, Salto Golondrinas (27°7'50,34"S 54°29'22,54"W) 23.11.2016 (fig. 4). MACN-Ict 12304, 2 exs., 103,8-117,0 mm SL, arroyo El Saltito, Salto Caracol (27°9'27,40"S 54°38'19,53"W), 25.11.2016. MACN-Ict 12305, 1 ex., 133,7 mm SL, arroyo Toro (26°36'32,8"S 53°44'13,9"W), 10.12.2014. IBIGEO-I 448, 5 exs., 88,0-118,6 mm SL, arroyo Toro (26°36'32,8" S, 53°44'13,9" W), 01.12.2016. IBIGEO-I 450, arroyo Melo (27°25'2,67"S 54°42'7,93"W), 13.11.2016 (fig. 7). CI-FML 7264, 4 ex., 39,2-108,8 SL, same collecting data as previous lot.



fig. 1. *Gymnogeophagus lipokarenos*, MLP 11251, arroyo Paraiso




fig. 2. *Gymnogeophagus lipokarenos*, MLP 11251, arroyo Paraiso, close up of color of caudal fin



fig. 3. *Gymnogeophagus lipokarenos*, MLP 11253, arroyo Guerrero



fig. 4. *Gymnogeophagus lipokarenos*, MACN-Ict 12306, arroyo El Saltito, Salto Golondrinas



fig. 5. Arroyo Fortaleza, middle Uruguay basin



fig. 6. Arroyo Toro, upper Uruguay basin

### Distribution

*Gymnogeophagus lipokarenos* is widely distributed in the Upper río Uruguay basin and has been described as endemic to this section of the river basin (Malabarba et al., 2015). However, some paratypes have been included in the original description (Malabarba et al., 2015) which were collected in the Middle río Uruguay basin (*sensu* Zaniboni Filho & Schulz; 2003; Lower Uruguay *sensu* Abell et al. (2008) in Brazil. Here we extend the distribution of the species to tributaries of both the Upper and Middle río Uruguay basin throughout Misiones, Argentina (fig. 9). The Middle Uruguay and Lower Uruguay *sensu* Zaniboni Filho & Schulz (2003) correspond to the Lower Uruguay Freshwater ecoregion *sensu* Abell et al. (2008).

### Habitat

All the above listed specimens of *Gymnogeophagus lipokarenos* have been captured in environments with clear to slightly turbid water and flat slab-like exposed rocky bottoms or above gravel (figs. 5-6 and 8).






fig. 7. *Gymnogeophagus lipokarenos*, arroyo Melo. A (CI-FML 7264) and B: males (IBIGEO-I 450), C: female (IBIGEO-I 450)




fig. 8. Arroyo Melo, middle Uruguay basin

### Remarks

The specimens of *G. lipokarenos* were identified following Malabarba et al. (2015). This species is diagnosed by a bright yellow color background on the unpaired fins, and the presence of white dots aligned between rays with the distal border of dorsal fin and dorsal and ventral contours of caudal fin red with white stripes and dots. The contrasting bright yellow and red coloration in unpaired fins is well observed in the specimens analyzed herein (figs. 1-4 and 7). Contrary to the holotype's photo and the original description that states that this species possesses a "lateroventral color bluish white", most of the adult male specimens analyzed herein have a yellow ventral half of the body (figs. 1-3 and 7) instead of bluish white (fig. 4). Both color morphs (yellow or blue) are however found in our material and were even collected syntopically (e.g. at Salto Caracol, arroyo El Saltito). These two color morphs may represent different reproductive or territorial strategies as it is known from other cichlid species, as in *Astatotilapia burtoni*, for example (Korzán & Fernald, 2006). Further studies involving manipulatory experiments would be of much interest to understand if these color morphs are involved in behavioral aspects of this species, if these differences are plastic, if they are reversible or if these are inherited and not environmentally affected.

Malabarba et al. (2015) described *G. lipokarenos* as endemic to the Upper Uruguay Ecoregion (Abell et al., 2008). Our records from Misiones, Argentina, show that *G. lipokarenos* is distributed in both the Upper and the Lower Uruguay Ecoregions (Abell et al., 2008; Hales & Petry, 2015). The distribution of *G. lipokarenos* in Misiones (fig. 9) reaches all the way South to the latitude from where *G. constellatus* Malabarba et al., 2015 and *G. missioneiro* Malabarba et al., 2015 are known. These species however are only known from left bank tributaries of the Uruguay river in Brazil and have so far not been collected from the right bank affluents of the río Uruguay in Argentina.




fig. 9. Collecting sites of *Gymnogeophagus lipokarenos* in Argentina: yellow: arroyo Toro, red: arroyo Guerrero, green: arroyo Fortaleza, light blue: arroyo El Saltito, Salto Caracol; pink: arroyo El Saltito, Salto Golondrinas; brown: arroyo Paraiso; Brown: lower arroyo Paraiso; orange: Arroyo Melo, and blue: arroyo Shangai

### Acknowledgements

We thank the authorities of the Ministerio de Ecología y Recursos Naturales Renovables de la Provincia de Misiones for the awarded fishing permits. We thank D. Baldo, J.M. Ferro, J.M. Mirande, G. Aguilera, M. Benítez and G. Terán for field support in the collection of specimens at Arroyo Melo. Financial support was provided by Comisión de Investigaciones Científicas de la provincia de Buenos Aires (CIC), Facultad de Ciencias Naturales y Museo (UNLP), and Grant Agency of the Czech Republic (GAČR) (grant number 14-26060P) to ŠŘ. In addition, we also thank Google Earth for the map used in this contribution.

### References

- Abell, R., M.L. Thieme, C. Revenga, M. Bryer, M. Kottelat, N. Bogutskaya, B. Coad, N. Mandrak, S.C. Balderas, W. Bussing, M.L.J. Stiassny, P. Skelton, G.R. Allen, P. Unmack, A. Naseka, R. Ng, N. Sindorf, J. Robertson, E. Armijo, J.V. Higgins, T.J. Heibel, E. Wikramanayake, D. Olson, H.L. López, R.E. Reis, J.G. Lundberg, M.H. Sabaj Pérez & P. Petry (2008): Freshwater ecoregions of the World: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience* 58: 403-414
- Hales, J. & P. Petry (2015): Freshwater Ecoregions of the World. 333 Upper Uruguay. Last updated 2.Oct.2015 when accessed on 4.Jul.2017 at [www.feow.org/ecoregions/details/upper\\_uruguay](http://www.feow.org/ecoregions/details/upper_uruguay)
- Korzan, W.J. & R.D. Fernald (2006): Territorial male color predicts agonistic behavior of conspecifics in a color polymorphic species. *Behavioral Ecology* 18 (2): 318-323
- Loureiro, M., M. Zarucki, L.R. Malabarba & I. González-Bergonzoni (2016): A new species of *Gymnogeophagus* Miranda Ribeiro from Uruguay (Teleostei: Cichliformes). *Neotropical Ichthyology* 14 (1): 155-164

- Malabarba, L.R., M.C. Malabarba & R.E. Reis (2015): Descriptions of five new species of the Neotropical cichlid genus *Gymnogeophagus* Miranda Ribeiro, 1918 (Teleostei: Cichliformes) from the rio Uruguay drainage. *Neotropical Ichthyology* 13 (4): 637-662
- Reis, R.E. & L.R. Malabarba (1988): Revision of the Neotropical cichlid genus *Gymnogeophagus* Ribeiro, 1918, with descriptions of two new species (Pisces, Perciformes). *Revista Brasileira de Zoologia* 4: 259-305
- Zaniboni Filho, E. & U.H. Schulz (2003): Migratory fishes of the Uruguay River. *In*: Carolsfeld, J., Harvey, B., Ross, C. & Baer, A. (Eds.), *Migratory Fishes of South America: Biology, Fisheries and Conservation Status*. International Development Research Center, pp. 161-194.

recommended form for reference:

- Casciotta, J., A. Almirón, Š. Říčanová, K. Dragová, L. Piálek, F. Alonso & O. Říčan (2017): First record of *Gymnogeophagus lipokarenos* Malabarba, Malabarba & Reis, 2015 (Teleostei: Cichliformes) from Argentina. *Ichthyological Contributions of PecesCriollos* 54: 1-5  
available as pdf-file at [www.pecescrilloos.de](http://www.pecescrilloos.de) since 26.Jul.2017