



Novel host report for *Catadiscus uruguayensis* Freitas & Lent, 1939 (Trematoda, Diplodiscidae) infecting *Austrolebias* Costa, 1998 species from Uruguay

Renzo I. Vettorazzi¹, Walter A. Norbis¹, Sergio R. Martorelli²

1 Universidad de la República, Facultad de Ciencias, Laboratorio de Fisiología de la Reproducción y Ecología de Peces, Iguá 4225 Malvín Norte, Montevideo, P.C. 11400, Uruguay. **2** Centro de Estudios Parasitológicos y Vectores (CEPAVE-UNLP-CONICET), Boulevard 120 s/n e/60 y 64, C.P. 1900, La Plata, Buenos Aires, Argentina.

Corresponding author: Renzo Vettorazzi, rvetto@fcien.edu.uy

Abstract

The genus *Catadiscus* Cohn, 1904 has a total of 16 known species that infect the intestinal tract of reptiles, amphibians, and mollusks. However, *Catadiscus* has never been found in teleosts. The annual fish *Austrolebias prognathus* (Amato, 1986) and *A. cheradophilus* (Vaz-Ferreira, Sierra de Soriano & Scaglia de Paulete, 1965) were collected from temporary ponds in the southeast of Uruguay. The specimens found in the intestinal tract of these hosts were morphologically identified as *Catadiscus uruguayensis* Freitas & Lent, 1939, which until now were only known to infect amphibians. This work represents the first report of the genus *Catadiscus* infecting and developing in a fish host.

Keywords

Annual fish, new host, parasite, South America, temporary pond.

Academic editor: Moisés Gallas | Received 22 June 2020 | Accepted 17 September 2020 | Published 29 September 2020

Citation: Vettorazzi RI, Norbis WA, Martorelli SR (2020) Novel host report for *Catadiscus uruguayensis* Freitas & Lent, 1939 (Trematoda, Diplodiscidae) infecting *Austrolebias* Costa, 1998 species from Uruguay. Check List 16 (5): 1277–1282. <https://doi.org/10.15560/16.5.1277>

Introduction

In South America, the genus *Catadiscus* Cohn, 1904 has a total of 16 known species which infect the intestinal tract of reptiles, amphibians and mollusks (two species from reptiles, 10 from amphibians, three known to infect both, and one reported in a mollusk) (Freitas 1943; Lent et al. 1946; Freitas and Dobbin 1956; Mañé-Garzón and Gortari 1965; Suriano 1970; Ostrowski 1978; Corrêa and Artigas 1979; Incorvaia 1983; Hamann 1992; Hamann et al. 2014). Particularly in Uruguay, three species have been reported: one from reptiles (*Catadiscus dolichocotyle* Cohn, 1913 in *Liophis miliaris* (Linnaeus, 1758) and *Chironius fuscus* (Linnaeus, 1785)), and two from

amphibians (*Catadiscus corderoi* Mañé-Garzón, 1958 in *Pseudis meridionalis* Miranda-Ribeiro, 1926 and *Pseudis minutus* Günther, 1858; and *Catadiscus uruguayensis* Freitas & Lent, 1939 in *Leptodactylus latrans* (Steffen, 1815)) (Freitas and Lent 1939; Mañé-Garzón and Gortari 1965). These hosts, specifically the amphibian ones, can be found in habitats associated with freshwater ecosystems, such as wetlands and temporary ponds, where tadpoles cohabit with teleost known as annual fishes.

Annual fishes (Cyprinodontiformes, Aplocheiloidei) stand as an attractive model worldwide for several topics including evolution and genetics, as well as being

bioindicators and pest regulators (Fletcher et al. 1992; Frenkel and Goren 2000; Arenzon et al. 2003; Berois et al. 2015). They exhibit unique annual life-cycles that comprise a rainy season during which they mature and reproduce, and a dry season where their habitats (temporary ponds) become depleted of water and thus only their burrowed draught-resistant eggs remain until the next rainy season in the following year, which triggers the completion of the eggs' development (Berois et al. 2015). In Uruguay, the annual fishes that are most widely used for research are those of the genus *Austrolebias* Costa, 1998, yet their parasitic fauna is extensively unknown, with only four parasitological reports in the South American continent (Luque et al. 2011; Delgado and García 2015; Montes et al. 2017; Marcotegui et al. 2018). Aiming to expand the knowledge of parasites infecting *Austrolebias*, we report the presence of a *Catadiscus* species in two annual fishes from Uruguay.

Methods

Fish hosts were collected manually from temporary ponds in Uruguay during the rainy season of 2018 and 2019 in the Department of Rocha (33°41'26"S, 054°41'15"W) and in the Department of Lavalleja (34°11'55"S, 053°55'45"W), Uruguay (Fig. 1).

Hosts were captured, manipulated, and euthanized under the approval of the institutional Ethics Council of Animal Experimentation (CHEA, Facultad de Ciencias, Uruguay), which follows the American Veterinary Medical Association (AVMA) international guidelines for fish euthanasia (with eugenol overdose before a cervical and cephalic incision) (Leary et al. 2013). Parasites were removed from the intestine of fish, heat-killed with slight pressure under coverslip, fixed in 10% formalin and then preserved in 70% ethanol. They were then stained with Langeron's Carmine, dehydrated in ethanol series, clarified with eugenol, and mounted in *Entellan*® medium. Whole-mount specimens were observed using an Olympus BX50 microscope and digital images were taken with adapted 318CU 3.2M CMOS camera and Micro-metrics SE Premium program. Illustration was done with digital tablet Genius i608X using the free license program GIMP2. The specimens were identified using the taxonomic keys proposed by Sey (1991) and Jones et al. (2005). Followingly, the description hereto was compared with the original description of the congeneric species. Distribution map was made with the program QGIS version 3.4 (QGIS Development Team 2018) and the program SimpleMappr (Shorthouse 2010). Specimens are deposited in the collection of Invertebrate Zoology of the Natural History National Museum (Museo Nacional de




Figure 1. Known records of *Catadiscus* Cohn, 1904 in South America and the new records in Uruguay; South American reports of *Catadiscus* (red dots). On the right, a closer look to Uruguay showing the location of the temporary ponds sampled in this study (black star—Department of Rocha, 33°41'26"S, 054°41'15"W; white star—Department of Lavalleja 34°11'55"S, 053°55'45"W).

Historia Natural, MNHN), Montevideo, Uruguay.

Measurements are reported in micrometers, with the format: mean (standard deviation; minimum-maximum). Due to the small sample size, standard deviation was calculated with bootstrap (9999 repetitions) using the free software PAST4 (Hammer et al. 2001). For this same reason, no confidence intervals are reported in the parasitological indexes. The latter were calculated according to Bush et al. (1997).

Results

Superfamily Paramphistomoidea

Family Diplodiscidae

Genus *Catadiscus* Cohn, 1904

Catadiscus uruguayensis Freitas & Lent, 1939

Figure 2

Materials examined. URUGUAY • 1 specimen; Rocha Department, Ruta Vieja; 33°41'26"S, 054°41'15"W; 14 Sep. 2018; R. Vettorazzi leg.; temporary pond, manual collection, host: *Austrolebias cheradophilus* (Vaz-Ferreira, Sierra de Soriano & Scaglia de Paulete, 1965); MNHN4215. • 1 specimen; Lavalleja Department, Retamosa; 34°11'55"S, 053°55'45"W; same data as for preceding, except host: *A. prognathus* (Amato, 1986); MNHN4216. • 3 specimens; same data as for preceding, except 27 Aug. 2019; MNHN4216.

Identification. Description based on three specimens stained and mounted.

Body sub-pyriform with smooth tegument, anterior surface region pigmented irregularly. Total length 1277 (218; 1100–1520), width at the equatorial level 405 (31; 370–430). Oral opening terminal, with undulated edges, diameter 133 (15; 120–150). Acetabulum sub-terminal, with medial constriction present but not well developed; 386 (23; 350–393) long, 323 (13; 310–335) wide. Pharynx strongly muscular, with extramural sacs, 211 (13; 200–225) long, 169 (4; 165–173) wide. Esophagus slim, 97 (15; 80–110) long; esophageal bulb present, 103 (20; 80–115) long, 90 (17; 70–100) wide. Intestinal ceca short, reaching equatorial level of the body, 301 (57; 250–405) long. Genital pore opens at the level of intestinal bifurcation. Cirrus sac globular, 141 (8; 134–150) long, 37 (8; 30–45) wide; distal extremity displaying seminal vesicle with sperm reservoir in living specimens. Vitelline follicles lateral, touching and surpassing posterior end of intestinal ceca. Testicle intercecal, oval; 112 (10; 100–120) long, 56 (21; 33–75) wide. Ovary post-testicular, diagonally opposite to testicle, oval; 108 (16; 90–120) long, 77 (13; 65–90) wide. Mehlis' glands visible at posterior margin of ovary. Uterus complex, forming coils which occupy entire region of body after caecal bifurcation, passing between testis and ovary. Eggs large, operculate, collapsed in fixed specimens; 81 (4; 78–86) long, 43 (2; 41–45) wide. Prevalence of 33% in *Austrolebias prognathus* ($n = 6$) and 3% in *Austrolebias cheradophilus* (n




Figure 2. *Catadiscus uruguayensis* Freitas & Lent, 1939 collected from fish host; *C. uruguayensis* from first intestinal portion of *Austrolebias prognathus* (Amato, 1986) and *A. cheradophilus* (Vaz-Ferreira, Sierra, Soriano, Scaglia & Paulete, 1965) captured in temporary ponds from Southwest Atlantic basin, Southeast Uruguay. Scale bar: 300 μ m.

= 30). Mean Intensity of 2 in *A. prognathus* and 1 in *A. cheradophilus*.

Remarks. Given the previous description, the morphological features place the species reported now under the genus *Catadiscus* due to the lumen of the acetabulum being constricted into anterior and posterior regions, lacking a central accessory sucker or peduncle.

The specimens here described were most comparable to four of the 16 species described and currently valid for the genus, given important characters such as overall body shape, undulated edges of the oral opening, presence of extramural sacs in pharynx, presence of esophageal bulb, acetabula morphology, vitelline follicles distribution, relative position of gonads and the uterus, position of the gonopore, and egg size (Sey 1991; Jones et al. 2005). These four species are *C. uruguayensis*, *C. rochai* Correa & Artigas, 1978, *C. hylae* Incorvaia, 1983, and *C. pomaceae* Hamann, 1992 (Table 1). All the cited species are known to infect either amphibians or reptiles from Argentina, Brazil, and Uruguay (except *C. pomaceae*, which was found in the snail *Pomacea canaliculata* (Lamarck, 1801)) (Sey 1991).

In contrast to the specimens of the present work, *C. pomaceae* was described as having the uterus coils reaching between the oral opening and the caecal bifurcation, bigger pharynx, and somewhat different acetabula to body length ratio (Hamann 1992). *Catadiscus hylae* differs by not presenting undulated edges of the oral opening and having fewer vitelline follicles which do not surpass the posterior end of the intestinal ceca (Incorvaia 1983). *Catadiscus rochai* is different from the specimens of the present work, in not having uterine coils between testis and ovary (Corrêa and Artigas 1979; Sey 1991). Thus, the specimens here described shared the most similarities with *C. uruguayensis* (Sey 1991) regardless of

some variations such as vitelline follicle size, egg size, and testis size (Table 1). These differences whatsoever, have all been recorded in the literature.

Suriano (1970) and Ostrowski (1978) redescribed *C. uruguayensis*, offering insight on disagreements with the original description from Freitas and Lent (1939). For instance, Suriano (1970) and Ostrowski (1978) established the vitelline follicles as being similar in size to the eggs, as opposed to Freitas and Lent (1939) who regarded the vitelline follicles to be remarkably smaller than the eggs. Both Suriano (1970) and Ostrowski (1978) also accepted egg sizes inferior to 0.1 mm as still pertaining to the species *C. uruguayensis*, while Freitas and Lent (1939) did not. Regarding the genitalia, Ostrowski (1978) mentioned that the testis and ovary had similar size in *C. uruguayensis* collected from tadpoles of *Hyla pulchella* (Duméril & Bibron, 1841), yet the ones found in adult frogs presented considerably bigger testis. In the specimens found in this work the testis and the ovary had similar size. A possible explanation is the testicular atrophy reported in *Catadiscus*, which occurs when the uterus is filled with eggs (Freitas and Lent 1939; Ostrowski 1978).

This work represents the first report of the genus *Catadiscus* infecting and developing (adult, gravid parasites, with food content in their ceca) in a fish host.

Catadiscus uruguayensis has been reported in Uruguay by Freitas and Lent (1939) parasitizing *Leptodactylus latrans*. This definitive amphibian host is associated throughout its life history with the same type of ecosystem where *A. prognathus* and *A. cheradophilus* populations occur (Naya et al. 2003; Lavilla et al. 2010; Berois et

al. 2015). Therefore, *A. prognathus* and *A. cheradophilus* could have been infected through ingestion of the tadpole of the definitive amphibian host (both *Austrolebias* species are reportedly piscivorous, thus capable of feeding on frog larvae; Costa 2009). However, this scenario would likely result in a case of concomitant predation (Johnson et al. 2010; Thieltges et al. 2013). Another possibility is infection by ingesting the metacercaria from the vegetation or substrate where they became encysted. It has not been studied if these fish have herbivore habits as juveniles, but it has been observed by Ostrowski (1978) that the substrate where the cercaria encysts is not specific, meaning they could encyst on the exterior of another prey such as an arthropod or a mollusk, which do form part of the diet of these species of *Austrolebias* (Laufer et al. 2009).

The mature specimens of *C. uruguayensis* found in this work are indicative of their ability to develop and fulfill their life cycle in freshwater fish hosts. This means that, despite the likelihood of the infection being accidental, this finding corroborates the generalist strategy of *Catadiscus*.

Acknowledgements

We thank the Agencia Nacional de Investigación e Innovación (ANII) and Programa de Desarrollo de las Ciencias Básicas (PEDECIBA), Uruguayan programs which finance the Master in Biological Science postgraduate studies of RV, for funding field sampling and laboratory materials. We are also thankful to Dr Uriel Koziol for

Table 1. Comparison of *Catadiscus uruguayensis* Freitas & Lent, 1939 from the present work with the four most similar South American *Catadiscus* species. Data expressed in micrometers, in the form “mean ± standard deviation (minimum-maximum)”, “minimum-maximum”, or “mean”, depending on the available data. AC: acetabula; L: length; TL: total length; W: width.

References	<i>C. uruguayensis</i>		<i>C. hylae</i>	<i>C. pomaceae</i>	<i>C. rochai</i>
	Present work	Suriano (1970); Ostrowski (1978)	Incorvaia (1983)	Hamann (1992)	Correa and Artigas (1979)
Total length	1277 ± 218 (1100–1520)	990–2100	1380–3150	700–2700	1850
Medial width	405 ± 31 (370–430)	480–880	570–1560	250–1080	750
Oral opening	133 ± 15 (120–150)	100–190	80–100	110–210	90
Extramural sacs	L	211 ± 13 (200–225)	190–390	80–140	180
	W	169 ± 4 (165–173)	130–150	110–220	270
Esophageal bulb	97 ± 15 (80–110)	210–240	90–170	250–470	380
Pharynx	L	103 ± 20 (80–115)	72–160	110–150	190–280
	W	90 ± 17 (70–100)	75–140	50–90	130–170
Acetabula	L	368 ± 23 (350–393)	460–670	500–900	200–560
	W	323 ± 13 (310–335)	390–540	400–900	160–460
Testis	L	112 ± 10 (100–120)	140–280	—	170
	W	56 ± 21 (33–75)	150–280	—	220
Ovary	L	108 ± 16 (90–120)	65–160	—	100
	W	77 ± 13 (65–90)	78–130	—	70
Cirrus sac	L	141 ± 8 (134–150)	130	130–170	—
	W	37 ± 8 (30–45)	20	100	—
Eggs	L	81 ± 4 (78–86)	72–100	54–87	86–90
	W	43 ± 2 (41–45)	42–50	18–39	47–54
TL:AC	3.1–3.9	2.1–3.2	3.5	3.5–6.5	3.8
AC:pharynx	3.4–4.4	3.5–4.6	5.0	1.5–2.7	5.4
Locality	Uruguay	Argentina, Brazil, Uruguay	Argentina	Argentina	Brazil
Host type	Fish	Amphibian	Amphibian	Mollusk	Ophidian

kindly donating a bibliographic collection about *Catadiscus*, and Dr Marcelo Loureiro for aiding in sampling the hosts used in this study. We thank the constructive criticism and contributions of the editor and three anonymous referees.

Authors' Contributions

RV collected, processed, and described the material, was the writer of the first draft, adapted it to the format of the journal, and made the major corrections throughout the peer-review process. WN coordinated and made the resources available. SM confirmed the righteousness of the identification. WN and SM worked as overseers of the research process and contributed to the correction of the draft.

References

- Arenzon A, Pinto RF, Colombo P, Raya-Rodriguez MT (2003) Assessment of the freshwater annual fish *Cynopoeilus melanotaenia* as a toxicity test organism using three reference substances. *Environmental Toxicology and Chemistry* 22 (9): 2188–2190. <https://doi.org/10.1897/02-292>
- Berois N, García G, de Sá RO (2015) Annual fishes: life history strategy, diversity, and evolution. CRC Press, New York, 327 pp.
- Bush AO, Lafferty KD, Lotz JM, Shostak AW (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *The Journal of Parasitology* 83 (4): 575–583. <https://doi.org/10.2307/3284227>
- Costa WJ (2009) Trophic radiation in the South American annual killifish genus *Austrolebias* (Cyprinodontiformes: Rivulidae). *Ichthyological Exploration of Freshwaters* 20 (2): 179–191.
- Corrêa AAS, Artigas PT (1979) *Catadiscus rochai* n. sp. (Trematoda; Paramphistomidae), parasito de *Dromicus typhlus* (L.) (Ophidia; Colubridae). *Memórias do Instituto Butantan* 42/43: 145–149.
- Delgado C, García G (2015) Coevolution between *Contracecum* (Nematoda, Anisakidae) and *Austrolebias* (Cyprinodontiformes, Rivulidae) host-parasite complex from SW Atlantic coastal basins. *Parasitology Research* 114 (3): 913–927. <https://doi.org/10.1007/s00436-014-4257-2>
- Fletcher M, Teklehaimanot A, Yemane G (1992) Control of mosquito larvae in the port city of Assab by an indigenous larvivorous fish, *Aphanius dispar*. *Acta Tropica* 52 (2–3): 155–166. [https://doi.org/10.1016/0001-706x\(92\)90032-s](https://doi.org/10.1016/0001-706x(92)90032-s)
- Freitas JFT (1943) *Catadiscus mirandai* n. sp., parasito de *Hemipipa carvalhoi* Mir-Rib. *Revista Brasileira de Biologia* 3 (4): 411–412.
- Freitas JFT, Dobbin JE (1956) Novo parasito de rã: *Catadiscus propinquus* sp. n. (Trematoda, Paramphistomoidea). *Revista Brasileira de Biologia* 16 (4): 439–441.
- Freitas JFT, Lent H (1939) Revisão do gênero *Catadiscus* Cohn 1904 (Trematoda, Paramphistomoidea). *Boletim Biológico* 4: 305–315.
- Frenkel V, Goren M (2000) Factors affecting growth of killifish, *Aphanius dispar*, a potential biological control of mosquitoes. *Aquaculture* 184 (3–4): 255–265. [https://doi.org/10.1016/S0044-8486\(99\)00326-9](https://doi.org/10.1016/S0044-8486(99)00326-9)
- Hamann MI (1992) *Catadiscus pomaceae* sp. n. (Trematoda, Paramphistomatidae) from *Pomacea canaliculata* (Lamarck, 1801) (Prosobranchia, Ampullariidae). *Memórias do Instituto Oswaldo Cruz* 87 (1): 9–14. <https://doi.org/10.1590/s0074-027619920001002>
- Hamann MI, Kehr AI, González CE (2014) Helminth community structure in the Argentinean bufonid *Melanophryniscus klappenbergi*: importance of habitat use and season. *Parasitology Research* 113: 3639–3649. <https://doi.org/10.1007/s00436-014-4029-z>
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4 (1): 1–9.
- Incorvaia IS (1983) *Catadiscus hylae* sp. nov. (Trematoda: Paramphistomidae) parasito intestinal de *Hyla pulchella* Dumeril y Bibron, 1841 (Anura; Hylidae) de la provincia de Buenos Aires, República Argentina. *Neotropica* 29: 91–95.
- Johnson PTJ, Dobson A, Lafferty KD, Marcogliese DJ, Memmott J, Orlofske SA, Poulin R, Thielges DW (2010) When parasites become prey: ecological and epidemiological significance of eating parasites. *Trends in Ecology & Evolution* 25 (6): 362–371. <https://doi.org/10.1016/j.tree.2010.01.005>
- Jones A, Bray RA, Gibson DI (2005) Keys to the Trematoda. Volume 2. CABI Publishing, Wallingford, 768 pp.
- Laufer G, Arim M, Loureiro M, Piñeiro-Guerra JM, Clavijo-Baquet S, Fagúndez C (2009) Diet of four annual killifishes: an intra and interspecific comparison. *Neotropical Ichthyology* 7 (1): 77–86. <https://doi.org/10.1590/S1679-62252009000100010>
- Lavilla EO, Langone JA, Caramaschi U, Heyer WR, de Sa RO (2010) The identification of *Rana ocellata* Linnaeus, 1758. Nomenclatural impact on the species currently known as *Leptodactylus ocellatus* (Leptodactylidae) and *Osteopilus brunneus* (Gosse, 1851) (Hylidae). *Zootaxa* 2346 (1): 1–16.
- Leary S, Underwood W, Raymond AS, Corey CD, Grandin T, Greenacre C, Gwaltney-Brant S, McCrackin MA, Meyer R, Miller D (2013) <https://www.avma.org/KB/Policies/Documents/euthanasia.pdf>. Accessed on 2020-9-22.
- Lent H, Freitas JFT, Proença MC (1946) Alguns helmintos de batráquios colecionados no Paraguai. *Memórias do Instituto Oswaldo Cruz* 44 (1): 195–214. <https://doi.org/10.1590/S0074-02761946000100007>
- Luque JL, Aguiar JC, Vieira FM, Gibson DI, Santos CP (2011) Checklist of Nematoda associated with the fishes of Brazil. *Zootaxa* 3082 (1): 1–88. <https://doi.org/10.11646/zootaxa.3082.1.1>
- Mañe-Garzón F, Gortari AM (1965) Sobre algunos trematodos de ofidios del Uruguay. *Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo* 8 (107): 1–21.
- Marcotegui PS, Montes MM, Barneche J, Ferrari W, Martorelli S (2018) Geometric morphometric on a new species of Trichodinidae. A tool to discriminate trichodinid species combined with traditional morphology and molecular analysis. *International Journal for Parasitology: Parasites and Wildlife* 7 (2): 228–236. <https://doi.org/10.1016/j.ijppaw.2018.06.004>
- Montes MM, Barneche J, García I, Preisz S, Martorelli SR (2017) New data on the acanthocephalan *Neoechinorhynchus villoldoi* Vizcaino, 1992 (Neoechinorhynchidae: Acanthocephala), based on specimens found in *Austrolebias bellottii* (Steindachner, 1881) (Rivulidae: Cyprinodontiformes) from Punta Indio, Argentina. *Check List* 13 (4): 53–59. <https://doi.org/10.15560/13.4.53>
- Naya D, Maneyro R, Camargo A, Canavero A, da Rosa I (2003) Seasonal changes in gut length of the South American common frog *Leptodactylus ocellatus* (Amphibia, Anura). *Biociências* 11 (1): 47–52.
- Ostrowski M (1978) Fauna de agua dulce de la República Argentina. IX. Sobre representantes de la familia Paramphistomatidae (Trematoda). *Physis* B 38 (95): 55–62.
- QGIS Development Team (2018) QGIS Geographic Information System version 3.4. Open Source Geospatial Foundation Project. <https://www.qgis.org/en/site/index.html>. Accessed on: 2020-9-22.
- Sey O (1991) Handbook of the zoology of amphistomes. CRC press, New York, 480 pp.
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net>. Accessed on: 2020-9-22.
- Suriano DM (1970) Estudio sobre la fauna parasitaria de *Leptodactylus ocellatus*.

tylus ocellatus (L.) (Amphibia, Leptodactylidae) de la República Argentina. I. Trematodes. Revista Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Zoología 10 (15): 215–239.
Thieltges DW, Amundsen PA, Hechinger RF, Johnson PTJ, Lafferty

KD, Mouritsen KN, Preston DL, Reise K, Zander CD, Poulin R (2013) Parasites as prey in aquatic food webs: implications for predator infection and parasite transmission. *Oikos* 122 (10): 1473–1482. <https://doi.org/10.1111/j.1600-0706.2013.00243.x>