1

2

ENVIRONMENTAL POLLUTANT HEXACHLOROBENZENE INDUCES HYPERTENSION IN A RAT MODEL

3	Castilla, Rocío*; Asuje, Agustín [†] ; Rivière, Stéphanie*; Romero Caimi, Giselle [‡] ; Martín,						
4	Pedro [†] ; Cao, Gabriel [*] ; Kleiman de Pisarev, Diana [‡] ; Milesi, Verónica [†] and Alvarez, Laura [‡] .						
5	*Universidad de Buenos Aires. CONICET. Instituto de Investigaciones Cardiológicas						
6	(ININCA). C1122AAJ Buenos Aires, Argentina. rcastillalozano@fmed.uba.ar,						
7	s.riviere@conicet.gov.ar; gabrielcao@fibertel.com.ar						
8	[†] Universidad Nacional de La Plata. Facultad de Ciencias Exactas. CONICET. Instituto de						
9	Estudios Inmunológicos y Fisiopatológicos (IIFP). 1900, La Plata, Buenos Aires, Argentina.						
10	agustin.asuaje@gmail.com; pedromartinv@gmail.com; vmilesi49@gmail.com						
11	[‡] Universidad de Buenos Aires. Facultad de Medicina. Departamento de Bioquímica Humana.						
12	Laboratorio de Efectos Biológicos de Contaminantes Ambientales. C1121ABG Buenos Aires						
13	Argentina. gisellerc@gmail.com; dianakleiman@yahoo.com.ar; laura6alvarez@yahoo.com.ar						
14	Running title: HCB INDUCES HYPERTENSION IN A RAT MODEL						
15	* Correspondence to: Rocío Castilla, PhD.						
16	Instituto de Investigaciones Cardiológicas ININCA-UBA-CONICET						
17	Marcelo T. de Alvear 2270- C1122AAJ, Buenos Aires, Argentina						
18	Tel: +54 +11 4508-3888/36						
19	rcastillalozano@fmed.uba.ar						
20	1						

Abbreviations

AhR: aromatic hydrocarbon receptorAch: AcetylcholineAT₁: angiotensin II type1 receptorBP: Systolic blood pressureDII: deiodinase IIeNOS: endothelial nitric oxidide synthaseERα: estrogen receptor alphaHCB: HexachlorobenzeneNA: noradrenalinNO: nitric oxidideTCDD: 2,3,7,8-tetrachlorodibenzo-p-dioxinTH: thyroid hormonesVSMC: vascular smooth muscle cells

22 ABSTRACT

23 Hexachlorobenzene (HCB) is a dioxin-like environmental pollutant, widely distributed in the environment. New research links exposure to high levels of persistent organic environmental 24 toxicants to cardiovascular disease, however little is known about the effect of HCB on 25 vascular function and on blood pressure. The purpose of the present study was to evaluate 26 biochemical and cardiovascular changes resulting from subchronic HCB exposure. Adult 27 female Sprague-Dawley rats were treated with vehicle or HCB (5 or 500 mg/kg b.w) for 45 28 days. Systolic blood pressure (BP), recorded by tail cuff plethysmography, was significantly 29 increased at 35, 40 and 45 days of 500 mg/kg HCB-treatment. HCB (500 mg/kg) increased 30 arterial thickness, while both 5 and 500 mg/kg HCB decreased proliferating cell nuclear 31 antigen (PCNA) protein levels and cellular nuclei in abdominal aortas indicating a 32 hypertrophic process. Also, aortas from both groups of HCB-treated rats presented higher 33 sensitivity to noradrenalin (NA) and a significant decrease in maximum contractile response. 34 Arteries from 500 mg/kg HCB-treated rats showed a significant increase in the levels of 35 transforming growth factor- β 1 (TGF- β 1) mRNA and angiotensin II type1 receptor (AT₁), and 36 a significant decrease in estrogen receptor alpha (ER α), endothelial nitric oxidide synthase 37 (eNOS) protein expression and deiodinase II (DII) mRNA levels. In conclusion, we have 38 demonstrated for the first time that subchronic HCB administration significantly increases BP 39 and alters associated cardiovascular parameters in rats. In addition, HCB alters the expression 40 of key vascular tissue molecules involved in BP regulation, such as TGF-β1, AT1, ERα, 41 eNOS and DII. 42

43

44 **Keywords**: hexachlorobenzene, hypertension, aorta, TGF- β 1, ER α , deiodinase II.

46 1. INTRODUCTION

Hypertension occurs in one out of three adults in urban populations and represents a
significant risk factor for life-threatening cardiovascular disease (Mills *et al.* 2016).
Nevertheless, and despite intensive research, only around 5% of cases have an identifiable
cause (Carretero and Oparil, 2000). The etiology of hypertension is very heterogeneous and
could involve exposure to environmental pollution as a risk factor (EPA, Environmental
Protection Agency USA, 2002; Huang *et al.*, 2006).

Dioxins are among the most harmful environmental toxins for humans. The Public Health Statement for Dioxins (1998), reported that 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) increases blood pressure (BP) and induces cardiac hypertrophy in laboratory animals (Kopf et al., 2008).

Hexachlorobenzene (HCB) is a widely distributed dioxin-like environmental pollutant. It has been used as a fungicide, and it is still released into the environment as a by-product in several industrial processes. It is very persistent and bioaccumulative, and belongs to the group of persistent organic pollutants (Xu et al., 2013). Chronic exposure of laboratory animals to HCB elicits a number of toxic effects including neurological symptoms, immunological disorders and endocrine disruption (ATSDR, 2002). However, its impact on the cardiovascular system has not yet been evaluated.

It has been reported that HCB alters the expression of molecules involved in BP regulation in several tissues, e.g. it affects the levels of growth factors, interleukins and cytokines such as transforming growth factor- β 1 (TGF- β 1) (Giribaldi *et al.*, 2011), which, in particular, has been associated to hypertension (Derhaschnig *et al.*, 2002). This growth factor also alters estrogen receptor alpha (ER α) levels in several types of cells (Li *et al.*, 2008; Peña *et al.*, 2012). This receptor is in turn involved in BP regulation, increasing the production of molecules like nitric oxidide (NO) with a relaxing action on vascular smooth muscle cells
(VSMC) (Jobe *et al.*, 2013; Pedram *et al.*, 2002).

BP can be altered by an imbalance between relaxation and contraction processes. In this sense, TGF- β 1 has been also reported to regulate the expression of angiotensin II type1 receptor (AT₁), the receptor by which angiotensin II acts as vasoconstrictor (Chaudhary and Chaudhary, 2017).

In addition, HCB is known to affect thyroid hormones (TH) homeostasis. In particular, modifications have been reported in the activity and expression of hepatic deiodinase DI and DII enzymes, respectively (Alvarez *et al.*, 2005; Kleiman de Pisarev *et al.*, 1990). TH have multiple effects on the cardiovascular system, including alterations in cardiovascular hemodynamics and the relaxation of coronary artery vascular smooth muscle cells (VSMC) (Gomberg-Maitland and Frishman, 1998).

⁸² Despite the evidence discussed above in different tissues, little is known about HCB effect on ⁸³ the expression of TGF- β 1, ER α , endothelial NO synthase (eNOS), AT1 and DII in the ⁸⁴ vascular system. In this context, the aim of this study was to determine whether subchronic ⁸⁵ exposure of laboratory animals to HCB alters BP, focusing on its possible effects on vascular ⁸⁶ hemodynamics and the expression of key molecules involved in BP regulation.

87 2. MATERIALS AND METHODS

88 2.1. Chemicals

HCB (>99% purity) was purchased from Sigma-Aldrich Co. The reagents used for cDNA
synthesis, reverse transcriptase, molecular weight markers, random primers, desoxi-nucleotide
triphosphate (dNTPs), and Taq enzyme polymerase, were purchased from Biodynamics SRL
(Buenos Aires, Argentina). Specific primers for TGF-β1 and glyceraldehyde 3-phosphate

93 dehydrogenase (GPDH) were purchased from Invitrogen Life Technology (Carlsbad, CA).
94 Polyvinylidene difluoride membranes (PVDF), goat anti-mouse IgG, and anti-rabbit IgG were
95 purchased from Bio-Rad Laboratories Inc. CP-BU plates were purchased from Agfa,
96 (Gevaert, Argentina S.A). Anti-β-actin immunoglobulin and anti-proliferating cell nuclear
97 antigen (PCNA) from Sigma-Aldrich Co., monoclonal anti-TGF-β1, anti-AT1 and anti-eNOS
98 from Abcam Inc.. All reagents present purification degree of 96% and molecular biology
99 grade.

100

101 **2.2.** Animals

All procedures involving animals were approved by the Institutional Committee of Animal
Care and Use, at the University of Buenos Aires (CICUAL, School of Medicine) and
conducted according to the principles of the Guide for the Care and Use of Laboratory
Animals (NIH Publications No. 80-23, revised 1996).

Sprague-Dawley rats were maintained on a 12 h/12 h light/dark cycle in a temperature- $(21 \pm 2 \text{ °C})$ and humidity-controlled (65 ± 5%) environment. Animals had access to food (Purina chow) and tap water *ad libitum*.

109

110 **2.3.** Animals and treatment

Female rats (160 g at the onset of the experiment) were gavage-administered HCB (5 or 500 mg/kg) as a suspension in water, containing Tween 20 (0.5 ml/100 ml), 5 days a week for 45 days. Control animals received equal volumes of vehicle by the same route. Six rats per group were used.

117 To reduce ambient variability in BP measurement, the animals were acclimatized to handling118 to reduce stress.

Systolic BP was measured between 11 and 12 h a.m., in rats treated with vehicle or HCB for 30, 35, 40, and 45 days, by anaesthesia-free tail cuff plethysmography. The average of at least 3 readings per session was recorded. A pneumatic pulse transducer positioned on the ventral surface of the tail, distal to the occlusion cuff, detected the return of the pulse following a slow deflation of the cuff using a programmed electro-sphygmomanometer PE-300 (Narco Bio-Systems, Austin, Texas). Pulses were recorded on a Physiograph MK-IIIS (Narco Bio-Systems, Austin, Texas).

126

127 **2.5.** Sample handling

Rats were sacrificed under anesthesia with an i.p. solution of sodium pentobarbital and sodium diphenylhydantoin (Euthanyl®) (100 mg/kg b.w.). Blood samples were obtained by ventricular puncture and plasma was separated. The aortic tissue was removed for further processing.

132

133 **2.6. Western blotting**

Total cellular protein lysates were electrophoresed in 10–12% SDS-polyacrylamide gel (SDSPAGE), and then transferred to PVDF in a semidry transfer cell at 18 V for 1.5 h. Membranes
were blocked for 1 h at 4 °C with 5% non-fat dry milk – 2.5% bovine serum albumin (BSA)
in TBST buffer (10 mM Tris–HCl, pH 8.0, 0.5% Tween 20, 150 mM NaCl). Membranes were

incubated overnight at 4 °C with the specific primary antibody (1:500 anti-β-actin 138 immunoglobulin (Sigma-Aldrich Co), 1:500 anti-proliferating cell nuclear antigen (PCNA) 139 (Sigma-Aldrich Co.), 1:500 monoclonal anti-TGF-B1 (Abcam Inc), 1:250 anti-AT1 (Abcam 140 141 Inc) and 1:500 anti-eNOS (Abcam Inc)) and then washed five times with TBST, and the suitable peroxidase-conjugated anti-specific antibodies were used for protein detection. After 142 washing, blots were developed using and ECL detection kit (Amersham Biosciences, Inc., 143 UK). The quantitative analysis of the integrated optical density (IOD) of bands was done 144 using the Image Quant 5.2. Inc. software. 145

146

147 **2.7. Semiquantitative RT-PCR**

Total RNA from arteries was extracted using Trizol reagent following the manufacturer's instructions (Life Technologies, Inc.-BRL, Grand Island, NY). The reverse transcription and PCR analyses were made using 2 or 4 μ g of total RNA. The cDNAs generated were further amplified by PCR under optimized conditions using the primer pairs listed below.

152 **TGF-**β1:

- 153 Forward: 5'-CTGCTGGCAATAGCTTCCTA-3'
- 154 Reverse: 5'-CGAGCCTTAGTTGGACAGGAT-3'
- 155 **D II:**
- 156 Forward: 5'-ACTCGGTCATTCTGCTCAAG-3'
- 157 Reverse: 5'-TTCAAAGGCTACCCCATAAG-3'
- 158 **GPDH** (glyceraldehyde 3-phosphate dehydrogenase):
- 159 Forward: 5'-ACCCAG AAG ACT GTG GAT GG-3'
- 160 Reverse: 5'-CAC ATT GGG GGT AGG AAC AC-3

161 The number of cycles used was optimized for each mRNA to fall within the linear range of162 PCR amplification.

PCR products were resolved on a 1.5% (wt/vol.) agarose gel. The gel images were acquired with the GelPro analyzer (IPS, North Reading, MA). The levels of mRNA were quantified using a computer-assisted image analyzer (ImageQuant 5.2) and the PCR results for each sample were normalized with GPDH mRNA as an internal control.

167

168 **2.8. Thyroid hormone assay**

169 Serum concentration of thyroid hormones were analyzed by RIA as previously described170 (Obregon *et al.*, 1981).

171

172 2.9. Vessel preparation and isometric tension recording

The abdominal aortas were carefully dissected, free of connective tissue, immersed in Krebs 173 solution (KS: 130 mM NaCl, 4.7 mM KCl, 1.17 mM Na₂HPO₄, 1.16 mM MgSO₄, 24.0 mM 174 HCO₃Na, 2.5 mM CaCl₂, and 6 mM glucose), and cut into 3-4 mm wide rings. The ring 175 vessel was gently suspended between two stainless steel wires in a water-jacketed organ bath 176 kept at 37 °C and filled with a KS, bubbled with a mixture of 5% CO₂ and 95% O₂, pH 7.4. 177 The lower wire was fixed to a vertical plastic rod immersed in the organ bath, while the upper 178 one was rigidly attached to a force transducer (Letica TRI-201). After the ring was suspended 179 in the organ bath, a passive force of 2 g was imposed and the preparation was stabilized for 1 180 h, being washed every 20 min. The signals from the force transducers were amplified and 181 driven into an analog-digital board (DT16EZ, Data Translation, Marlboro, MA, USA) 182

mounted in a computer. On-line recordings and files for later processing were obtained withthe software Labtech Notebook Pro (Laboratory Technology, Wilmington, MA, USA).

To assess the contractile response of the abdominal aorta, cumulative doses $(10^{-9} \text{ to } 4 \times 10^{-6} \text{ M})$ of noradrenaline (NA) were added to the organ bath sequentially. To avoid variations due to differences in ring size, the measured maximal contractile force for every NA concentration, was normalized to the wet tissue weight and expressed in grams of developed force divided by grams of wet tissue (gF/gW).

KS with a potassium solution was used to elicit maximal contractions by depolarization of the
VSMC. The solution was prepared increasing KCl to 80 mM in the KS; NaCl was lowered to
keep normal osmolarity.

The Acetylcholine (Ach) relaxation was studied on, aorta precontracted with NA. At the
contraction plateau, the effect of cumulative concentrations (10⁻⁸ -10⁻⁵) M of Ach was tested.
Maximal relaxant effect was expressed as percent of the maximal contraction obtained with
NA.

197 **2.10. Immunohistochemistry**

After antigen retrieval with 0.01 M citrate buffer, pH 6, endogenous peroxidase activity was blocked by incubation with 3% hydrogen peroxide in 60% methanol. Nonspecific binding sites were blocked with 5% BSA. Sections were incubated overnight at 4 °C with 1:200 rabbit polyclonal anti-ER α antibody (Santa Cruz Biotechnology, Inc.) or 1:200 rabbit polyclonal anti-AhR antibody (Santa Cruz Biotechnology, Inc.). After several washes, sections were incubated with secondary antibody biotinylated anti-rabbit IgG (Vector Laboratories Inc., Burlingame, CA, USA) for 2 h at room temperature. To reveal the sites of antigen/antibody binding, an avidin-horseradish peroxidase complex (Vector Laboratories, Burlingame, CA, USA) and the chromogen 3,3'diaminobenzidine tetrahydrochloride (DAB; Sigma, St. Louis, MO, USA) were used.

For control staining, some sections were incubated with PBS instead of primary antibody and no immunoreactivity was detected. Sections were dehydrated in ethanol, cleared in xylene and mounted in Canada balsam, and then photographed using Nikon Eclipse 80i microscope and Visiopharm Integrator System software.

212

213 2.11. Histological evaluations

Abdominal aorta samples (5 mm width) and ventricles from control and HCB-treated rats were fixed with 4% paraformaldehyde. Samples were dehydrated, and embedded in paraffin, serially sectioned in transversal orientation, and stained with Masson Trichrome. Planimetry was performed using a Nikon Eclipse E400 microscope and image analysis software (Image J, National Institute of Health, Bethesda, MD). Sections were magnified and digitalized. Vessel thickness and right and left ventricle cavities were then measured.

Arterial wall thickness and of ventricle areas were analysed in 10 slices of the same sampleand, at least, 5 measurements of each photograph were done.

222

223 2.12. Creatinine analysis

Serum creatinine was measured using the enzymatic method Creatinina enzimatica AA(Wiener Lab.) according to the manufacturer instructions.

227 2.13. Statistical analysis

Data were analyzed by one-way analysis of variance followed by *post hoc* tests (Bonferroni multiple comparison *t*-test) in order to evaluate selected pairs of groups. p < 0.05 was considered significant. SPSSTM version 15.0 software was used to analyze data.

231

3. RESULTS

233 **3.1. Blood pressure**

As dioxin TCDD induces hypertension in rats and HCB is considered a dioxin-like environmental pollutant, we evaluated whether HCB could induce alterations in BP in HCB (5 and 500 mg/kg)-treated rats, for 30, 35, 40 and 45 days (Figure 1).

When animals were treated with the lower dose, 5 mg/kg HCB, BP remain unchanged, but, when animals were treated with the highest dose, 500 mg/kg HCB, BP showed a significant increase at 35 days of HCB treatment which remained until 45 days (Figure 1). Similar BP increase by HCB treatment was observed in a model using male rats (data not shown).

Of note, no differences in neither mortality rate nor rat body weigh was observed between
groups (227.6 ± 2.7, 230.3 ± 14.0, 222.0 ± 8.3 g for control, 5 mg/kg HCB and 500 mg/kg
HCB respectively).

244

245 **3.2.** Arterial remodeling

It is well known that increased peripheral vascular resistance to blood flow, largely due to vascular remodeling, induces hypertension. In order to investigate whether the effect of HCB on BP can be attributed to this phenomenon, intima-media layer thickness was measured in abdominal aortae. Representative cross-sections of the aortic wall are shown in Figure 2A. Image analyses showed a significant increase (34.6%, $p \le 0.01$) in the intima-media layer thickness only in 500 mg/kg HCB-treated rats after 45 days of treatment, as compared to control rats (Figure 2B).

253

254 **3.3. PCNA protein level and VSMC number in abdominal aorta**

Considering the effect of HCB on abdominal aorta wall thickness, we analyzed PCNA protein expression as an index of cell proliferation and found a decrease in aortae from both 5 and 500 mg/kg HCB-treated rats (40.9%, p<0.01 and 36.6%, p<0.01, respectively) (Figure 2C and D). In addition, the number of cell nuclei per area in the arterial wall was analyzed. A 29.7% (p<0.05) decrease was observed at the highest dose of HCB (Figure 2E). Worth pointing out, 5 mg/kg HCB-treatment decreased the cell number, even though statistical significance was not reached.

262

263 **3.4. Heart remodeling**

Hypertension also causes compensatory structural changes in the heart. For this reason, we analysed the ratio between the left ventricle area, including the interventricular septum, and the left ventricular cavity (Figure 3A). No changes were observed between control and HCBtreated rats, which indicates the absence of adaptive mechanisms (Figure 3B).

In addition, the ventricle weights were analyzed and no significant differences between groups were observed $(3.35 \pm 0.02, 3.49 \pm 0.23 \text{ and } 3.56 \pm 0.10 \text{ mg}$ of ventricles/g of rat for control, 5 mg/kg HCB and 500 mg/kg HCB, respectively).

271 **3.5. Kidney function**

Hypertension also impacts kidney function, damaging kidney blood vessels and reducing their ability to work properly. Therefore, we evaluated kidney function by determining creatinine concentration in serum of HCB-treated and control rats. Our results show that creatinine levels remain unchanged from control levels, regardless of HCB dose used (Figure 3C).

276

277 **3.6.** Vascular reactivity

With the aim to evaluate arterial function, the contractile response of rings of abdominal aorta to NA was evaluated. In both groups of HCB-treated rats, the aortic contractile response presents higher sensitivity to NA, evidenced by a significant reduction (41.3%, p<0.01 and 64.8%, p<0.001; respectively) in the half maximal effective concentration (EC₅₀).

In turn, the maximal contractile response to NA was significantly decreased in arteries from both HCB-treated groups compared to control group. Interestingly, 5 mg/kg HCB was more effective than 500 mg/kg HCB, to induce a decrease in the maximal contractile response, (52.4%, p<0.001 and 28.6%, p<0.05 respectively) (Figure 4).

We also evaluated the effect of Ach on arterial relaxation after NA contraction. It was observed a significant (p< 0.05) diminution in the maximal percentage of relaxation in arteries rings of both 5 and 500 mg/kg HCB-treated rats compared to control (22.93 \pm 2.82%, 11.4 \pm 2.81% and 12.8 \pm 1.6% for control, 5 mg/kg HCB and 500 mg/kg HCB, respectively)

290

291 **3.7.** Key molecules involved in blood pressure regulation

292 **TGF-**β1

As abundant evidence shows that TGF- β 1 is involved in the pathogenesis of hypertension (Ohta et al., 1994), we analyzed its arterial expression by semiquantitative RT-PCR (Figure 5). Rats treated with 500 mg/kg HCB showed a significant increase in TGF- β 1 mRNA levels (35%, *p* < 0.05). Worth pointing out, treatment with 5 mg/kg HCB exhibited an increasing – though statistically non-significant– tendency in growth factor expression.

298 **ΕR**α

An increase in BP could be the consequence of arterial function alterations. The arterial 299 function is finely controlled by contraction and relaxation processes of VSMC. In this sense, a 300 direct relationship has been widely established between vasodilation and ERa expression 301 (Reslan et al., 2013). Therefore, and given that HCB has been shown to alter ERa protein 302 content in mammary gland (Li et al., 2008; Pena et al., 2012), we evaluated ER α expression 303 in aortae from both groups of HCB-treated rats. Immunohistochemistry shows that ERa 304 expression decreased in 500 mg/kg HCB-treated rats (Figure 6A). Similarly, Western blot 305 analysis showed that ER α protein content was significantly decreased (23.6%, p < 0.05) by 306 500 mg/kg HCB, whereas it was unaffected in arteries from HCB (5 mg/kg)-treated rats 307 (Figure 6B). 308

309 eNOS

It is well known that estradiol has vessel relaxation effects through the release of several factors. Among them, NO is generated by an increase in eNOS activity, in turn produced by protein induction through ER α . For this reason, we evaluated eNOS expression in arteries from HCB-treated rats. As shown in Figure 6D and E, a significant dose-dependent decrease of 25 (p < 0.05) and 40% (p < 0.01) eNOS expression was observed in arteries of rats treated with 5 and 500 mg/kg HCB, respectively.

316 AT1

One of the main mechanisms involved in arterial vasoconstriction is an increment in AT1 expression. Therefore, we evaluated whether HCB also modifies the expression of this receptor. As indicated in Figure 6F and G, a significant increase in AT1 (48.2%, p < 0.05) was observed in arteries from rats treated with 500 mg/kg HCB.

321 TH

TH are involved in the regulation of contraction and relaxation mechanisms. HCB is a hormonal disruptor which impacts TH homeostasis. For this reason, studies were carried out on possible TH alterations in aortic arteries from HCB-treated rats (Table 1). Results rendered no changes in T₃, or TSH levels in serum of rats treated with either dose of HCB. However, T₄ levels were significantly lower in rats with 500 mg/kg HCB (50.9%, ** p< 0.01).

Since DII is crucial in the regulation of local T₃ concentrations, HCB effect on abdominal aortae DII mRNA expression (the deiodinase isoform expressed in arteries) was evaluated by semiquantitative RT-PCR. Our results demonstrate that a significant dose-dependent decrease of 31.5 (p < 0.05) and 55.4% (p < 0.01) in DII mRNA was observed in arteries of rats treated with 5 and 500 mg/kg HCB, respectively (Table 1).

332 **3.8.** AhR activation

As a dioxin-like toxicant, HCB exerts both AhR-dependent and independent effects. We studied arterial AhR activation by analysis of its translocation to the nucleus by immunohistochemestry in rats treated with 5 and 500 mg/kg HCB (Figure 6 panel H).

AhR activity in arteries from 5 mg/kg HCB-treated rats showed an increasing tendency although not statically significant, whereas, a statistically greater expression in the nuclear zone in 500 mg/kg HCB treated rats was observed (Figure 6 panel I).

340 4. DISCUSSION

Even when several studies have informed about the effect of HCB on different tissues, data on its influence on the cardiovascular system are scarcely documented (Sjoberg Lind et al., 2013). In this context, the present work constitutes an original initial approach to the analysis of the effect of HCB on the cardiovascular system and unveils some of the molecular mechanisms involved.

The effect of HCB on BP regulation was evaluated using two different gavage-administered 346 doses of HCB during 45 days. The criteria used for dose selection were based on TH levels. 347 The lower dose (5 mg/kg) does not alter TH levels but the highest dose (500 mg/kg) generates 348 hypothyroxinemia (Kleiman de Pisarev et al., 1990). According to previous reports, the 349 administration of 500 mg/kg HCB for 30 days results in an HCB serum concentration of 6 350 351 ng/ml, which is equivalent to HCB serum of a widely exposed population (4.1 - 10.6 ng/ml)(Ohta et al., 1994). For this reason, our findings show a new pathological aspect to take into 352 353 account in the evaluation of human populations with high HCB exposure.

Worth pointing out, the high discrepancy observed between the HCB administered and the serum concentrations responds to the fact that only 2-5% HCB is absorbed when administered as an aqueous suspension (Koss and Koransky, 1975).

We observed an increase in systolic BP in rats exposed to the high dose of HCB, which allowed us to classify this toxic as a hypertensor. Worth pointing out, this increase in systolic BP was evident as late as 35 days of HCB treatment, suggesting an indirect effect of the toxic on vessels. Of note, our results are in line with those reported by Kopf et al., who demonstrated that dioxin TCDD induces hypertension *in vivo* and *in vitro* models (Kopf et al., 2008) and also in agreement with previous work by Dalton et al. which did not show significant increases in BP until day 23–29 of TCDD treatment (Dalton et al., 2001).

The persistent increment in vascular resistance is the main hemodynamic characteristic of 364 hypertension (Intengan and Schiffrin, 2000). This can be produced by functional 365 vasoconstriction or structural vessel wall thickening as a consequence of hypertrophy, 366 367 hyperplasia or both phenomena together. In this work, 500 mg/kg HCB-administered rats exhibited wall thickening and a decrease in PCNA expression in abdominal aorta. In 368 369 agreement, the number of aorta nuclei was decreased in HCB-treated rats. Although we have not studied the arterial extracellular matrix content, these results suggest that wall thickening 370 may be due to the hypertrophic effect of HBC on VSMC and not to hyperplasic growth. 371

Hypertension is also known to participate in heart remodeling. Sjoberg *et al.*, (2013) showed that long exposure to HCB is related to increased wall thickness of the left ventricle and concentric left ventricular remodeling (Sjoberg Lind et al., 2013) . However, in this work, no compensatory structural changes were observed in the hearts of HCB-treated rats. It is possible that this absence of heart remodeling responds to the relatively short duration of the hypertensive state.

In turn, kidney plays a key role in and is affected by BP regulation, as high BP can damage 378 kidney blood vessels, reducing their ability to work properly. In this work a conserved kidney 379 function was observed in HCB-treated rats. These results are in line with previous studies 380 covering less than 7 weeks of intoxication (Andrews et al., 1988; Richter et al., 1981) 381 although an increase in kidney weight was reported in longer HCB exposure (Andrews et al., 382 1989; Den Besten et al., 1993; Smith et al., 1985). Furthermore, HCB nephrotoxicity is a sex-383 dependent phenomenon, as it is observed in male but not female rats (Smith et al., 1985; 384 Arnold et al., 1985). 385

Alterations in arterial function, finely controlled by contraction and relaxation processes of VSMC, can trigger an increase in BP. In this work, these alterations were observed in abdominal aorta from HCB-treated animals, reflected by an increase in NA sensitivity and a

decrease in maximal response to NA. An increase in NA sensitivity could contribute to 389 vascular resistance when the sympathetic neurotransmitter is released and, in this way, 390 increase BP in HCB treated animals. The decrease in maximal response to NA could indicate 391 a imbalance in molecules with relaxing or contractile actions or a toxic effect of HCB on the 392 contractile machinery of VSMC. It is worth highlighting that this decrease was sharper in 393 arteries from 5 mg/kg than 500 mg/kg HCB-administered rats, suggesting that HCB could act 394 by different mechanisms depending on the dose used. In addition, the fact that BP is not 395 modified at 5 mg/kg HCB despite the lower capacity of contraction by NA could be explained 396 by compensatory responses that lead to BP normalization. In contrast, the increase in BP in 397 the 500 mg/kg HCB group may be explained by structural changes in the arterial wall 398 interfering with such compensatory mechanisms. 399

Among the components of the vascular wall, the endothelium plays an important role in vascular remodeling as a reaction to disruptive stimuli. Shear stress induced by blood flow on endothelial cells can result in structural and functional modifications affecting the production or release of several vasoactive factors, as well as a response to these factors (Urschel et al., 2012). In this sense, endothelial cells produce TGF- β 1, among others, which exerts its effect on VSMC (Simionescu et al., 2005).

In previous work we demonstrated an increase in TGF- β 1 expression in liver from HCBtreated rats which correlated with a imbalance between cell proliferation and apoptosis (Giribaldi et al., 2011). In this work, an increment in TGF- β 1 was observed in abdominal aortae from HCB-treated rats, which could constitute an adaptive endothelial homeostatic response to structural alterations in order to regulate cell proliferation. The increase in TGF- β 1 could also explain, in part, the reduction observed in aortic cell proliferation in the 500 mg/kg HCB group. This is in line with results reported by Sato *et al.*, (1995) showing a role

- 413 for TGF-β1 in the control of VSMC growth. TGF-β1 inhibits serum-induced proliferation of
- 414 rat aortic VSMC and concurrently induces cellular hypertrophy (Sato *et al.*, 1995).

In this study, a significant reduction was observed in ER α expression in abdominal arteries of 415 high-dose HCB-treated rats, in line with previous results indicating that HCB alters ERa 416 protein content in mammary gland (Li et al., 2008; Pena et al., 2012). Of note, it has also been 417 demonstrated that TGF- β 1 diminishes the expression of ER α in several tissues (Petrel and 418 Brueggemeier, 2003). Moreover, we have recently reported that an increase in TGF-B1 419 expression in the intimal layer of coronary arteries correlates with a decrease in ERa 420 expression (Castilla et al., 2014). Also, TGF-B1 has been shown to produce increased 421 proteasome-dependent degradation of ERa in cancer cell lines (Petrel and Brueggemeier, 422 2003). ER α is a relevant molecule in the mechanism of arterial relaxation and it is involved in 423 the protective effect of estrogens on the endothelium (Arnal et al., 2007; Jobe et al., 2013). 424 Estradiol increases eNOS activity through ERa by both Akt-mediated phosphorylation and 425 protein induction. This results in the generation and release of NO (Chambliss et al., 2000; 426 Rosenfeld et al, 2003, Jobe, S.O. 2013), which in turn relaxes VSMC, thus producing vessel 427 relaxation. Accordingly, in this work, a decrease was also observed in eNOS expression in 428 HCB treated rats. The decrease in NO levels in HCB-treated rats was also indirectly 429 corroborated by the Ach relaxation tests, since it acts through NO production. 430

In addition, an increase in arterial AT1 expression, the receptor by which angiotensin II acts as a vasoconstrictor, was observed in HCB-treated rats, in agreement with results reported in previous work in hypertensive rats (Romero-Nava *et al.*, 2016). Worth highlighting, TGF- β 1 was also shown to enhance AT1 gene expression in several tissues such as lung fibroblasts (Renzoni *et al.*, 2004), adrenal cells (Lebrathon *et al.*, 1994.), and trophoblasts (Tower *et al.*, 2005.) but to decrease AT1 levels in VSMCs (Zhang *et al.*, 2012), which indicates the importance of the physiological cell microenvironment in the final expression pattern. Altogether, these findings could partly explain the decrease in NA-induced vessel contraction observed in arteries from 500 mg/kg HCB-treated rats. Further studies should be conducted to clarify the mechanisms underlying the lower contraction observed at low doses of toxic in aorta rings. Likewise, such alterations observed *in vitro* assays at 5 mg/kg HCB did not correlate with a BP increase in the *in vivo* model.

Furthermore, and given that HCB is a hormonal disruptor, analyses were conducted on TH 443 concentration, of strong impact on vascular resistance and contraction. Yoneda et al. (1998) 444 have demonstrated that a single T_3 or T_4 injection promotes fast vasodilation in rat coronary 445 arteries. In contrast, the loss of vasodilation has been associated with an increase in systemic 446 vascular resistance, as observed in hypothyroidism (Duntas and Biondi, 2011; Klein and 447 Ojamaa, 2001). In this work, we also observe that HCB generates hypothyroxinemia, which 448 agrees with our previous report (Kleiman de Pisarev et al., 1990). Hypothyroxinemia, in turn, 449 450 can affect tissue concentration of TH and their possible effects on downstream intracellular signal transduction. 451

This work also evaluated the levels of arterial DII expression, as it is responsible for T4 metabolism to generate T3. A dose-dependent decrease in DII expression was observed in abdominal arteries from HCB-treated rats, which, together with the lower levels of T4 in serum, could be responsible for the decrease in arterial T3 levels. Moreover, it has been widely established that a decrease in T3 causes a reduction in ER α expression (Davis *et al.*, 2002; Faustino *et al.*, 2015) which could be in line with the decrease in ER α expression observed in this work.

As a dioxin-like toxicant, HCB exerts both AhR-dependent and independent effects. AhR is a transcription factor that modulates processes such as apoptosis, cell proliferation and migration. Even though AhR is located in the cytosol, the binding of HCB to AhR can trigger the toxic- AhR complex translocation to the nucleus in order to modulate the expression of 463 genes with dioxin response elements (DREs) in their promoters. In this work we have shown 464 artery AhR activation by 500mg/kg HCB but not by 5 mg/kg HCB, which indicates that the 465 mechanism by which these doses act is different. In addition, we conclude that at least part of 466 500 mg/kg HCB effects on the vascular system are exerted through AhR.

467 CONCLUSIONS

This work shows for the first time that subchronic administration of HCB generates arterial hypertension in rats, associated with impaired vascular reactivity. In this scenario, TGF- β 1, ER α , eNOS, AT1 and DII could constitute key molecules involved in the mechanism of HCB action and the development of hypertensive pathology. These findings may facilitate an experimental model for the development of more precise therapeutic targets in the treatment of HCB-induced hypertension.

474

475 **5. ACKNOWLEDGEMENTS**

This work was supported by grants from the Concejo Nacional de Investigaciones Científicas
y Tecnológicas (CONICET) (PIP 6060 and 0739) and Universidad de Buenos Aires,
Argentina (M041 and 6932/13).

479

480 **6. REFERENCES**

- 481 Agency for Toxic Substances and Disease Registry (ATSDR), (1998). Public Health
 482 Statement Dioxins.
- Agency for toxic substances, disease registry (ATSDR), (2002). Toxicological Profile for
 Hexachlorobenzene, USDOHAH Services, Atlanta, CA. USA.
- 485 Alvarez, L., Hernandez, S., Martinez-de-Mena, R., Kolliker-Frers, R., Obregon, M.J.,
- 486 Kleiman de Pisarev, D.L., (2005) The role of type I and type II 5' deiodinases on

- 487 hexachlorobenzene-induced alteration of the hormonal thyroid status. *Toxicology*.
 488 207(3), 349-362.
- Andrews, J.E., Courtney, K.D., Donaldson, W.E., (1988) Impairment of calcium homeostasis
 by hexachlorobenzene (HCB) exposure in Fischer 344 rats. *J Toxicol Environ Health*.
 23(3), 311-320.
- 492 Andrews, J.E., Courtney, K.D., Stead, A.G., Donaldson, W.E., (1989) Hexachlorobenzene-
- 493 induced hyperparathyroidism and osteosclerosis in rats. *Fundam Appl Toxicol.* 12(2),
 494 242-251.
- Arnal, J.F., Scarabin, P.Y., Tremollieres, F., Laurell, H., Gourdy, P., (2007) Estrogens in
 vascular biology and disease: where do we stand today? *Curr Opin Lipidol*. 18(5),
 554-560.
- 498 Arnold, D.L., Moodie, C.A., Charbonneau, S.M., Grice, H.C., McGuire, P.F., Bryce, F.R.,
- Collins, B.T., Zawidzka, Z.Z., Krewski, D.R., Nera, E.A., et al., (1985) Long-term
 toxicity of hexachlorobenzene in the rat and the effect of dietary vitamin A. *Food Chem Toxicol.* 23(9), 779-793.
- Carretero, O.A., Oparil, S., (2000) Essential hypertension. Part I: definition and aetiology.
 Circulation. 101(3), 329-335.
- 504 Castilla, R., Cao, G., Frers, R.K., Muller, A., Ambrosio, G., Milei, J., (2014) Inverse
- 505 expression of estrogen receptor alpha and apolipoprotein B in coronary intimal
- 506 hyperplasia of surgically repaired congenital heart disease: a pre-atherosclerotic
- 507 condition? *Int J Cardiol.* **177**(2), 548-550.
- 508 Chambliss, K.L., Yuhanna, I.S., Mineo, C. *et al.*, (2000) Estrogen receptor alpha and
- endothelial nitric oxide synthase are organized into a functional signaling module in
 caveolae. *Circ Res.* 87, E44–E52
- 511 Chaudhary, M., Chaudhary, S., (2017) Unravelling the Lesser Known Facets of Angiotensin
- 512 II Type 1 Receptor. *Curr Hypertens Rep.* **19**(1), 1.

- Davis, P.J., Tillmann, H.C., Davis, F.B., Wehling, M., (2002) Comparison of the mechanisms
 of nongenomic actions of thyroid hormone and steroid hormones. *J Endocrinol Invest*.
 25(4), 377-388.
- 516 Dalton, T.P., Kerzee, J.K., Wang, B., Miller, M., Dieter, M.Z., Lorenz, J.N., Shertzer, H.G.,
- 517 Nerbert, D.W., Puga, A., (2001). Dioxin exposure is an environmental risk factor for
 518 ischemic heart disease. *Cardiovasc Toxicol* 1, 285-298.
- den Besten, C., Bennik, M.H., Bruggeman, I., Schielen, P., Kuper, F., Brouwer, A., Koeman,
- 520 J.H., Vos, J.G., Van Bladeren, P.J., (1993) The role of oxidative metabolism in
- 521 hexachlorobenzene-induced porphyria and thyroid hormone homeostasis: a
- 522 comparison with pentachlorobenzene in a 13-week feeding study. *Toxicol Appl*
- 523 *Pharmacol.* **119**(2), 181-194.
- 524 Derhaschnig, U., Shehata, M., Herkner, H., Bur, A., Woisetschläger, C., Laggner, A.N.,
- Hirschl, M.M., (2002) Increased levels of transforming growth factor-beta1 in
 essential hypertension. *Am J Hypertens*. 15(3), 207-211.
- 527 Duntas, L.H., Biondi, B., (2011) New insights into subclinical hypothyroidism and
 528 cardiovascular risk. *Semin Thromb Hemost.* 37(1), 27-34.
- Environmental Toxicology Branch, Agency for toxic substances, disease registry (ATSDR),
 (2002).
- Faustino, L.C., Gagnidze, K., Ortiga-Carvalho, T.M., Pfaff, D.W., (2015) Impact of Thyroid
 Hormones on Estrogen Receptor α-Dependent Transcriptional Mechanisms in
- 533 Ventromedial Hypothalamus and Preoptic Area. *Neuroendocrinology*. 101(4), 331534 346.
- 535 Giribaldi, L., Chiappini, F., Pontillo, C., Randi, A.S., Kleiman de Pisarev, D.L., Alvarez, L.,
- 536 (2011) Hexachlorobenzene induces deregulation of cellular growth in rat liver.
- 537 *Toxicology*. **289**(1), 19-27.

- Huang, X., Lessner, L., Carpenter, D.O., (2006). Exposure to persistent pollutants and
 hypertensive disease. *Environ. Res.* 102, 101–106.
- 540 Intengan, H.D., Schiffrin, E.L., (2000) Structure and mechanical properties of resistance
 541 arteries in hypertension, role of adhesion molecules and extracellular matrix

542 determinants. *Hypertension*. **36**(3), 312-318.

- Jobe, S.O., Ramadoss, J., Wargin, A.J., Magness, R.R., (2013) Estradiol-17beta and its
- 544 cytochrome P450- and catechol-O-methyltransferase-derived metabolites selectively
 545 stimulate production of prostacyclin in uterine artery endothelial cells: role of estrogen
 546 receptor-alpha versus estrogen receptor-beta. *Hypertension*. **61**(2), 509-518.
- 547 Kleiman de Pisarev, D.L., Rios de Molina, M.C., San Martin de Viale, L.C., (1990) Thyroid
- function and thyroxine metabolism in hexachlorobenzene-induced porphyria. *Biochem Pharmacol.* **39**(5), 817-825.
- Klein, I., Ojamaa, K., (2001) Thyroid hormone: targeting the vascular smooth muscle cell.
 Circ Res. 88(3), 260-261.
- 552 Kopf, P.G., Huwe, J.K., Walker, M.K., (2008) Hypertension, cardiac hypertrophy, and
- impaired vascular relaxation induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin are
 associated with increased superoxide. *Cardiovasc Toxicol.* 8(4), 181-193.
- Koss, G, Koransky, W., (1975) Studies on the toxicology of hexachlorobenzene. I.
 Pharmacokinetics. *Arch Toxicol.* 34(3):203-12.
- Lebrathon, M., Jaillard, C., Naville, D., Begeot, M., Saez, J., (1994) Effects of transforming
 growth factor- β 1 on human adrenocortical fasciculate-reticularis cell differentiated
 functions. *J Clin Endocrinol Metab.* **79**, 1033–1039.
- Li, J., Li, N., Ma, M., Giesy, J.P., Wang, Z., (2008) In vitro profiling of the endocrine
 disrupting potency of organochlorine pesticides. *Toxicol Lett.* 183(1-3), 65-71.

562	Mills K.T., Bundy J.D., Kelly T.N., Reed J.E., Kearney P.M., Reynolds K., Chen J., He J.,
563	(2016) Global Disparities of Hypertension Prevalence and Control: A Systematic
564	Analysis of Population-Based Studies From 90 Countries. Circulation. 134(6), 441-
565	450
566	Obregon, M.J., Mallol, J., Escobar Del Rey, F., Morreale de Escobar, G., (1981). Presence of
567	L-thyroxine and 3,5,3'-triiodothyronine in tissues from thyroidectomized rats.
568	Endocrinology. 109, 908-918.
569	Ohta, K., Kim, S., Hamaguchi, A., Yukimura, T., Miura, K., Takaori, K., Iwao, H., (1994)
570	Role of angiotensin II in extracellular matrix and transforming growth factor-beta 1
571	expression in hypertensive rats. Eur J Pharmacol. 269(1), 115-119.
572	Pedram, A., Razandi, M., Aitkenhead, M., Hughes, C.C., Levin, E.R., (2002) Integration of
573	the non-genomic and genomic actions of estrogen. Membrane-initiated signaling by
574	steroid to transcription and cell biology. J Biol Chem. 277(52), 50768-50775.
575	Peña, D., Pontillo, C., Garcia, M.A., Cocca, C., Alvarez, L., Chiappini, F., Bourguignon, N.,
576	Frahm, I., Bergoc, R., Kleiman de Pisarev, D., Randi, A., (2012) Alterations in c-
577	Src/HER1 and estrogen receptor alpha signaling pathways in mammary gland and
578	tumors of hexachlorobenzene-treated rats. <i>Toxicology</i> . 293(1-3), 68-77.
579	Petrel, T.A., Brueggemeier, R.W., (2003) Increased proteasome-dependent degradation of
580	estrogen receptor-alpha by TGF-beta1 in breast cancer cell lines. J Cell Biochem.
581	88 (1), 181-190.
582	Renzoni, E.A., Abraham, D.J., Howat, S., Shi-Wen, X., Sestini, P., Bou-Gharios, G., Wells,
583	A.U., Veeraraghavan, S., Nicholson, A.G., Denton, C.P., Leask, A., Pearson, J.D.,
584	Black, C.M., Welsh, K.I., deBois, R.M., (2004) Gene expression profiling reveals
585	novel TGF-β1 targets in adult lung fibroblasts. <i>Respir Res.</i> 5, 24,

586	Reslan, O.M., Yin, Z., do Nascimento, G.R., Khalil, R.A., (2013) Subtype-specific estrogen
587	receptor-mediated vasodilator activity in the cephalic, thoracic, and abdominal
588	vasculature of female rat. J Cardiovasc Pharmacol. 62(1), 26-40.
589	Ribas-Fito, N., Sunyer, J., Sala, M., Grimalt, J.O., (2003) Changes in concentrations of
590	organochloride compounds in women from Flix, Tarragona [Spain]. Gac Sanit. 17(4),
591	309-11.
592	Richter E, Renner G, Bayerl J, (1981). Differences in the biotransformation of
593	hexachlorobenzene (HCB) in male and female rats. Chemosphere. 10:779-785.
594	Romero-Nava, R., Rodriguez, J.E., Reséndiz-Albor, A.A., Sánchez-Muñoz, F., Ruiz-
595	Hernandéz, A., Huang, F., Hong, E., Villafaña, S., (2016) Changes in protein and gene
596	expression of angiotensin II receptors (AT1 and AT2) in aorta of diabetic and
597	hypertensive rats. Clin Exp Hypertens. 38(1), 56-62,
598	Rosenfeld, C.R., Chen, C., Roy, T., and Liu, X., (2003) Estrogen selectively up-regulates enos
599	and nnos in reproductive arteries by transcriptional mechanisms. J Soc Gynecol
600	Investig. 10, 205–215.
601	Sato, M., Ohsaki, Y., Tobise, K., (1995) Transforming growth factor-beta 1 proliferated
602	vascular smooth muscle cells from spontaneously hypertensive rats. Am J Hypertens.
603	8 (2), 160-166.
604	Simionescu, A., Philips, K., Vyavahare, N., (2005) Elastin-derived peptides and TGF-beta1
605	induce osteogenic responses in smooth muscle cells. Biochem Biophys Res Commun.
606	334 (2), 524-32.
607	Sjoberg Lind, Y., Lind, L., Salihovic, S., van Bavel, B., Lind, P.M., (2013) Persistent organic
608	pollutants and abnormal geometry of the left ventricle in the elderly. J Hypertens.

09 **31**(8), 1547-1553.

610	Smith, A.G., Francis, J.E., Dinsdale, D., Manson, M.M., Cabral J.R., (1985)
611	Hepatocarcinogenicity of hexachlorobenzene in rats and the sex difference in hepatic
612	iron status and development of porphyria. Carcinogenesis. 6(4), 631-636.
613	Tower, C.L., Chappell, S.L., Morgan, K., Kalsheker, N., Baker, P.N., Morgan, L.J., (2005)
614	Transforming growth factor β 1 regulates angiotensin II type 1 receptor gene
615	expression in the extravillous trophoblast cell line SGHPL-4. Mol Hum Reprod. 11,
616	847-852.
617	Urschel, K., Cicha, I., Daniel, W.G., Garlichs, C.D., (2012) Shear stress patterns affect the
618	secreted chemokine profile in endothelial cells. Clin Hemorheol Microcirc. 50(1-2),
619	143-152.
620	Xu, W., Wang, X., Cai, Z., (2013) Analytical chemistry of the persistent organic pollutants
621	identified in the Stockholm Convention: A review. Anal Chim Acta. 790, 1-13.
622	Yoneda, K., Takasu, N., Higa, S., Oshiro, C., Oshiro, Y., Shimabukuro, M., Asahi, T., (1998)
623	Direct effects of thyroid hormones on rat coronary artery: nongenomic effects of
624	triiodothyronine and thyroxine. Thyroid. 8(7), 609-613.
625	Zhang, X.H., Zheng, B., Gu, C., Fu, J.R., Wen, J.K., (2012) TGF-B1 downregulates AT1
626	receptor expression via PKC-δ-mediated Sp1 dissociation from KLF4 and Smad-
627	mediated PPAR-γ association with KLF4. Arterioscler Thromb Vasc Biol. 32, 1015–
628	1023.

FIGURE LEGENDS

Figure 1: **Systolic BP**. Systolic arterial pressure was measured at the indicated days of HCB treatment. White bands indicate control animals; grey and black bands indicate rats treated with 5 and 500 mg/kg HCB, respectively. Data represent the mean \pm SEM of three independent experiments * p< 0.05; ** p< 0.01; *** p< 0.001 vs. control at each time; + p< 0.05; ++ p< 0.01,+++ p< 0.001 vs. 5 mg/kg HCB group (n = 6 per group).

Figure 2: **Abdominal aorta structure and PCNA protein level:** Arteries from control and HCB-treated rats were stained with Masson Trichrome. Panel A: Representative microphotographs. Artery wall thickness was measured in the arterial photographs using Image J software (bars indicate how measurements were done). Panel B: Quantification of arterial wall thickness. Panel C: Representative Western blot of PCNA from control and HCB-treated rats. Panel D: Quantification of PCNA protein levels by densitometric scanning of immunoblots. Panel E: Quantification of VSMC number in the arterial wall. Data represent the mean \pm SEM of three independent experiments * p< 0.05; ** p< 0.01; *** p< 0.001 vs. control (n = 6 per group).

Figure 3: Heart remodeling and serum creatinine levels. Heart sections from control and HCB-treated rats were stained with Masson Trichrome. Panel A: Representative photographs. Panel B: Morphometric quantification. Panel C: serum creatinine concentration from control and HCB-treated rats. Data represent the mean \pm SEM of three independent experiments (n = 6 per group).

Figure 4: Abdominal aorta ring response to NA. The arterial contractile response to NA was analyzed *in vitro* Data represent the mean \pm SEM of three independent experiments * p< 0.05, ** p< 0.01 vs. control; + p < 0.05 vs. 5 mg/kg HCB group (n = 6 per group).

Figure 5: TGF- β 1 expression. Rats were treated with HCB (5 and 500 mg/kg) for 45 days. Panel A: Representative pattern of semiquantitative RT-PCR amplification of TGF- β 1 and GPDH cDNA. Panel B: Quantification of cDNAs normalized to GPDH cDNA. Panel C. Representative Western blot of TGF- β 1 \square . Panel C: Quantification of western blot TGF- β 1 IOD normalized to the corresponding β -actin signal. Data represent the mean \pm SEM of three independent experiments. * *p* < 0.05 *vs.* control; + *p* < 0.05 vs. 5 mg/kg HCB group (n = 6 per group).

Figure 6: ER α , eNOS and AT1 expression, and AhR activation. Rats were treated with HCB (5 and 500 mg/kg) for 45 days. Panel A: Representative image of ER α arterial immunohistochemistry. Panel B: Representative Western blot of ER α \square . Panel C: Quantification of western blot ER α IOD normalized to the corresponding β -actin signal. Panel D: Representative Western blot of eNOS. Panel E: Quantification of eNOS IOD normalized to the corresponding β -actin signal. Panel F: Representative Western blot of AT1. Panel G: Quantification of AT1 IOD normalized to the corresponding β actin signal. Panel H: Representative image of AhR arterial immunohistochemistry. Panel I: Quantification of cell number with nuclear AhR signal. Arrows indicate positive signal. Data represent the mean \pm SEM of three independent experiments. * p <0.05, ** p < 0.01 vs. control; + p < 0.05 vs. 5 mg/kg HCB group (n = 6 per group).





в













HCB (mg/kg)



А



HCB (mg/kg)



← Control
 ← HCB 5 mg/kg
 ← HCB 500 mg/kg





С









D

н









ACCEPTED MANUSCRIPT

1 TABLE 1: TH serum levels and aortic DII expression

	TSH (ng/ml)	T₄ (μg/dl)	T₃ (μg/dl)	mRNA DII/GPDH	
Control	1.90 ± 0.30	4.16 ± 0.43	74.00 ± 4.60	9.2 ± 0.2	K
HCB 5 (mg/kg)	1.99 ± 0.38	3.95 ± 0.27	76.00 ± 2.90	6.3 ± 0.1*	
HCB 500 (mg/kg)	2.07 ± 0.27	2.01 ± 0.20**+	72.50 ± 5.50	4.1 ± 0.1**+	

Table 1: Data represent the mean \pm SEM of three independent experiments, * p < 0.05 and **

p < 0.01 vs. control; + p < 0.05 vs. 5 mg/kg HCB group (n = 6 per group).