

# THE TANGO BETWEEN PKA AND CaMKII SIGNALING IN CARDIAC PACEMAKER CELLS

**Yael Yaniv**

Biomedical Engineering Faculty, Technion-IIT, Haifa, Israel

**Correspondence to:**

Yael Yaniv, PhD (yaely@bm.technion.ac.il)

**ABSTRACT**

The sinoatrial node is the primary pacemaker that controls the heart rate under normal conditions. Although the heart rate was originally measured thousands of years ago, the mechanisms that control the spontaneous beating of the sinoatrial node (SAN) are still under debate. In the last century, SAN function was mostly investigated by electrophysiological tools. Therefore, not surprisingly, the major mechanisms that control SAN function were thought to be related only to membranal ionic modulations. Recent biophysical, biochemical and imaging techniques have shed new light on the role of intrinsic pacemaker mechanisms on SAN function. Specifically, the role of post-translational modification signaling on SAN function has been explored using numerical and experimental tools. We describe here the major breakthroughs related to these signaling mechanisms in SAN cells. We conclude that the recent findings are only the tip of the iceberg in the fascinating world of downstream post-translational modification signaling, and we point out future research directions that may increase our knowledge of pacemaker function.

**Keywords:** Coupled-clock system, Mathematical modeling, Pacemaker.


## Introduction

During hibernation a bear's heart rate can decrease to 10 beats/min [1] while 700 beats/min was recorded in short-tailed shrews [2]. The primary pacemaker in these mammals and others is the sinoatrial node (SAN). Although the heart pulse was originally measured more than thousands of years ago, the mechanisms that control the spontaneous beating of SAN cells are still under debate. The SAN consists of hundreds to thousands (depending on the mammal species) of pacemaker cells that can beat spontaneously even without any neural stimulation. Using recent biophysical, biochemical and imaging techniques, it has been revealed that pacemaker cell function is controlled by internal and membrane mechanisms. Moreover, the identity of the nodes that connect these mechanisms has been revealed. Here, we focus on the role of post-translational modification signaling on SAN function, as discovered using both numerical and experimental approaches. We describe here the major breakthroughs related to these signaling mechanisms in SAN cells.

### A coupled-clock system controls pacemaker function

Pacemaker function is orchestrated by two clocks (Fig. 1): the surface membrane clock (M clock), an ensemble of sarcolemmal electrogenic molecules, and the  $\text{Ca}^{2+}$  clock (the sarcoplasmic reticulum (SR)) (for a review see [3]). A numerical model (for a review see [4]) predicted that these clocks are coupled, as was confirmed by experimental measurements (for a review see [5]). The two clocks are coupled through  $\text{Ca}^{2+}$  signaling: local  $\text{Ca}^{2+}$  releases (LCRs) activate an inward  $\text{Na}^+/\text{Ca}^{2+}$  exchange (NCX) current and other  $\text{Ca}^{2+}$  dependent mechanisms (e.g.,  $\text{Ca}^{2+}$ -activated potassium channels [3] and  $\text{Ca}^{2+}$ -dependent inactivation of L-type  $\text{Ca}^{2+}$  channels) that prompt the M clock to generate an action potential (AP). By definition, the LCR signal is regulated by the SR  $\text{Ca}^{2+}$  content and thus by the  $\text{Ca}^{2+}$  clock. However, because the L-type  $\text{Ca}^{2+}$  channels regulate cell  $\text{Ca}^{2+}$  available for SR pumping, the LCR signal is controlled by both clocks, and its magnitude indicates the degree of cross-talk between the clocks [6, 7]. Thus, the  $\text{Ca}^{2+}$  is a node that connects the two clocks, where an amplification of LCR signal leads to an increase in the spontaneous AP firing rate and vice versa.  $\text{Ca}^{2+}$  not only directly couples the clocks, but also does so indirectly through the phosphorylation cascade:  $\text{Ca}^{2+}$  activates calmodulin-adenylyl cyclase (AC)-dependent protein kinase A (PKA) and  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II (CaMKII) [8, 9]. These phosphorylation signaling cascades act on the same  $\text{Ca}^{2+}$  (ryanodine receptors and phospholamban) and M-clock proteins (L type  $\text{Ca}^{2+}$  channels and  $\text{K}^+$  channels). The increase in phosphorylation activity leads to amplification of the LCR  $\text{Ca}^{2+}$  signal and therefore to an increase in the AP firing rate. Thus, both  $\text{Ca}^{2+}$  cycling and phosphorylation activity levels indicate the degree of clock coupling: when the signal is further amplified, so are the activity levels of the membrane and SR  $\text{Ca}^{2+}$  molecules, and thus the AP firing rate is higher as well

to the mammal each group worked with (mice vs. rabbits). Alternatively, it might be related to other non-specific gene changes in these mutant mice (e.g., L-type) that contribute to AP firing.




**Figure 1.** Schematic figure illustrating the cross-talk between M clock (membrane molecules) and the  $\text{Ca}^{2+}$  clock (sarcoplasmic reticulum (SR)). Two major nodes connect between the clocks:  $\text{Ca}^{2+}$  and CaMKII-cAMP/PKA signaling. Local  $\text{Ca}^{2+}$  release through the ryanodine receptor (RyR) increases inward sodium- $\text{Ca}^{2+}$  exchange current that ignites together with  $I_f$  the action potential. Increase in membrane potential lead to opening of L-type channels that reload the SR. Thus, both clocks components contribute to  $\text{Ca}^{2+}$  signaling.  $\text{Ca}^{2+}$  also works as indirect node: it activates calmodulin-adenylyl cyclase (AC)-dependent protein kinase A (PKA) and  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II (CaMKII). These phosphorylation signaling cascades act on the same  $\text{Ca}^{2+}$  (RyR and phospholamban (PLB)) and M-clock proteins (L type  $\text{Ca}^{2+}$  channels,  $\text{K}^+$  channels and NaK pump). Note that a different type of AC is activated by brain receptors.

### PKA and CaMKII are coupled

As discussed above, PKA and CaMKII are linked upstream (through  $\text{Ca}^{2+}$ -calmodulin activation). Because both PKA and CaMKII phosphorylate the same  $\text{Ca}^{2+}$  and M-clock proteins, and these protein activities affect the  $\text{Ca}^{2+}$  cycling, PKA and CaMKII are also linked downstream. Thus, theoretically, any change in either PKA or CaMKII activity should affect the other mechanism in steady state. Indeed, when a battery of drugs that disturb the M ( $I_f$  blocker) or  $\text{Ca}^{2+}$  clock (SERCA inhibitor or ryanodine blocker) activity is administered, the AP firing rate is reduced, as is the cAMP level in steady state; the same effect was obtained by administering drugs that disturb the interconnected signaling (PKA inhibitor and  $\text{Ca}^{2+}$  chelate) or the brain signaling (cholinergic receptor activator) [14] (Fig. 2A). Interestingly, application of CaMKII inhibitors (KN-93 or AIP) leads not only to a reduction in the AP firing rate but also to a reduction in cAMP level (Fig. 2B). Importantly, the relationship between spontaneous AP firing rate and cAMP level in response to the afferent battery of drugs is similar to the relationship between spontaneous AP firing rate and cAMP level in response to the CaMKII blocker [11]

(Fig. 2C). Thus, reduction in CaMKII signaling reduces the  $\text{Ca}^{2+}$  cycling, leading to a reduction in  $\text{Ca}^{2+}$ -activated AC activity and thus cAMP-PKA signaling. Theoretically, the opposite cascade from reduction in PKA signaling to reduction in CaMKII activity should exist; however, no experimental tools exist today to prove or disprove this theory.



**Figure 2.** The relationship between spontaneous action potential firing rate and steady state cAMP in response to (A) a battery of drugs: blebbistatin (blocks spontaneous contractions, 10  $\mu\text{M}$ ), CsCl ( $I_f$  blocker, 2mM), cyclopiazonic acid (inhibiting the sarcoplasmic reticulum  $\text{Ca}^{2+}$  pump, 5  $\mu\text{M}$ ), carbachol (activating cholinergic receptors, 0.2 $\mu\text{M}$ ), ryanodine (disabling ryanodine receptors, 30 $\mu\text{M}$ ), PKI (PKA inhibitor, 25 $\mu\text{M}$ ) and BABTA (buffering intracellular  $\text{Ca}^{2+}$ , 25  $\mu\text{M}$ ); (B) KN-92 (inactivated CaMKII inhibitor, 3  $\mu\text{M}$ ), AIP (inhibiting CaMKII, 2 and 10 $\mu\text{M}$ ), KN-93 (inhibiting CaMKII, 0.5 and 3  $\mu\text{M}$ ), and a combination of low AIP and low KN-93 (denoted in the figure as **both**). (C) Note the resemblance between the action potential beating interval in response to the battery of drugs in (A) (black line) and in response to CaMKII inhibition in (B) (red line). Data are from [11, 14].

### AC-cAMP-PKA signaling regulates SANC automaticity in all physiological ranges

Although the role of PKA as a key nodal signaling pathway that interacts with both M and  $\text{Ca}^{2+}$  clock functions has been known for more than 10 years [15], its magnitude, dynamics and spatiotemporal regulation have only recently been described in heart pacemaker cells [16]. In cultured adult rabbit pacemaker cells infected with an adenovirus expressing the FRET sensor AKAR3, the magnitude and time response of PKA activity and the spontaneous AP firing rate were similar in response to graded increases in the intensity of  $\beta$ -adrenergic receptor ( $\beta$ -AR) stimulation (by isoproterenol) or in response to phosphodiesterase (PDE) inhibition. Thus, the kinetics and stoichiometry of increases in PKA activity in response to a physiological ( $\beta$ -AR stimulation) or pharmacological (PDE inhibitor) stimuli match those of changes in the AP firing rate. Recent work has also shown that the kinetics and stoichiometry of decreases in PKA activity in response to a physiological stimulus (cholinergic receptor stimulation) match those of changes in the AP firing rate [17]. Therefore, AC-cAMP-PKA signaling

can play a similar role to that of PKA in pacemaker function. Future experiments on CaMKII activity, using approaches similar to those described here for measuring PKA signaling are needed.

### **PLB phosphorylation is an important mediator of changes in the spontaneous AP firing rate**

Although the experimental results described in the previous section revealed for the first time PKA kinetics and its importance to pacemaker function, the main molecular targets that mediate between PKA signaling and pacemaker function are not known. To identify these targets, a novel mechanistic model was developed [16, 17]. In addition to a description of membrane molecules and internal pacemaker mechanisms, the model includes a description of autonomic-nervous receptors and post-translation signaling cascades. The model reproduces the experiments described in section 4 and yields two important predictions: [1] AC-cAMP-PKA signaling is the core cascade between autonomic receptor stimulation and pacemaker function and [2] directly activating internal clock mechanisms can mimic the effect of autonomic receptor stimulation. However, an even more important prediction was that phospholamban is an important PKA target in allowing the fight-or-flight response, and an important component in maintaining the changes in AP firing rate in response to cholinergic stimulation. A similar conclusion on the important role of phospholamban in mediating the PKA effect on pacemaker function was reached before [18]. However, the model in [18] includes a phenomenological description of AC-cAMP-PKA signaling. Others also attempted to find the main molecular targets that mediate between PKA signaling and pacemaker function [19]. Although the Noma group [19] included in their model a description of cAMP/PKA signaling, they did not include the connection between  $\text{Ca}^{2+}$  and AC, a description of cholinergic receptor stimulation, and correct  $\text{Ca}^{2+}$  cycling dynamics.

The authors of [20] numerically modeled the effect of CaMKII level on pacemaker function. The model predicts how SANC death and fibrosis are expressed in myocardial infarction by oxidizing CaMKII affect pacemaker function. However, lack of experimental results on CaMKII dynamics prevents the development of a numerical mechanistic model similar to that described above (reviewed in [21]).

### **Mutations in the coupled clock pacemaker system involve changes in CaMKII-cAMP/PKA signaling**

As described above, alteration in CaMKII and/or cAMP/PKA signaling can lead to disturbances in clock function. However, any alteration in one clock mechanism will be transmitted to the other clock mechanism via changes in CaMKII and cAMP/PKA signaling, and all of these together will affect pacemaker function [22]. For example, ivabradine, a specific  $I_f$  blocker, can indirectly alter the release of LCR, a coupled-clock function parameter. The numerical model predicts that the ivabradine effect on coupled-clock system function is through the effect on  $\text{Ca}^{2+}$  cycling and  $\text{Ca}^{2+}$  activated AC-cAMP/PKA signaling [7]. Similarly, a mutation that increases the funny channel's sensitivity to cAMP (R524Q) leads to a shorter AP firing rate than that of wild-type cells, and this gain in pacemaker function is the net effect of the R514Q mutation on the functioning of the coupled-clock system through  $\text{Ca}^{2+}$  cycling and  $\text{Ca}^{2+}$ -activated AC-cAMP/PKA signaling (23). On the other hand, a mutation that induces decrease in  $I_f$  function (mutant S672R channels respond normally to cAMP but they are activated at more negative voltages) leads to a longer AP firing rate than that of wild-type cells, and this loss in pacemaker

function is the net effect of the mutation on the functioning of the coupled-clock system through  $\text{Ca}^{2+}$  cycling and  $\text{Ca}^{2+}$  activated AC-cAMP/PKA signaling [23]. Thus, AC-cAMP-PKA signaling is involved in both gain and loss of pacemaker cell function.

### Summary

PKA and CaMKII signaling are dancing the tango. By means of this dance, they control coupled-clock system function, either by directly changing their dynamics or by transmitting changes from one coupled clock mechanism to the others. Future works will reveal more about the spatial dynamics of PKA and CaMKII signaling in healthy pacemaker cells, in deteriorated pacemaker cells associated with different cardiac diseases and aging [24].

### References

- [1] **Evans, A.L., N.J. Singh, A. Friebe, J.M. Arnemo, T.G. Laske, O. Fröbert, J.E. Swenson, and S. Blanc.** Drivers of hibernation in the brown bear. *Front. Zool.* 2016. 13: 7.
- [2] **Pfeiffer, C.J., and J.C. Keith.** Cardiac ultrastructure and electrocardiogram of the short-tailed shrew, *Blarina brevicauda*. *J. Submicrosc. Cytol. Pathol.* 1993. 25: 571–81.
- [3] **Weisbrod, D., S.H. Khun, H. Bueno, A. Peretz, and B. Attali.** Mechanisms underlying the cardiac pacemaker: the role of SK4 calcium-activated potassium channels. *Acta Pharmacol. Sin.* 2016. 37: 82–97.
- [4] **Maltsev, V.A., Y. Yaniv, A. V Maltsev, M.D. Stern, and E.G. Lakatta.** Modern perspectives on numerical modeling of cardiac pacemaker cell. *J Pharmacol Sci.* 2014. 125: 6–38.
- [5] **Yaniv, Y., E.G. Lakatta, and V.A. Maltsev.** From two competing oscillators to one coupled-clock pacemaker cell system. *Front Physiol.* 2015. 6: 28.
- [6] **Yaniv, Y., A.E. Lyashkov, S. Sirenko, Y. Okamoto, T.R. Guiriba, B.D. Ziman, C.H. Morrell, and E.G. Lakatta.** Stochasticity intrinsic to coupled-clock mechanisms underlies beat-to-beat variability of spontaneous action potential firing in sinoatrial node pacemaker cells. *J Mol Cell Cardiol.* 2014. 77: 1–10.
- [7] **Yaniv, Y., S. Sirenko, B.D. Ziman, H.A. Spurgeon, V.A. Maltsev, and E.G. Lakatta.** New evidence for coupled clock regulation of the normal automaticity of sinoatrial nodal pacemaker cells: bradycardic effects of ivabradine are linked to suppression of intracellular  $\text{Ca}^{2+}$  cycling. *J Mol Cell Cardiol.* 2013. 62: 80–89.
- [8] **Mattick, P., J. Parrington, E. Odia, A. Simpson, T. Collins, and D. Terrar.**  $\text{Ca}^{2+}$ -stimulated adenylyl cyclase isoform AC1 is preferentially expressed in guinea-pig sinoatrial node cells and modulates the I(f) pacemaker current. *J Physiol.* 2007. 582: 1195–1203.
- [9] **Younes, A., A.E. Lyashkov, D. Graham, A. Sheydina, M. V Volkova, M. Mitsak, T.M. Vinogradova, Y.O. Lukyanenko, Y. Li, A.M. Ruknudin et al.**  $\text{Ca}^{2+}$ -stimulated basal adenylyl cyclase activity localization in membrane lipid microdomains of cardiac sinoatrial nodal pacemaker cells. *J Biol Chem.* 2008. 283: 14461–14468.
- [10] **Sirenko, S., D. Yang, Y. Li, A.E. Lyashkov, Y.O. Lukyanenko, E.G. Lakatta, and**

- T.M. Vinogradova.** Ca<sup>2+</sup>-dependent phosphorylation of Ca<sup>2+</sup> cycling proteins generates robust rhythmic local Ca<sup>2+</sup> releases in cardiac pacemaker cells. *Sci Signal.* 2013. 6: ra6.
- [11] **Yaniv, Y., H.A. Spurgeon, B.D. Ziman, and E.G. Lakatta.** Ca(2+)/calmodulin-dependent protein kinase II (CaMKII) activity and sinoatrial nodal pacemaker cell energetics. *PLoS One.* 2013. 8: e57079.
- [12] **Wu, Y., H.H. Valdivia, X.H.T. Wehrens, and M.E. Anderson.** A Single Protein Kinase A or Calmodulin Kinase II Site Does Not Control the Cardiac Pacemaker Ca<sup>2+</sup> Clock. *Circ. Arrhythm. Electrophysiol.* 2016. 9: e003180.
- [13] **Wu, Y., Z. Gao, B. Chen, O.M. Koval, M. V Singh, X. Guan, T.J. Hund, W. Kutschke, S. Sarma, I.M. Grumbach et al.** Calmodulin kinase II is required for fight or flight sinoatrial node physiology. *Proc Natl Acad Sci U S A.* 2009. 106: 5972–5977.
- [14] **Yaniv, Y., M. Juhaszova, A.E. Lyashkov, H. Spurgeon, S.J. Sollott, and E.G. Lakatta.** Ca<sup>2+</sup>-regulated-cAMP/PKA signaling in cardiac pacemaker cells links ATP supply to demand. *J Mol Cell Cardiol.* 2011. 51: 740–748.
- [15] **Vinogradova, T.M., A.E. Lyashkov, W. Zhu, A.M. Ruknudin, S. Sirenko, D. Yang, S. Deo, M. Barlow, S. Johnson, J.L. Caffrey et al.** High basal protein kinase A-dependent phosphorylation drives rhythmic internal Ca<sup>2+</sup> store oscillations and spontaneous beating of cardiac pacemaker cells. *Circ Res.* 2006. 98: 505–514.
- [16] **Yaniv, Y., A. Ganesan, D. Yang, B.D. Ziman, A.E. Lyashkov, A. Levchenko, J. Zhang, and E.G. Lakatta.** Real-time relationship between PKA biochemical signal network dynamics and increased action potential firing rate in heart pacemaker cells: Kinetics of PKA activation in heart pacemaker cells. *J Mol Cell Cardiol.* 2015. 86: 168–178.
- [17] **Behar, J., A. Ganesan, J. Zhang, and Y. Yaniv.** The Autonomic Nervous System Regulates the Heart Rate through cAMP-PKA Dependent and Independent Coupled-Clock Pacemaker Cell Mechanisms. *Front. Physiol.* 2016. 7: 419.
- [18] **Maltsev, V.A., and E.G. Lakatta.** A novel quantitative explanation for the autonomic modulation of cardiac pacemaker cell automaticity via a dynamic system of sarcolemmal and intracellular proteins. *Am J Physiol Hear. Circ Physiol.* 2010. 298: H2010-23.
- [19] **Himeno, Y., N. Sarai, S. Matsuoka, and A. Noma.** Ionic mechanisms underlying the positive chronotropy induced by beta1-adrenergic stimulation in guinea pig sinoatrial node cells: a simulation study. *J Physiol Sci.* 2008. 58: 53–65.
- [20] **Luo, M., X. Guan, E.D. Luczak, D. Lang, W. Kutschke, Z. Gao, J. Yang, P. Glynn, S. Sossalla, P.D. Swaminathan et al.** Diabetes increases mortality after myocardial infarction by oxidizing CaMKII. *J Clin Invest.* 2013. 123: 1262–1274.
- [21] **Yaniv, Y., and V.A. Maltsev.** Numerical Modeling Calcium and CaMKII Effects in the SA Node. *Front Pharmacol.* 2014. 5: 58.
- [22] **Yaniv, Y., and E.G. Lakatta.** Pacemaker gene mutations, bradycardia, arrhythmias and the coupled clock theory. *J Cardiovasc Electrophysiol.* 2013. 24: E28-9.
- [23] **Behar, J., and Y. Yaniv.** Dynamics of PKA phosphorylation and gain of function in cardiac pacemaker cells: a computational model analysis. *Am J Physiol Hear. Circ Physiol.* 2016. 310: H1259-66.
- [24] **Tsutsui, K., O.J. Monfredi, and E.G. Lakatta.** A general theory to explain heart rate and cardiac contractility changes with advancing age. *Phy Mini Rev.* 2016. 9(2): 9-25.

## About the authors



**Prof. Yael Yaniv** is an Assistant Professor in Biomedical Engineering Faculty in the Technion-Israel Institute of Technology. She received her BSc in both Electrical and Mechanical Engineering (cum laude, 2002), her MSc in Electrical Engineering (summa cum laude 2003), and her PhD in Biomedical Engineering (2007) all from the Technion-IIT, Israel. She was a postdoctoral fellow (2007-2012) and a Research Fellow (2012-2013) in cardiovascular science at the National Institute on Aging, National Institutes of Health, under Dr. Edward Lakatta. Her research focus on: (i) the identity of clock mechanisms and their coupling signal in sinoatrial, atrial and ventricular cells, (ii) identification of the entrainment between energy balance and clock mechanism functions and (iii) prediction of arrhythmogenic events by identifying in vivo changes in the degree of clock coupling. She has written several highly cited original and review papers in these fields.