

Accepted Manuscript

Suspected poisoning in beef cattle from ingestion of *Prosopis nigra* pods in north-western Argentina

Juan F. Micheloud, Luis A. Colque Caro, Luciana A. Cholich, Olga G. Martínez, Eduardo J. Gimeno



PII: S0041-0101(18)31023-7

DOI: <https://doi.org/10.1016/j.toxicon.2018.11.302>

Reference: TOXCON 6028

To appear in: *Toxicon*

Received Date: 6 August 2018

Revised Date: 10 November 2018

Accepted Date: 19 November 2018

Please cite this article as: Micheloud, J.F., Colque Caro, L.A., Cholich, L.A., Martínez, O.G., Gimeno, E.J., Suspected poisoning in beef cattle from ingestion of *Prosopis nigra* pods in north-western Argentina, *Toxicon* (2018), doi: <https://doi.org/10.1016/j.toxicon.2018.11.302>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Case Report**

2 **SUSPECTED POISONING IN BEEF CATTLE FROM INGESTION OF *Prosopis***
3 ***nigra* PODS IN NORTH-WESTERN ARGENTINA**

4 Juan F. Micheloud ^{a,b}, Luis A. Colque Caro ^a, Luciana A. Cholich ^{c,f}, Olga G. Martínez ^d,
5 Eduardo J. Gimeno ^{e,f}.

6 ^a *Grupo de Trabajo de Patología, Epidemiología e Investigación Diagnóstica, Área de*
7 *Sanidad Animal-IIACS/INTA, Salta, Argentina.*

8
9 ^b *Facultad de Ciencias Veterinarias y Agrarias-UCASAL, Salta, Argentina.*

10 ^c *Facultad de Ciencias Veterinarias-UNNE, Corrientes, Argentina.*

11 ^d *Cátedra de Diversidad de las Plantas, Universidad Nacional de Salta, Salta, Argentina.*

12 ^e *Cátedra de Patología General, Facultad de Ciencias Veterinarias-Universidad Nacional*
13 *de La Plata, Bs As, Argentina.*

14 ^f *Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.*

15 * *E-mail: micheloud.juan@inta.gob.ar*

16

17 **Abstract**

18 The aim of this paper was to present the first report of *Prosopis nigra* poisoning of
19 cattle in Argentina. Outbreaks occurred in five farms located in Salta and Santiago del
20 Estero provinces. All animals were examined, euthanized and necropsied. Clinical signs
21 included tongue protrusion, twitches and tremors of muscles of mastication, weight loss and
22 lethargy. Severe atrophy of the masseter, buccinator and lingual muscles was observed,
23 along with neuronal vacuolation in the nuclei of the trigeminal, facial, and hypoglossus
24 nerves. These findings and the clinical signs are consistent with results obtained in animals,
25 spontaneously and experimentally intoxicated with *Prosopis juliflora* in previous studies.
26 Several species of this genus are native to Argentina. Farmers should be warned about the
27 suspected toxicity by *Prosopis nigra*, since this species has wide geographical distribution
28 in the country.

29 **Keywords:** cattle; mesquite pods; neuronal damage; nervous signs; *Prosopis*-poisoning.

1. Introduction

The genus *Prosopis*, family *Leguminosae* (Fabaceae), subfamily *Mimosoideae*, is native to the Americas, Africa and Asia, and comprises a great number of species (Pasicznik et al., 2001). In Argentina, the species of this genus are known as “Algarrobos” (Verga et al., 2009). Its pods are a source of animal feed in many regions of the world due to their nutritional value (Silva et al., 1981; Silva et al., 1990; Riet-Correa et al., 2012). Despite this, spontaneous cases of intoxication have been observed in animals that remain in areas occupied by “algarrobos” for more than a year or longer than a fruiting period (Lima et al., 2004, Assis et al., 2009).

To date, all reports of toxicity by the genus *Prosopis* in animals have been associated with consumption of *P. juliflora* pods (Lima et al., 2004; Tabosa et al., 2004; Silva et al., 2006; Assis et al., 2009; Câmara et al., 2009). Spontaneous intoxication by *P. glandulosa* in goats was also reported (Washburn et al., 2002). Moreover, a spontaneous poisoning of goats by *Prosopis* sp. pods in Argentina has been recently reported (Micheloud et al., 2018).

The animals poisoned by *P. juliflora* pods exhibited oral dysphagia, masseter atrophy, tongue protrusion, mandible slackening, and progressive weight loss, leading to death (Câmara et al., 2009; Almeida et al., 2017). Experimental studies concluded that lesion in cranial nervous nuclei was the primary damage and that it induced the clinical signs (Tabosa et al., 2000; Almeida et al., 2017).

This paper describes five outbreaks of nervous disease in cattle associated with consumption of *Prosopis nigra* pods, including clinical and epidemiological aspects of the disease, and the gross and microscopic lesions identified.

2. Case report

54 2.1 *Clinical and epidemiological findings*

55 A diagnostic visit was made to five farms due to suspected signs of *P. nigra*
56 poisoning reported by the owners. Three of the affected herds were in common grazing
57 areas, located in the region of San Carlos in Salta province. The other two outbreaks
58 occurred in commercial breeding herds located in Tolombón (Salta province) and Frías
59 (Santiago del Estero province). The five holdings were dedicated to beef cattle production
60 based on grazing of natural pastures and fruits of native trees (Fig 1 A-D).

61 2.2 *Botanical identification*

62 In the grazing areas the species most closely associated with poisoning was
63 *Prosopis nigra*. The estimated consumption time, and incidence, mortality and lethality
64 data are shown in Table 1. Part of the plant material from grazing areas was collected and
65 submitted to herborization. All samples were identified as *Prosopis nigra* and recorded as
66 specimen MCNS 12880 at the MCNS Herbarium of National University of Salta. Data
67 about sampling sites, *P. nigra* tree/ha and relative abundance are presented in Table 2.
68 Abundance was calculated by counting the percentage of adult trees (*P. nigra*) per unit area
69 and is expressed as the average of the count of 10 randomly selected areas.

70 2.3 *Clinical findings*

71 Animals showing symptoms of intoxication were subjected to clinical examination,
72 which included evaluation of the posture, behaviour and spontaneous and induced
73 movements. The salient clinical signs consisted of weight loss and head tilting during
74 chewing; some of these animals had twitches and tremors of the masticatory muscles.
75 Decreased lingual tone was observed, and even 20-25% of the animals exhibited tongue
76 protrusion of about 5 and 10 cm (Fig. 2A, B). Abnormal movements of mastication and

77 notorious masseter atrophy were observed. Finally, the most severely affected animals had
78 total paralysis of the lower jaw and inability to close the mouth. Thus, the jaw was loose
79 and showed pendulous movements.

80 2.2. Pathological findings

81 Four of these animals were euthanized and subsequently necropsied during the visit.
82 At necropsy, general poor condition, serous fat atrophy in the omentum, and perirenal and
83 pericardial fat were observed, In addition, varying degrees of atrophy of the muscles of
84 mastication (masseter and temporal muscles) were identified (Table 3). In all cases, the
85 presence of *Prosopis* sp. seed in the rumen content was abundant. Later, fragments of
86 tissue, including the central nervous system and muscles of mastication, were collected in
87 10% buffered formalin, processed by routine histological techniques and stained with
88 hematoxylin and eosin (HE). Degenerative changes were observed, with cytoplasmic
89 hypereosinophilia, loss of striations and fragmentation of muscle fibres. In some muscle
90 fascicles, degeneration was so severe that muscle tissue was replaced with abundant
91 connective tissue and hyaline floccules. All cases had focal or diffuse mononuclear
92 infiltration of varying degrees. In the processed brain samples, focal gliosis was observed
93 with diffuse vacuolization and swelling of some neuronal bodies in the trigeminal, facial
94 and hypoglossus motor nuclei (Fig. 2.C, D).

95 3. Discussion

96 Diagnosis of intoxication by *Prosopis* sp. is based on epidemiological data, clinical
97 signs and pathological findings. The average time of consumption of *Prosopis nigra* pods
98 in the five outbreaks was estimated in 92 days. In bovines experimentally fed with *Prosopis*
99 *juliflora* pods, clinical signs were observed between days 45 and 75 after the beginning of

100 the experiment (Tabosa et al., 2006). Goats seem to be more resistant than cattle, since the
101 signs were observed after 240 days of exposure to diets with 60% of *P. juliflora* pods
102 (Tabosa et al., 2000). However, Almeida et al. (2017) showed that sheep became
103 intoxicated after consuming a diet composed of about 80% of *Prosopis* for 21 months (>
104 600 days), suggesting that sheep are much more resistant to poisoning than cattle and goats.
105 The epidemiological data (incidence, mortality and lethality) were similar to those reported
106 elsewhere for a natural outbreak of *P. juliflora* poisoning in cattle (Câmara et al., 2009,
107 Kingsbury 1964).

108 All the observed outbreaks occurred at sites where *P. nigra* was the most abundant
109 species (90% approximately), with a very high density (106.8 adult trees/ha). This plant
110 density ensures a good supply of pods for the animals. This assumption agrees with
111 findings obtained by experimentally poisoning bovines with *P. juliflora* pods (Tabosa et al.,
112 2006); in that study, the disease was evident when the supply of pod exceeded 60% for a
113 prolonged period.

114 In the present study, one of the early clinical signs was faint tremors of the lower
115 jaw, which progressed to severe disorders characterized by rapid mastication and alterations
116 in chewing movements. In more advanced stages of intoxication, evidence of masseter
117 muscle atrophy, weight loss and lethargy was observed. Tongue protrusion was observed in
118 20% of the animals, which may have been due to paralysis of the lower jaw. All the clinical
119 signs described are in agreement with information obtained from cattle naturally and
120 experimentally intoxicated with pods of *P. juliflora* (Figueiredo et al. 1995; Tabosa et al.,
121 2006; Câmara et al., 2009).

122 Histopathological studies of the masseter, buccinator and lingual muscles revealed
123 different atrophy degrees. These lesions can explain the disorders in chewing and

124 swallowing in affected animals; and muscle atrophy is, most probably, due to disorders of
125 innervation, which were confirmed through lesions found in nuclei of the trigeminal,
126 hypoglossal and facial nerves. These lesions were described in cattle (Tabosa et al., 2006;
127 Câmara et al., 2009), sheep (Almeida et al., 2017) and goat (Tabosa et al., 2000) intoxicated
128 with *Prosopis juliflora*.

129 Neuropathic muscular atrophy has been reported as one of the most common causes
130 of muscle atrophy in animals (Jubb et al., 2007; Valberg 2010). Muscles fibres require a
131 neural stimulation of low tone for maintenance; when there is no neural stimulation, the
132 fibres get retracted and can suffer degeneration and pyknosis (Valberg, 2010). All the
133 findings reported here are consistent with those mentioned by Tabosa et al. (2006) in
134 experimental poisoning with *Prosopis juliflora* in Brazil.

135 *P. juliflora* contains piperidine alkaloids, such as juliflorine, julifloricine,
136 julifloridine, juliprosinene, juliflorinine and juliprosopine (Ahmad et al., 1986, Ahmad et
137 al., 1989). According to those authors, these alkaloids are responsible for producing the
138 damage of the mitochondria in neurons of central nuclei (Maioli et al., 2012; Silva et al.,
139 2013). It seems reasonable to speculate that these alkaloids are also the toxic compounds in
140 *P. nigra*. However, the toxic components were not isolated from the plant in this study.
141 Thus, further studies are needed to confirm this hypothesis and to identify the specific toxin
142 and its concentration in the plant. Despite the positive effects of pods as source of food in
143 animals (Riet-Correa et al., 2012), these results suggest that consumption, in large
144 quantities and over a long period can be dangerous. The findings described suggest that the
145 consumption of pods of native *Prosopis* species (*P. nigra*) produces the nervous damage
146 and clinical signs, as observed in cattle poisoning by *Prosopis juliflora* in others countries.

147 To date, the only *Prosopis* species reported as toxic have been *P. juliflora* (Tabosa
148 et al., 2006, Câmara et al., 2009) and *P. glandulosa* (Washburn et al., 2002). In Argentina,
149 there are 47 native species of this genus (Flora Argentina, 2017). Farmers should be warned
150 about the suspicion of the toxicity of this species, since *P. nigra* has wide geographical
151 distribution in the country.

152

153 **CONFLICT OF INTEREST**

154 No competing financial interest exists for any of the coauthors of this manuscript.

155 **ACKNOWLEDGEMENTS**

156 This study was supported by grants from INTA (Instituto Nacional de Tecnología
157 Agropecuaria) and Consejo de Investigación de la Universidad Católica de Salta. LACH
158 and EJJ are members of CONICET (Consejo Nacional e Investigaciones Científicas y
159 Técnicas). We thank Dr Rubén Bravo and Dr Rafael Piedrasanta for their contribution to
160 field work during outbreaks.

161 **REFERENCES**

- 162 Ahmad A., Ali Khan K., Ahmad V.U., Qazi, S. 1986. Antibacterial activity of juliflorine
163 isolated from *Prosopis juliflora*. *Planta Medica*, 1, 285-288.
- 164 Ahmad V.U., Sultana A., Qazi, S. 1989. Alkaloids from the leaves of *Prosopis juliflora*. *J.*
165 *Nat. Prod.*, 52, 497-501.
- 166 Almeida V. M., Rocha B. P., Pfister J. A., Medeiros R.M., Riet-Correa F., Chaves H.A.,
167 Silva Filho G.B., Mendonça F.S. 2017. Spontaneous poisoning by *Prosopis juliflora*
168 (Leguminosae) in sheep. *Pesq. Vet. Bras.*, 37, 110-114.

- 169 Assis, T.S., Medeiros, R.M.T., Riet-Correa, F., Araújo, J.A.S., Dantas, A.F.M. 2009.
170 Intoxicações por plantas em ruminantes e equídeos no sertão paraibano. *Pesq. Vet. Bras.*,
171 29, 919-924.
- 172 Câmara, A.C.L., Costa, N.A., Riet-Correa, F., Afonso, J.A.B., Dantas, A.F.M., Mendonça,
173 C.L., Souza, M.I. 2009. Intoxicação espontânea por vagens de *Prosopis juliflora* (Leg.
174 Mimosoideae) em bovinos em Pernambuco. *Pesq. Vet. Bras.*, 29, 233-240.
- 175 Jubb, Kennedy, Palmer's. 2007. *Pathology of Domestic Animals*, 5th ed., Saunders Elsevier,
176 Philadelphia.
- 177 Figueiredo L.J.C., Ferreira M.M., Távora J.P.F., Dantas J. & Simões S.D. 1995. Estudo
178 clínico e anatomopatológico da doença "cara torta" em bovinos no nordeste brasileiro. *Arq.*
179 *Med. Vet. UFBA* 18:175-183.
- 180 Flora Argentina. 2017. *Plantas Vasculares de la República Argentina*.
181 www.floraargentina.edu.ar [consulted in September 2018].
- 182 Kingsbury J.M. 1964. *Poisonous Plants of the United States and Canada*. Prentice-Hall,
183 Englewood Cliffs, New Jersey, pp.349-351.
- 184 Lima E., Riet-Correa F., Amorin S.L., Sucupira Júnior G. 2004. Intoxicação por favas de
185 *Prosopis juliflora* (algaroba) em caprinos no Nordeste do Brasil. *Pesq. Vet. Bras.*, 24, 36-
186 37.
- 187 Maioli, M.A., Lemos D.E.C.V., Guelfi, M., Medeiros, H.C.D., Riet-Correa, F., Medeiros,
188 R.M.T., Barbosa-Filho, J.M., Mingatto, F.E. 2012. Mechanism for the uncoupling of
189 oxidative phosphorylation by juliprosopine on rat brain mitochondria. *Toxicon*, 60, 1355-
190 1362.

- 191 Micheloud, J.F., Vera, T.A., Caro, L.C., Gimeno, E.J. 2018. Clinico-pathological findings
192 in natural cases of “mascadera” in goats. Trop. Anim. Health. Prod., 50, 441-444.
- 193 Pasiecznick, N.M., Felker, P., Harris, P.J.C., Harsh, L. N., Cruz, G., Tewari, J. C., Cadoret,
194 K., Maldonado, L.J. 2001. The *Prosopis juliflora*- *Prosopis pallida* Complex. A
195 monograph. HDRA Coventry, UK pp. 162.
- 196 Riet-Correa, F., Andrade, F.R.M., Carvalho, F.K.L., Tabosa, I.M., Galiza, G.J., Bernardino,
197 N., Simões, S.V.D. & Medeiros, R.M.T. 2012. Use of *Prosopis juliflora* pods as food for
198 sheep and goats. Pesq. Vet. Bras., 32, 987-989.
- 199 Silva, D.S. 1981. Substituição progressiva do farelo de trigo pela vagem da algaroba na
200 alimentação de bovinos em engorda. Universidade Federal da Paraíba; Areia (PB).
- 201 Silva, A. M., Rodrigues, M. E., De Silva, J.F. 1990. Nutritive value of mesquite beans
202 (*Prosopis juliflora*) in the diet of rabbit. Vet. Zoot., 2, 9-16.
- 203 Silva, D.M., Riet-Correa, F., Medeiros, R. M. T., Oliveira O.D. 2006. Oliveira. Plantas
204 tóxicas para ruminantes e eqüídeos no Seridó Ocidental e Oriental do Rio Grande do Norte.
205 Pesq. Vet. Bras., 26, 223-236.
- 206 Silva, V. D. A., Pitanga, B. P., Nascimento, R. P., Souza, C. S., Coelho, P. L. C., Menezes-
207 Filho, N., Silva A.M., Costa M de F, El-Bachá R.S., Velozo E.S., Costa, S. L. 2013
208 Juliprosopine and Juliprosine from *Prosopis juliflora* leaves induce mitochondrial damage
209 and cytoplasmic vacuolation on cocultured glial cells and neurons. Chem. Res. Toxicol, 26,
210 1810-1820.

- 211 Tabosa, I.M., Souza, J.C.A., Barbosa-Filho, J.M., Almeida, R.N., Graça, D.L., Riet-
212 Correa, F. 2000. Neuronal vacuolation of the trigeminal nuclei in goats caused by the
213 ingestion of *Prosopis juliflora* pods (mesquite Supply of). *Vet Hum Toxicol*, 42, 155–158.
- 214 Tabosa, I.M., Riet-Correa F., Simões S.V.D., Medeiros R.M.T., Nobre V.M.T. 2004.
215 Intoxication by *Prosopis juliflora* pods (mesquite beans) in cattle and goats in Northeastern
216 Brazil, in: T. Acamovic, C.S. Stewart, T.W. Pannycott (Eds.), *Toxic Plants and other*
217 *Natural Toxicants*. CAB International Publishing, Wallingford, Oxon, UK. pp. 341-346.
- 218 Tabosa, I. M., Riet-Correa, F., Barros, S. S., Summers, B. A., Simões, S. V. D., Medeiros,
219 R. M. T., Nobre, V. M. T. 2006. Neurohistologic and ultrastructural lesions in cattle
220 experimentally intoxicated with the plant *Prosopis juliflora*. *J Vet Pathol* , 43, 695-701.
- 221 Valberg, S. J. 2010. Enfermedades Musculares, fourth ed, in: B.P. Smith (Ed.), *Medicina*
222 *interna de los grandes animales*. Barcelona, Spain, pp.1388-1418.
- 223 Verga A., López Lauenstein D., López C., Navall M., Joseau J., Gómez C., Royo O.,
224 Degano W., Marcó M. 2009. Caracterización morfológica de los algarrobos (*Prosopis sp.*)
225 en las regiones fitogeográficas Chaqueña y Espinal norte de Argentina. *Quebracho*, 17, 31-
226 40.
- 227 Washburn, K. E., Breshears, M. A., Ritchey, J. W., Morgan, S. E., Streeter, R. N. 2002.
228 Honey mesquite toxicosis in a goat. *J Am Vet Med Assoc*, 220, 1837-1839.
- 229

Figure legends

Fig. 1. A- Grazing areas, locally characterized as "algarrobales", where the dominant species was *P. nigra*. B- Detail of an adult specimen of *P. nigra*. C and D- Inflorescence and fruits of the plant.

Fig. 2. A- Bovine affected by the disease showing partial protrusion of tongue. B- Adult cow tilting the head to prehend the forage. Microscopic image: C- Fibers of masseter muscles, showing degeneration and fragmentation. Abundant mononuclear infiltration is also observed between fibers (H&E 20X). D- Vacuolation and degenerative changes in neurons (insert) of the trigeminal nucleus (H&E 20X).

Table 1. Epidemiological data (mortality, incidence and lethality) of outbreaks.

Outbreak	Estimated time of <i>P. nigra</i> pod consumption (days)	Incidence (%)	Mortality (%)	Lethality (%)
1	100	15% (9/60)	6.6 % (4/60)	44% (4/9)
2	100	21% (8/38)	13% (5/38)	62% (5/8)
3	100	12.5% (1/8)	12.5% (1/8)	100% (1/1)
4	75	40 % (60/150)	40% (60/150)	100 % (60/60)
5	85	20 (74/300)	20 (70/300)	100% (70/74)
average	92	21.7	18.42	81.2

Table 2. Outbreak sites, *P. nigra* density (adult trees /ha) and relative abundance.

Outbreak	Location	GPS	<i>P. nigra</i> trees /ha	Adult (<i>P. nigra</i>) trees per unit area (%)
1	San Carlos (SAL)*	25°57'15.49"S 65°56'51.82"W	80	88%
2	San Carlos (SAL)	25°55'44.26"S 65°57'20.82"W	75	92%
3	San Carlos (SAL)	25°56'7.71"S 65°57'10.96"W	150	90%
4	Tolombón (SAL)	26°19'10.51"S 65°57'27.57"W	135	99%
5	Frías (SDE)*	28°42'2.59"S 65° 3'46.38"W	94	85%
average			106.8	90.8%

*Salta province (SAL)




*Santiago del Estero province (SDE)

Table 3. Degrees of muscle atrophy in HE-stained tissue sections from cattle spontaneously poisoned with *Prosopis nigra*

Necropsies	Levels of atrophy		
	Masseter muscle	Buccinators muscle	Extrinsic tongue muscles
1	++++	+++	++
2	+++	+	+
3	++++	++	+++
4	++++	++	+

- = No lesion + = Minors lesions ++ = moderate lesions +++ = severe lesions

A**B****C****D**



Highlights

- First report providing local information on outbreaks of poisoning by *P. nigra* pods in cattle.
- The macroscopic and microscopic findings were consistent with those observed in poisoning by other *Prosopis* species.
- *P. nigra* has wide geographical distribution in the Argentina, Paraguay, Brazil and Uruguay region.