



ISSR markers to explore entomopathogenic fungi genetic diversity: Implications for biological control of tobacco pests

MARIA FLORENCIA VIANNA¹, SEBASTIAN PELIZZA¹, MARÍA LETICIA RUSSO^{1*} ,
ANDREA TOLEDO², CECILIA MOURELOS² and ANA CLARA SCORSETTI¹

¹Instituto Spegazzini, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina

²Centro de Investigaciones en Fitopatología, Universidad Nacional de La Plata,
La Plata, Buenos Aires, Argentina

*Corresponding author (Email, leticia.russo@conicet.gov.ar)

MS received 15 March 2020; accepted 6 October 2020

Tobacco is one of the major industrial crops cultivated worldwide. Chemical control is the main method employed to reduce damage by insect pests. The use of entomopathogenic fungi represents an alternative to replace insecticides. The search for effective strains in the field constitutes a first step when developing a formulation. The objective of this work was to study genetic differences among isolates of entomopathogenic fungi obtained from tobacco grown soils using ISSR markers. The pathogenicity of the strains towards *Helicoverpa gelotopoeon* and *Diabrotica speciosa* was also assessed in order to search for a relationship between virulence and genetic diversity. Nineteen isolates were identified according to morphological features and molecular techniques as *Beauveria bassiana* (11) and *Purpureocillium lilacinum* (8). The diversity tree generated by ISSR analysis showed a high diversity among the strains. The pathogenicity towards *H. gelotopoeon* and *D. speciosa* was assessed and the logistic models generated showed that *B. bassiana* isolates LPSc1215 and LPSc1364 were the most pathogenic against both insect pests tested. In the diversity tree, these strains were grouped in a same cluster with a similarity level of approximately 85%, indicating a possible relationship between virulence and the band pattern generated.

Keywords. *Beauveria bassiana*; *Diabrotica speciosa*; genetic polymorphism; *Helicoverpa gelotopoeon*; pathogenicity; *Purpureocillium lilacinum*

1. Introduction

Tobacco (*Nicotiana tabacum* L.) is one of the major industrial crops, with a cultivated area of approximately 3.9 million hectares worldwide. Many insect pests attack this crop at different stages of its life cycle, affecting the development of the plant and therefore decreasing crop yield. The most harmful insect affecting this crop are those belonging to Lepidoptera and Coleoptera (Mercado Cardenas *et al.* 2013; Yang *et al.* 2019). The main method to reduce damage is the use of chemical insecticides that seriously affect the environment and human health, especially of farmers that are in direct contact with them (Da Silva *et al.* 2014).

Biological control has arisen to amend adverse effects of conventional plant protection strategies,

particularly fungal entomopathogens are promising organisms among these ecofriendly practices. Species within the genera *Beauveria*, *Lecanicillium*, *Metarhizium* and *Purpureocillium* have been commercially formulated with more than 200 products available in the market (de Barros *et al.* 2015; de Faria and Wraight 2007; Mishra *et al.* 2015; Pathan *et al.* 2019; Sandhu *et al.* 2017).

Accurate taxonomic determination and knowledge about genetic diversity of strains constitutes the first step when searching for a suitable biocontrol agent (Fernández Bravo 2017; Zimmermann 2007). In this sense, a promising application of fingerprints techniques is that they may allow tracing organisms in the field when these are utilized as formulations with biological control purposes and therefore the way strains perform and persist

on the field (Takatsuka 2007). Tigano-Milani *et al.* (1995) proposed the hypothesis that more than one haplotype may be required to initiate and maintain a fungal epizootic in an insect population. According to this, the genetic diversity of the strains would be the factor that would allow the fungi to adapt to changing environmental conditions and to successfully attack different insect populations (Cruz *et al.* 2006). In line with this, PCR molecular markers such as Amplified Fragment Length Polymorphism (RFLP), Random Amplified Polymorphic DNA (RAPD) and Inter Simple Sequence Repeats (ISSR) have been used as effective tools to detect genetic variability among fungal populations (Enkerli and Widmer 2010; Fernandes *et al.* 2006; Serna-Domínguez *et al.* 2019; Sevím *et al.* 2012; Trissi *et al.* 2013). DNA fingerprinting with ISSR markers is considered to provide superior discrimination among fungal isolates compared to other methods because it produces quite variable fingerprints with a large number of fragments throughout the genome (Taylor *et al.* 1999). Concerning fungal entomopathogens, ISSR markers have been successfully used to detect genetic variability within several species of Hypocreales like *Beauveria bassiana* (Bals.-Criv.) Vuill. (Estrada *et al.* 2007; Rehner and Buckley 2005; Takatsuka 2007; Toledo *et al.* 2019), *Metarhizium anisopliae* (Metsch.) Sorokin (Enkerli *et al.* 2005; Oulevey *et al.* 2009) and *Paecilomyces fumosoroseus* (= *Isaria fumosorosea*) Wize (Dalleau-Clouet *et al.* 2005), constituting a first step in the selection of pathogenic strains according to their genotype.

The objective of this investigation was to study genetic differences among entomopathogenic fungi associated to tobacco crop using ISSR markers and to determine their entomocidal capacity against to major tobacco pests, *Helicoverpa gelotopoeon* Dyar (Lepidoptera: Noctuidae) and *Diabrotica speciosa* Germar (Coleoptera: Chrysomelidae) under laboratory conditions in order to establish a connection between fungal genetic profiles and their capacity to kill insects.

2. Material and methods

2.1 Fungal isolation and morphological characterization

Fungal isolates were obtained from soil samples from seven tobacco (*N. tabacum* L.) crops. Samples were collected during summer in Perico, Jujuy province (24°23'S – 65°08'W). Five compound soil samples from each crop (from a depth up to 12 cm) were collected with a sterile shovel, placed into

polyethylene plastic bags and stored at 4°C in the laboratory for no longer than a month until they were examined for the presence of entomopathogenic fungi.

To accomplish entomopathogenic fungi isolation from soil, serial dilution protocol was performed according to Lecuona (1996). For each sample, six dilutions with five repetitions each were obtained and 100 µl were spread with a Drigalsky spreader in Petri dishes containing selective medium for entomopathogenic fungi (Posadas *et al.* 2012). A total of 1,050 plates were analyzed for the presence of entomopathogenic fungi. These were stored at 24°C in the darkness for a period of 10 to 15 days. Entomopathogenic fungal colonies were detected under binocular microscope and transferred to potato dextrose agar (PDA) (Britania® S.A., Buenos Aires, Argentina) plates. Microscopical mounts were made to confirm fungal morphological identity according to taxonomic keys and monographs of Nagamani *et al.* (2006); Domsch *et al.* (2003); Samson *et al.* (1988); Seifert *et al.* (2011). For each isolate, monosporic colonies were obtained and preserved at the Micological Collection of the Botanical Institute Carlos Spegazzini, Argentina (LPSc), under their respective accession numbers.

2.2 Diversity analysis

DNA was extracted from eleven strains of *B. bassiana* and eight strains of *Purpureocillium lilacinum* (Thom) Luangsa-ard, Houbraken, Hywel-Jones & Samson grown on liquid Sabouraud dextrose plus 1% yeast extract (SDY 1%) medium after 7 day's incubation, without agitation, at 25°C in the darkness. Extraction was performed according the methodology proposed by Aljanabi and Martinez (1997). DNA quality was assessed by electrophoresis on 0.7% w/v agarose gels supplemented with ethidium bromide (100 ng ml⁻¹) and its quantity was estimated by comparison with a molecular marker control of known concentration (Lambda Phage Genome digested with Hind III – Promega Biotech) using the image analyzer (SYNGENE GeneTools).

Diversity analysis was performed using the ISSR primers 826 (5'-ACCACACACACACC-3'), BA3 (5'-ACACACACACACACACCT-3') and KA5 (5'-CTACACACACACACACAC-3') selected considering previous results obtained by Toledo *et al.* (2019). The ISSR-PCR reactions were performed on a MJ Research (PTC-150 MiniCycler) and each amplification reaction

had a total volume of 15 μ l containing 1 \times buffer (500 mM KCl; 100 mM Tris-HCl, pH 9.0 at 25°C; 1% Triton X-100), 2 mM (primers 826 and KA5) or 2.5 mM (primer BA3) of MgCl₂, 0.25 mM each dNTP, 50 ng of each primer, 1 U Taq DNA polymerase and approximately 50 ng of template DNA (reagents Inbio Highway®, Tandil, Buenos Aires, Argentina). PCR program was as follows: initial denaturation at 94°C for 5 min, followed by 30 cycles of denaturation at 94°C for 45 s, annealing at 48°C for 45 s and extension at 72°C for 3 min, with an additional final extension step at 72°C for 5 min. PCR products were resolved by electrophoresis on 1.5% w/v agarose gels supplemented with ethidium bromide (100 ng ml⁻¹) using as control an internal standard the DNA marker 100–1000 bp (Inbio Highway®, Tandil, Buenos Aires, Argentina). The ISSR banding patterns generated were analyzed with the software Past3 and a dendrogram was built using the similarity matrix of DICE and the Unweighted Pair Group Method with Arithmetic Mean (UPGMA).

2.3 Molecular identification

For further isolates identification, their molecular identity was confirmed by choosing eight isolates to amplify and sequence the ribosomal internal transcribed spacers (ITS) and the translation elongation factor 1-alpha (TEF1- α) genes. The isolates were selected according to their location in the previously built diversity analysis dendrogram.

DNA extraction was performed as describe above for the diversity analysis. The universal primers ITS5 (5'-GGAAGTAAAAGTCGTAACAAGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White *et al.* 1990) and primers 983F (5'-GCYCCYGGH-CAYCGTGAYTTYAT-3') and 2218R (5'- ATGACACCRACRGCACRGTGTG-3') (Rehner and Buckley 2005) were used to amplify ITS and TEF1- α regions, respectively. The reaction mix contained Buffer 1 \times (500 mM KCl; 100 mM Tris-HCl, pH 9.0 at 25°C; 1% Triton X-100), 1.5 mM of MgCl₂, 0.2 mM of each dNTP, 50 ng of each primer, 1 U of Taq DNA polymerase and approximately 20 ng of DNA (Inbio Highway®, Tandil, Buenos Aires, Argentina reagents). The PCR reactions were performed in a MJ Research (PTC-150 MiniCycler) termocycler, and the program consisted in an initial DNA denaturation at 94°C for 4 min followed by 33 cycles of denaturation at 94°C for 1 min, 45 s at 56°C for ITS and 1 min at 48°C for TEF1- α , 1 min at 72°C and finally an elongation step at

72°C for 5 min. The amplified PCR products were precipitated in an isopropanol volume and a 0.1 volume of ammonium acetate 3M, this mixture was incubated at -18°C for 12 h and later was centrifuged at 15,000g for 15 min. The DNA pellet was washed with ethanol 70%, dried and dissolved in distilled sterile water. The sequencing was performed at the CERELA-Institute (CONICET, Tucumán, Argentina). The sequences obtained were submitted to the National Center for Biotechnology Information (NCBI) GenBank database (www.ncbi.nlm.nih.gov) for gene annotation. Sequences were edited using BioEdit software version 7.0.5 (Hall 1999), and posteriorly used to perform a phylogenetic analysis that included the sequences of the isolates of *Beauveria* and *Purpureocillium* from this work with sequences of some species for the genera available at the GenBank database including type material of *B. bassiana* ITS= NR_111594.1, TEF1- α = HQ880974.1 and *P. lilacinum* ITS= NR_111432.1, TEF1- α = EF468792.1. All sequences were aligned with the ClustalW tool of the program Mega5 (Tamura *et al.* 2011) and the phylogenetic study was carried out using the analysis of maximum likelihood. The statistical support for the nodes was evaluated through 1000 bootstrap replicates.

2.4 Pathogenicity tests

Pathogenicity bioassays were performed against third instar larvae of *H. gelotopoeon* and adults of *D. speciosa*. *Helicoverpa gelotopoeon* eggs were provided by AgIdea Pergamino, Buenos Aires, Argentina (<https://www.agidea.com.ar/>) and larvae were fed with artificial diet according to Patana (1977). *Diabrotica speciosa* adults were collected from the field and were kept in laboratory for a period of fifteen days to prevent any disease from the field (Pilz *et al.* 2007).

All entomopathogenic strains were tested for their pathogenicity. Each fungal colony was grown on PDA at 25°C in the darkness during 14 days. After two weeks, 10 ml of Tween 80 (Merck®) 0.01 % v/v were added to Petri plates to obtain conidia by scraping strains surfaces with a sterile scalpel. The resulting conidial suspensions were filtered with a sterile cloth and subsequently were vortexed for five minutes. Conidial concentration of each suspension was adjusted to 1 \times 10⁸ conidia/ml using a Neubauer chamber. Conidial viability was assed according to Ayala Zermeño *et al.* (2015).

To test strains pathogenicity the immersion method was carried out according to Nussenbaum and Lecuona (2012). Three replicates of 10 individuals for each

treatment were performed. Insects were submerged individually for fifteen seconds in conidial suspensions, subsequently were lean on sterile filter paper to dry off excess moisture, and finally placed on plastic Petri dishes. Controls were treated only with Tween 80. Insects were kept on a breeding chamber under uniform and controlled conditions ($25 \pm 2^\circ\text{C}$, 75% RH, photoperiod 14:10 L: D). Food was provided on daily bases, for *H. gelotopoeon* caterpillars artificial diet according to Patana (1977) was used and for *D. speciosa* adults, zucchini (*Cucurbita pepo*) pieces of approximately 0.5 cm^3 were offered (Pecchioni 1989).

The number of dead insects on each treatment was recorded for a period of fourteen days. Mortality due to mycosis was asserted by placing cadavers on humid chambers and stored at 25°C in the darkness to promote mycelia outgrowth (Powell et al. 2007).

Mortality registered for each strain was adjusted to a logistic model to estimate their pathogenicity as a function of time using StatSoft Statistica software (version 7.1) (2005).

3. Results

3.1 Fungal isolates

A total of nineteen entomopathogenic fungal isolates were obtained from tobacco grown soil samples using selective media for this type of fungi. According to morphological features, eleven of them were assigned to *B. bassiana* (Accession numbers: LPSc1210, LPSc1211, LPSc1212, LPSc1213, LPSc1214, LPSc1215, LPSc1216, LPSc1217, LPSc1363, LPSc1364, LPSc1364, and LPSc1265) and eight to *P. lilacinum* (Accession numbers: LPSc1367, LPSc1368, LPSc1369, LPSc1370, LPSc1371, LPSc1372, LPSc1373, and LPSc1374).

3.2 Diversity analysis

The diversity study employing three ISSR primers generated a total of 55 fragments with sizes that ranged between 300 and 1200 pb. The primer 826 generated a pattern of 18 bands, the primer BA3 a pattern of 13 bands and the primer KA5 a pattern of 24 bands (figure 1). The topology of the UPGMA-tree showed a high diversity among the strains. The three primers used confirmed the presence of four clusters inside *B. bassiana* group, showing the existence of at least four different haplotypes. Isolate LPSc1363 appeared in the first clade. Isolates LPSc1214,

LPSc1213, LPSc1211, LPSc1212 and LPSc1365 were associated in the second clade, in which the first three isolates were grouped in a subclade and the other two isolates in another subclade. Isolates LPSc1215 and LPSc1364 formed the third clade, while isolates LPSc1217, LPSc1216 and LPSc1210 were grouped in the fourth clade. Similar relationship was observed inside the *P. lilacinum* group, where four different haplotypes were also recorded. The first clade was formed by isolates LPSc1368, LPSc1369, LPSc1370 and LPSc1367, with the two first isolates grouped in a subclade. The second clade was formed by isolates LPSc1372 and LPSc1373, while the third and the fourth clades were formed by isolates LPSc1374 and LPSc1371, respectively (figure 1).

3.3 Molecular identification

ITS amplicons from the eight selected strains (*B. bassiana* LPSc1213, LPSc1216, LPSc1363 and LPSc1364 and *P. lilacinum* LPSc1370, LPSc1371, LPSc1373 and LPSc1374) considered as representative for each cluster in the ISSR dendrogram had a length of 522 to 576 bp for *B. bassiana* and of 474 to 486 bp for *P. lilacinum*. TEF1- α amplicons of the same selected isolates had a size of 954 to 956 bp for *B. bassiana* and of 917 to 920 bp for *P. lilacinum*. In the maximum likelihood tree resulting from the phylogenetic analysis of the ITS region, the four isolates of *B. bassiana* were clustered together with the material type of *B. bassiana* NR_111594 in a monophyletic group supported by a bootstrap value of 99%. While *Beauveria amorphia*, *B. caledonica*, *B. malawiensis* and *Cordyceps brongniartii*, all morphologically-related species, formed an independent clade supported by a bootstrap value of 95% (figure 2A). In the same tree the *P. lilacinum* isolates selected were clustered in a monophyletic group supported by a bootstrap value of 99%, which included only *P. lilacinum* representatives, including the sequences of the type material NR_111432.1 (figure 2A). On the other hand, the maximum likelihood tree resulting from the phylogenetic analysis of the TEF1- α showed a similar topology to that observed for ITS regions. All isolates of *B. bassiana* selected in this work were clustered together with *B. bassiana* type material HQ880974.1 in a same clade supported by a bootstrap value of 82%. Likewise, the four isolates of *P. lilacinum* were grouped together with the type material EF468792.1 in a clade integrated only by this species and supported by a bootstrap value of 100% (figure 2B).




Figure 1. Dendrogram built using the similarity matrix of DICE and the UPGMA including the patterns of bands of *Beauveria bassiana* and *Purpureocillium lilacinum* isolates generated by ISSR primers 826, BA3 and KA5. M = DNA marker 100–1000 bp.

3.4 Pathogenicity tests

All isolates tested against *H. gelotopoeon* were adjusted to a logistic model for their pathogenicity as a function of time. Beta values were considered as 0, and p values for all models showed values <0.005 (table 1

and figure 3). Those isolates that were not pathogenic to larvae were not adjusted to any model.

The strains LPSc1212, LPSc1214, LPSc1215, LPSc1364 and LPSc1365 belonging to *B. bassiana* exhibited pathogenicity levels above 50%. The isolate LPSc1215 was the most pathogenic to the insect with




Figure 2. Phylogenetic tree obtained by the analysis of (A) ITS and (B) TEF1- α sequences from four *Beauveria bassiana* isolates and four *Purpureocillium lilacinum* and related groups. Isolates belonging to this work are identified by an * symbol. GeneBank accession numbers of the isolates are provided next to each strain. The bar at the bottom indicates the number of substitutions per site.

100% infectivity at 7 days after treatment. Also, isolates LPSc1212 and LPSc1364 showed high levels of pathogenicity with 97% and 94% at 11 and 10 days post treatment, respectively. With respect to *P. lilacinum* isolates, only LPSc1371 was pathogenic against *H. gelotopoeon* larvae, showing a low pathogenicity of 33% after 7 days post treatment. In addition, a logistic model was chosen as well to analyze the behavior (pathogenicity as a function of time) of isolates towards *D. speciosa*. Beta values were considered as 0, and p values for all models showed values

<0.005. Alfa and Gama values for the models generated for each strain are presented in table 1 and figure 4. If strains were not pathogenic, no model was adjusted. Only four isolates displayed pathogenicity levels over 50% against *D. speciosa*. The isolate *B. bassiana* LPSc1215 was the most pathogenic to the insects reaching 98% mortality at 11 days while isolate *B. bassiana* LPSc1364 reached 86% mortality at 12 days. Likewise, *P. lilacinum* isolates LPSc1368 and LPSc1369 showed 81% and 78% mortality at 10 and 9 days, respectively.

Table 1. Pathogenicity of *Beauveria bassiana* and *Purpureocillium lilacinum* towards *Helicoverpa gelotopoeon* and *Diabrotica speciosa*

Species	Strain	<i>Helicoverpa gelotopoeon</i>			<i>Diabrotica speciosa</i>		
		Maximum pathogenicity reached	Pathogenicity rate	Maximum pathogenicity time	Maximum pathogenicity reached	Pathogenicity rate	Maximum pathogenicity time
<i>B. bassiana</i>	LPSc 1365	0.64	1.09	10	0.16	1.6	8
<i>B. bassiana</i>	LPSc 1212	0.97	0.6	11	0.2	1.46	10
<i>B. bassiana</i>	LPSc 1217	0.2	1.48	7	0	0	0
<i>B. bassiana</i>	LPSc 1214	0.54	1.05	10	0.07	1.52	8
<i>B. bassiana</i>	LPSc 1216	0	0	0	0	0	0
<i>B. bassiana</i>	LPSc 1213	0.37	1.15	12	0.48	1.02	10
<i>B. bassiana</i>	LPSc 1215	1.02	1.08	6	0.98	1.43	11
<i>B. bassiana</i>	LPSc 1210	0.13	1.46	9	0.1	1.52	11
<i>B. bassiana</i>	LPSc 1211	0.31	1.31	10	0.34	1.28	9
<i>B. bassiana</i>	LPSc 1363	0.24	1.64	5	0	0	0
<i>B. bassiana</i>	LPSc 1364	0.94	1.32	10	0.86	1.29	12
<i>P. lilacinum</i>	LPSc 1368	0	0	0	0.81	0.96	10
<i>P. lilacinum</i>	LPSc 1369	0	0	0	0.78	1.34	9
<i>P. lilacinum</i>	LPSc 1370	0	0	0	0.17	1.3	7
<i>P. lilacinum</i>	LPSc 1367	0	0	0	0.24	1.44	9
<i>P. lilacinum</i>	LPSc 1371	0.33	1.07	7	0	0	0
<i>P. lilacinum</i>	LPSc 1372	0	0	0	0.07	1.38	8
<i>P. lilacinum</i>	LPSc 1373	0	0	0	0	0	0
<i>P. lilacinum</i>	LPSc 1374	0	0	0	0.13	1.11*	11

4. Discussion

In the present study, soil samples associated to *N. tabacum* crops were analyzed for the presence of entomopathogenic fungi. A total of 19 isolates were identified according to morphological features and molecular techniques as *B. bassiana* (11) and *P. lilacinum* (8). Both species are frequently isolated from natural and agricultural soils (Vega 2008) where their

survival is influenced by various factors such as UV radiation, air mobility, water content, temperature (Flint *et al.* 2002; Thompson *et al.* 2006; Yang *et al.* 2019), fertilizers and other chemical inputs (Bruck 2009, Shapiro-Ilan *et al.* 2013), processing methods (Boetel *et al.* 2012; Hummel *et al.* 2015) and soil and crop types (Shapiro-Ilan *et al.* 2012). The results of our investigations revealed the presence of two of the most effective biological control species among




Figure 3. Accumulated mortality (%) as a function of time (days) of *Helicoverpa gelotopoeon* larvae after inoculation of different strains of *Beauveria bassiana* and *Purpureocillium lilacinum*.




Figure 4. Accumulated mortality (%) as a function of time (days) of *Diabrotica speciosa* adults after inoculation of different strains of *B. bassiana* and *P. lilacinum*.

entomopathogenic fungi, capable to survive in tobacco soils in Jujuy province and constituting potential agents for the control of pests in this crop.

The diversity tree generated by ISSR analysis allowed to separate strains of *B. bassiana* and *P. lilacinum*, but also permitted to discriminate isolates within the same species. Studies by Takatsuka (2007), Wang *et al.* (2005) and Estrada *et al.* (2007) also demonstrated that this type of technique provides valuable information when discriminating isolates of *B. bassiana*. Furthermore, Aquino de Muro *et al.* (2005) who studied *Beauveria* species isolated from Middle East and West Asia, established that ISSR and AFLP analyses allowed correlating intraspecific groupings with genetic diversity between strains, recognizing then that this kind of methodologies are quick and reliable when describing genetic diversity of entomopathogenic fungi. In this study, three ISSR markers were used to generate banding patterns that enabled characterizing strains diversity among *B. bassiana* and *P. lilacinum*. Usually, the number of markers employed is higher but only a few of them are useful to discriminate between isolates (Estrada *et al.* 2007; Fontecha *et al.* 2011; Takatsuka 2007; Wang *et al.* 2005). According to this, the present work was based on a previous selection of three markers that were the most adequate to separate haplotypes in *B. bassiana* collected from different locations in Argentina (Toledo *et al.* 2019). The results indicated that markers 826, BA3 and KA5 were also effective to detect genetic variability between isolates of *B. bassiana* and *P. lilacinum* collected from tobacco soils. It is interesting to mention that this study constitutes the first report for these markers to discriminate between *P. lilacinum* strains.

According to the relationship between fungal genetic diversity and pathogenicity the results show that *B. bassiana* isolates LPSc1215 and LPSc1364, recorded as the most pathogenic against both insect pests tested, were grouped in a same cluster with a similarity level of approximately 85%. This cluster was located next to another cluster that included the other two isolates (LPSc1212 and LPSc1365) highly pathogenic against *H. gelotopoeon*. Respect to *P. lilacinum*, isolates LPSc1368 and LPSc1369, both the most pathogenic against *D. speciosa* adults, formed a single cluster with a similarity level of 98%, while isolate LPSc1371, only pathogenic against *H. gelotopoeon*, was not grouped with any other isolate, but it was remained along forming a last cluster within the ISSR dendrogram. The results obtained in this work are in accordance with those recorded by Almeida Carneiro *et al.* (2008) who, through the use of RAPD markers, observed that *B.*

bassiana and *B. brongniartii* (Sacc.) isolates were grouped in three different clusters according to their levels of pathogenicity against *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). These authors detected that three RAPD polymorphic bands among *Beauveria* isolates were associated with the levels of pathogenicity against *S. frugiperda* and suggested a potential use of these markers as a first screening strategy of *Beauveria* isolates for biological control purposes against this pest. Similar results were reported by Amer *et al.* (2008). These authors evaluated the pathogenicity of *B. bassiana*, *B. brongniartii*, *M. anisopliae*, *M. flavoviridae* (Gams & Roszypal) and *Paecilomyces farinosus* (= *Isaria farinosa*) (Holmsk.) Fr. against *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae) and observed that only the highly virulent isolates *M. anisopliae* and *M. flavoviridae* showed two unique RAPD markers that were not present in the other pathogens. Their results show that RAPD markers would have diagnostic potential in identifying highly virulent isolates. Similarly, the results obtained in our investigation not only provide valuable information regarding genetic diversity of native fungal isolates and their pathogenicity against to *H. gelotopoeon* and *D. speciosa*, but also suggest the potential use of ISSR markers to identify isolates with different virulence against these two important pests of tobacco crops. Although future studies will be needed to establish a link among amplified bands and genes related to the ability of entomopathogenic fungi to kill insects.

References

- Aljanabi SM and Martinez I 1997 Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucleic. Acids Res.* **25** 4692–4693
- Almeida Carneiro A, Gomes EA, Teixeira Guimarães C, Tavares Fernandes F, Portilho Carneiro N and Cruz I 2008 Molecular characterization and pathogenicity of isolates of *Beauveria* spp. to fall armyworm. *Pesqui. Agropecu. Bras.* **43** 513–520
- Amer M., El-Sayed TI, Bakheit HK, Moustafa SA, El-Sayed YA 2008 Pathogenicity and genetic variability of five entomopathogenic fungi against *Spodoptera littoralis*. *Res. J. Agric. Biol. Sci.* **4** 354–367
- Aquino De Muro M, Elliott S, Moore D, Parker BL, Skinner M, William REID and El Bouhssini M 2005 Molecular characterisation of *Beauveria bassiana* isolates obtained from overwintering sites of Sunn Pests (*Eurygaster* and *Aelia* species). *Mycol. Res.* **109** 294–306

- Ayala-Zermeño MA, Gallou A, Berlanga-Padilla AM, Serna-Domínguez MG, Arredondo-Bernal HC and Montesinos-Matías R 2015 Characterisation of entomopathogenic fungi used in the biological control programme of *Diaphorina citri* in Mexico. *Biocontrol Sci. Techn.* **25** 1192–1207
- Boetel MA, Majumdar A, Jaronski ST and Horsley RD 2012 Cover crop and conidia delivery system impacts on soil persistence of *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) in sugarbeet. *Biocontrol Sci. Techn.* **22** 1284–1304
- Fernández Bravo MDC 2017 Diversity, population dynamics and ecology of entomopathogenic fungi from soil and phylloplane of Mediterranean agroforestry systems and the effect of UV-B over their virulence, Doctoral Thesis, Cordoba University, Spain.
- Bruck DJ 2009 Impact of fungicides on *Metarhizium anisopliae* in the rhizosphere, bulk soil and in vitro. *BioControl* **54** 597–606
- Cruz LP, Gaitan AL and Gongora CE 2006 Exploiting the genetic diversity of *Beauveria bassiana* for improving the biological control of the coffee berry borer through the use of strain mixtures. *Appl. Microbiol. Biotechnol.* **71** 918–926
- Da Silva FR, Kvitko K, Rohr P, Abreu MB, Thiesen FV and Da Silva J 2014 Genotoxic assessment in tobacco farmers at different crop times. *Sci. Total Environ.* **490** 334–341
- Dalleau-Clouet C, Gauthier N, Risterucci AM, Bon MC and Fargues J 2005 Isolation and characterization of microsatellite loci from the entomopathogenic hyphomycete, *Paecilomyces fumosoroseus*. *Mol. Ecol. Notes.* **5** 496–498
- de Barros NM, Fronza E and Bertholdo-Vargas LR 2015 Use of fungi as biopesticides; In: *Biopesticides handbook* (eds) L. M Nollet and H. S.Rathore (London, New York: CRC Press, Taylor & Francis Group) pp 247–281
- de Faria MR and Wraight SP 2007 Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biol. Control.* **43** 237–256
- Domsch KH, Gams W and Anderson TH 2003 *Compendium of soil fungi* (New York: Academic Press) pp 894
- Enkerli J, Kölliker R, Keller S and Widmer F 2005 Isolation and characterization of microsatellite markers from the entomopathogenic fungus *Metarhizium anisopliae*. *Mol. Ecol. Notes.* **5** 384–386
- Enkerli J and Widmer F 2010 Molecular ecology of fungal entomopathogens: molecular genetic tools and their applications in population and fate studies. *BioControl.* **55** 17–37
- Estrada ME, Camacho MV and Benito C 2007 The molecular diversity of different isolates of *Beauveria bassiana* (Bals.) Vuill. as assessed using intermicrosatellites (ISSRs). *Cell. Mol. Biol. Lett.* **12** 240
- Fernandes EKK, Costa GL, Lage Moraes AM, Zahner V and Pinheiro Bittencourt VRE 2006 Study on morphology, pathogenicity, and genetic variability of *Beauveria bassiana* isolates obtained from *Boophilus microplus* tick. *Parasitol. Res.* **98** 324–332
- Flint SD, Miller CD, Anderson AJ and Roberts DW 2002 Damage and recovery from UV-B exposure in conidia of the entomopathogens *Verticillium lecanii* and *Aphanocladium album*. *Mycologia.* **94** 912
- Fontecha G, Trabanino R, Pérez-Borrero B, Catalán P, Aguilar E, Gallego FJ and Benito C 2011 Molecular characterization of Centro-american isolates of *Beauveria bassiana* for the control of the coffee drill. *Revista Ciencia y Tecnología.* **8** 39–61
- Hall TA 1999 BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98NT. *Nucleic Acids Symp. Ser.* **41** 95–98
- Hummel RL, Walgenbach JF, Barbercheck ME, Kennedy GG, Hoyt GD and Arellano C 2015 Effects of production practices on soilborne entomopathogens in western North Carolina vegetable systems. *Environ. Entomol.* **31** 84–91
- Lecuona RE 1996 *Pathogenic microorganisms used in the microbial control of pest insects* (Buenos Aires, Argentina: Talleres gráficos Mariano Mas) pp 338
- Mercado Cárdenas G; Gimenez Monge JL and Ortega AE 2013 *Manual of recognition of pests and biotic and abiotic diseases in Tobacco* (Salta: INTA ediciones) 188pp
- Mishra J, Tewari S, Singh S and Arora NK 2015 Biopesticides: where we stand? In: *Plant microbes symbiosis: applied facets* (eds) N. K., Arora (New Delhi: Springer) pp 37–75
- Nagamani A, Kunwar IK and Manoharachary C 2006 *Handbook of soil fungi* (New Delhi: I.K International Pvt. Ltd) pp 436
- Nussenbaum AL and Lecuona RE 2012 Selection of *Beauveria bassiana* sensu lato and *Metarhizium anisopliae* sensu lato isolates as microbial control agents against the boll weevil (*Anthonomus grandis*) in Argentina. *J. Invert. Pathol.* **110** 1–7
- Oulevey C, Widmer F, Kölliker R and Enkerli J 2009 An optimized microsatellite marker set for detection of *Metarhizium anisopliae* genotype diversity on field and regional scales. *Mycol. Res.* **113** 1016–1024
- Patana R 1977 Rearing selected western cotton insects in the laboratory. U.S. Department of Agriculture, Agricultural Research Service, ARS W-51, 8
- Pathan EK, Patil AV and Deshpande MV 2019 Bioprospecting of fungal entomo- and myco-pathogens. In: *Advancing frontiers in mycology and mycotechnology* (eds) T Satyanarayana, S Deshmukh and M Deshpande (Singapore: Springer) pp 497–513
- Pecchioni MTD 1989 Rearing *Diabrotica speciosa* (Coleoptera: Chrysomelidae) under laboratory conditions. *Revista Peruana de Entomología* **31** 86–90

- Pilz C, Wegensteiner R and Keller S 2007 Selection of entomopathogenic fungi for the control of the western corn rootworm *Diabrotica virgifera virgifera*. *J. App. Entomol.* **131** 426–431
- Posadas JB, Comerio RM, Mini JI, Nussenbaum AL and Lecuona RE 2012 A novel dodine-free selective medium based on the use of cetyl trimethyl ammonium bromide (CTAB) to isolate *Beauveria bassiana*, *Metarhizium anisopliae* sensu lato and *Paecilomyces lilacinus* from soil. *Mycologia* **104** 974–980
- Powell WA, Klingeman WE, Ownley BH, Gwinn KD, Dee M and Flanagan PC 2007 Endophytic *Beauveria bassiana* in tomatoes yields mycosis in tomato fruitworm larvae. *HortScience* **42** 933–933
- Rehner SA and Buckley E 2005 A *Beauveria* phylogeny inferred from nuclear ITS and EF1- α sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* **97** 84–98
- Samson RA, Evans HC and Latgé JP 1988 *Atlas of entomopathogenic fungi* (Berlin: Springer-Verlag) pp 187
- Sandhu SS, Shukla H, Aharwal RP, Kumar S and Shukla S 2017 Efficacy of entomopathogenic fungi as green pesticides: current and future prospects. In: *Microorganisms for Green Revolution* (eds) DG Panpatte, YK Jhala, RV Vyas and HN Shelat (Singapore: Springer) pp 327–349
- Seifert K, Morgan-Jones G, Gams W and Kendrick B 2011 *The genera of Hyphomycetes* (Utrecht: CBS-KNAW Fungal Biodiversity Centre) pp 997
- Shapiro-Ilan DI, Gardner WA, Wells L and Wood BW 2012 Cumulative impact of a clover cover crop on the persistence and efficacy of *Beauveria bassiana* in suppressing the pecan weevil (Coleoptera: Curculionidae). *Environ. Entomol.* **41** 298–307
- Shapiro-Ilan DI, Gardner WA, Wells L, Cottrell TE, Behle RW and Wood BW 2013 Effects of entomopathogenic fungus species, and impact of fertilizers, on biological control of Pecan weevil (Coleoptera: Curculionidae). *Environ. Entomol.* **42** 253–261
- Serna-Domínguez MG, Andrade-Michel GY, Rosas-Valdez R, Castro-Félix P, Arredondo-Bernal HC. and Gallou, A 2019 High genetic diversity of the entomopathogenic fungus *Beauveria bassiana* in Colima, Mexico. *J. Inverteb. Pathol.* **163** 67–74
- Sevim A, Höfte M and Demirbağ Z 2012 Genetic variability of *Beauveria bassiana* and *Metarhizium anisopliae* var. *anisopliae* isolates obtained from the Eastern Black Sea Region of Turkey. *Turk J. Biol.* **36** 255–265
- StatSoft Inc 2005 STATISTICA (data analysis software system), version 7.1. (www.statsoft.com)
- Tamura K, Peterson D, Peterson N, Nei M and Kumar S 2011 MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28** 2731–2739
- Takatsuka J 2007 Characterization of *Beauveria bassiana* isolates from Japan using inter-simple-sequence-repeat-anchored polymerase chain reaction (ISSR-PCR) amplification. *Appl. Entomol. Zool.* **42** 563–571
- Taylor JW, Jacobson DJ and Fisher MC 1999 The evolution of asexual fungi: reproduction, speciation and classification. *Annu. Rev. Phytopathol.* **37** 197–246
- Thompson SR, Brandenburg RL and Arends JJ 2006 Impact of moisture and UV degradation on *Beauveria bassiana* (Balsamo) Vuillemin conidial viability in turfgrass. *Biol. Control.* **39** 401–407
- Tigano-Milani MS, Honeycutt RJ, Lacey LA, Assis R, McClellan M and Sobral BWS 1995 Genetic variability of *Paecilomyces fumosoroseus* isolates revealed by molecular markers. *J. Inverteb. Pathol.* **65** 274–282
- Toledo AV, Franco MEE, Medina R, Marino de Remes Lenicov AM and Balatti PA 2019 Assessment of the genetic diversity of Argentinean isolates of *Beauveria bassiana* (Ascomycota: Hypocreales) using ISSR markers. *J. King Saud Uni. Sci.* **31** 1264–1270
- Trissi AN, El Bouhsini M, Alsalti MN, von Korff M, Hamwieh A, Skinner M, Parker BL and Baum M 2013 Genetic diversity among summer and winter *Beauveria bassiana* populations as revealed by AFLP analysis. *J. Asia-Pac. Entomol.* **16** 269–273
- Vega FE 2008 Insect pathology and fungal endophytes. *J. Invertebr. Pathol.* **98** 277–279
- Wang S, Miaox X, Zhao W, Huang B, Fan M, Li Z and Huang Y 2005 Genetic diversity and population structure among strains of the entomopathogenic fungus, *Beauveria bassiana*, as revealed by inter-simple sequence repeats (ISSR). *Mycol. Res.* **109** 1364–1372
- White TJ, Bruns T, Lee S and Taylor J 1990 Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols. a guide to methods and applications* (eds) MA Innis, DH Gelfand, JJ Sninsky and TJ White (San Diego: Academic Press) pp 315–322
- Yang H, Qin C, Chen Y, Zhang G, Dong L and Wan S 2019 Persistence of *Metarhizium* (Hypocreales: Clavicipitaceae) and *Beauveria bassiana* (Hypocreales: Clavicipitaceae) in tobacco soils and potential as biocontrol agents of *Spodoptera litura* (Lepidoptera: Noctuidae). *Environ. Entomol.* **48** 147–155
- Zimmermann G 2007 Review on safety of the entomopathogenic fungi. *Beauveria bassiana* and *Beauveria brongniartii*. *Biocontrol Sci. Tech.* **17** 553–596