

Supplementary File:

***RHBDD2*: a 5-Fluorouracil responsive gene overexpressed in the advanced stages
of colorectal cancer.**

Lacunza E^{1*}, Canzoneri R¹, , Rabassa ME¹, Zwenger A^{1,2}, Segal-Eiras A¹, Croce MV¹,
Abba MC¹

¹ CINIBA, Facultad de Ciencias Médicas, Universidad Nacional de La Plata. La Plata, Argentina.

² Hospital Provincial Neuquén “Dr Castro Rendón”. Servicio de Oncología. Neuquén, Argentina.

*Corresponding Author: Dr. Ezequiel Lacunza. CINIBA, Facultad de Ciencias Médicas, Universidad Nacional de La Plata. E-mail: ezequiellacunza@hotmail.com




Fig 1: *RHBDD2* mRNA expression in colon cancer cell lines. a) *RHBDD2* mRNA expression in 30 colon cancer cell lines; data obtained from Mariadason et al. 2003. Code bars of 5FU sensitivity and Tp53 status are shown below. References of 5FU bar: Red squares, high; pink squares, moderate; light-pink squares, low; white squares, resistant. References of Tp53 bar: gray squares, mutant; white squares, wilde type b) *RHBDD2* expression in colon cancer cell line HCT116 exposed to 5 FU, according to Boyer et al (2006).