

The geophilomorph centipedes (Chilopoda) of Brazilian Amazonia

DONATELLA FODDAI*
LUIS ALBERTO PEREIRA**
ALESSANDRO MINELLI*

Resumen. Se presenta un catálogo de los Chilopoda Geophilomorpha conocidos para el territorio brasileño correspondiente a la Región Amazónica. Se conocen treinta y una especies para esta región; excepto seis, todas las demás son endémicas como lo es un género (*Hyphydrophilus*) entre los once conocidos de Amazonia hasta el presente.

Palabras clave: Chilopoda-Geophilomorpha, Amazonia brasileña, composición faunística.

Abstract. A catalogue of the Chilopoda Geophilomorpha known from Brazilian Amazonia is here presented. Thirty-one species are known from this region; all but six of these species are endemic to it, as is one (*Hyphydrophilus*) of the eleven genera hitherto known from Amazonia.

Key words: Chilopoda - Geophilomorpha, Brazilian Amazonia, faunal composition.

Introduction

Geophilomorph centipedes are currently classified into 14 families, of which 11 occur in the Neotropical Region, the remaining families, not further mentioned in

*Dipartimento di Biologia, Università degli Studi di Padova, Via Ugo Bassi 58B, I 35131 Padova, Italia. foddai@civ.bio.unipd.it, almin@civ.bio.unipd.it

**Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s.n., (1900) La Plata, R. Argentina. lpereira@museo.fcnym.unlp.edu.ar

this paper, being the Eucratonychidae, Gonibregmatidae and Dignathodontidae. Some 1100 species in 180 genera are known worldwide and ca. 310 have been hitherto described from the Neotropical Region including the whole of México (Foddai & Minelli 2000; Foddai, Minelli & Pereira 2002; Foddai, Pereira & Minelli 2000, 2002).

Morphology and biology

The main traits of geophilomorph morphology and biology, with particular reference to the Amazonian fauna, have been recently summarized by Foddai, Minelli & Pereira (2002); therefore we will limit this account to very few details.

The number of leg-bearing segments in the Neotropical representatives of this group range between 27, hitherto recorded from a single specimen of *Schendylops oligopus* (Pereira, Minelli & Barbieri 1995) from Amazonia (Minelli *et al.* 2000) and 151 in *Notiphilides grandis* Brölemann 1904, also from Amazonia; the size ranges between the 5 mm of *Dinogeophilus oligopodus* Pereira 1984 from Argentina and the 14 cm of *Notiphilides grandis*.

In the Neotropics, geophilomorphs occur both in forest and in open habitats, especially in the litter and soil, with peculiar adaptations to periodically inundated forest. All species are blind and primarily adapted for life in dark environments and preying on small invertebrates.

The females lay a group of eggs, less than ten to about eighty, in a cavity in the soil, and remain coiled around them until the hatchlings are able to feed for themselves.

The hatching juveniles are already provided with the full number of legs.

Amazonian geophilomorphs

As for Amazonia, some biotopes in the vicinity of Manaus have been intensively investigated during the last two decades. Some species are well-adapted to survive in periodically inundated forests (Adis *et al.* 1996; Morais *et al.* 1997) and morpho-anatomical modifications for surviving submersion have been recorded in the respiratory system of a couple of species (Adis *et al.* in prep.).

Most Amazonian species belong to the large families Geophilidae and Schendylidae. The genera with the highest number of Amazonian species are *Ribautia* and *Schendylops*, each with six Amazonian species.

At the species level, Amazonian geophilomorphs show a high level of endemism, but there is only one endemic genus, the geophilid *Hyphydrophilus* Pereira, Minelli & Barbieri 1994.

Pereira, Foddai & Minelli (1997) attempted to reconstruct the recent history of this centipede group in Amazonia on basis of the hypothesis of Pleistocene refugia (*sensu* Haffer 1982). As far as we know, half of the Neotropical species belonging to

the genera *Schendylops*, *Pectiniunguis* and *Ribautia* are endemic to areas that have been identified as Pleistocene refugia, based on evidence from woody Angiosperms (Nelson *et al.* 1990, Prance 1982). The main routes of subsequent distribution did probably follow the valleys of the major rivers (Amazon, Orinoco, Paraná, Paraguay).

Unfortunately, the current level of knowledge of the Amazonian geophilomorphs, and those of the Neotropical Region at large, is still very limited and future research may modify the current picture to a significant extent.

A list of the Neotropical genera of Geophilomorpha, with the number of species hitherto known from Brazilian Amazonia is given in Table 1.

Table 1. Neotropical genera of Chilopoda - Geophilomorpha: total number of species, number of species in the Neotropical Region and in Brazilian Amazonia

<i>Family</i>	<i>Genus</i>	<i>Total number of species</i>	<i>Number of species in the Neotropical Region</i>	<i>Number of species in the Brazilian Amazonia</i>
Aphilodontidae	<i>Aphilodon</i>	14	4	-
	<i>Mecistauchenus</i>	1	1	-
	<i>Mecophilus</i>	1	1	-
Ballophilidae	<i>Ballophilus</i>	41	3	-
	<i>Caritohallex</i>	1	1	-
	<i>Cerethmus</i>	1	1	-
	<i>Clavophilus</i>	1	1	-
	<i>Diplethmus</i>	6	6	-
	<i>Ityphilus</i>	22	17	6
	<i>Koinethmus</i>	1	1	-
	<i>Leucolinum</i>	1	1	-
	<i>Taeniolinum</i>	6	6	1
	<i>Tanophilus</i>	1	1	-
	<i>Zygethmus</i>	1	1	-
Eriphantidae	<i>Eriphantes</i>	1	1	-
Geophilidae	<i>Apogeophilus</i>	2	2	-
	<i>Aztekophilus</i> (<i>Aztekophilus</i>)	1	1	-
	<i>Aztekophilus</i> (<i>Thylakiophilus</i>)	1	1	-
	<i>Barrophilus</i>	1	1	-
	<i>Chilenophilus</i>	4	4	-
	<i>Chomatophilus</i>	3	3	-

Table 1. *Continues*

<i>Family</i>	<i>Genus</i>	<i>Total number of species</i>	<i>Number of species in the Neotropical Region</i>	<i>Number of species in the Brazilian Amazonia</i>
	<i>Dinogeophilus</i>	2	2	-
	<i>Ecuadoron</i>	1	1	-
	<i>Erithophilus</i>	1	1	-
	<i>Eurytion</i>	29	8	-
	<i>Filipponus</i>	1	1	-
	<i>Garrina</i>	13	11	-
	<i>Geoperingueyia</i>	11	1	-
	<i>Gospina</i>	2	1	-
	<i>Hyphydrophilus</i>	2	2	2
	<i>Ketampa</i>	1	1	-
	<i>Nabocodes</i>	1	1	-
	<i>Nesidiphilus</i>	7	7	-
	<i>Nicopus</i>	1	1	-
	<i>Oligna</i>	1	1	-
	<i>Orinomerium</i>	1	1	-
	<i>Pachymerinus</i>	9	6	-
	<i>Pachymerium</i>	38	6	-
	<i>Pandineum</i>	10	10	-
	<i>Peruphilus</i>	1	1	-
	<i>Piestophilus</i>	2	2	-
	<i>Polycricus</i>	16	16	-
	<i>Portoricona</i>	2	2	-
	<i>Ribautia</i>	Ca. 45	23	6
	<i>Schendyloides</i>	2	2	-
	<i>Schizonampa</i>	3	1	1
	<i>Schizonium</i>	6	6	-
	<i>Sogona</i>	5	3	-
	<i>Suturodes</i>	4	4	-
	<i>Synerium</i>	1	1	-
	<i>Taiyuna</i>	5	1	-
	<i>Telocricus</i>	5	5	-
	<i>Tuoba</i>	10	3	-
Himantariidae	<i>Arcophilus</i>	2	2	-
	<i>Californiphilus</i>	2?	1	-
	<i>Causerium</i>	1	1	-
	<i>Chomatobius</i>	9	4	-
	<i>Geoballus</i>	2	2	-
	<i>Straberax</i>	1	1	-
Linotaeniidae	<i>Araucania</i>	1	1	-

Table 1. *Continues*

<i>Family</i>	<i>Genus</i>	<i>Total number of species</i>	<i>Number of species in the Neotropical Region</i>	<i>Number of species in the Brazilian Amazonia</i>
Macronicophilidae	<i>Pagotaenia</i>	1	1	-
	<i>Tomotaenia</i>	10	1	-
	<i>Macronicophilus</i>	4	4	2
Mecistocephalidae	<i>Mecistocephalus</i>	Ca. 130	5	1
	<i>Tygarrup</i>	15	1	-
Neogeophilidae	<i>Evallogeophilus</i>	1	1	-
	<i>Neogeophilus</i>	3	3	-
Oryidae	<i>Henioria</i>	1	1	-
	<i>Metaxythus</i>	1	1	-
	<i>Notiphilides</i>	3	3	1
	<i>Orphnaeus</i>	20	6	2
	<i>Pentorya</i>	4	1	-
	<i>Titanophilus</i>	4	4	-
	<i>Trematorya</i>	1	1	-
Schendylidae	<i>Algunguis</i>	1	1	-
	<i>Bimindyla</i>	1	1	-
	<i>Ctenophilus</i>	12	1	-
	<i>Cymochilus</i>	1	1	-
	<i>Marsikomerus</i>	5	1	-
	<i>Mexiconyx</i>	1	1	-
	<i>Morunguis</i>	1	1	-
	<i>Nannopodellus</i>	1	1	-
	<i>Nyctunguis</i>	18	3	-
	<i>Orygmadyla</i>	1	1	-
	<i>Parunguis</i>	4	3	-
	<i>Pectiniunguis</i>	23	20	3
	<i>Portoricellus</i>	1	1	-
	<i>Schendylellus</i>	1	1	-
	<i>Schendylops</i>	63	51	6
	<i>Sogodes</i>	1	1	-
	<i>Sogolabis</i>	1	1	-
	<i>Thindyla</i>	1	1	-

Catalogue of the Chilopoda Geophilomorpha from Brazilian Amazonia

Family BALLOPHILIDAE

Genus *Ityphilus* Cook

Ityphilus Cook 1899: 306. Type-species, *lilacinus* Cook (orig. des.).

Thalthybius Attems 1901: 139-140. Type-species, *melanostigma* Attems (mon.).

Thalthybius, subg. *Prionothalthybius* Brölemann 1909a: 334. Type-species, *perrieri* Brölemann 1909 (mon.).

calinus Chamberlin 1957: 25, fig. 7. Type-locality: "Columbia: 13 mi. W. of Cali, Valle". Distr. - Colombia (Cali), Brazil (AM [Reserva Ducke]).

crabilli Pereira, Minelli & Barbieri 1994: 163, 164-166, figs. 1-28. Type-locality: Brazil, AM, Rio Tarumã Mirim. Distr. - Brazil (AM [Rio Tarumã Mirim, Reserva Ducke]).

demoraisi Pereira, Minelli & Barbieri 1995: 325, 327-328, figs. 3-42. Type-locality: Brazil, AM, Manaus [Reserva Ducke].

guianensis Chamberlin 1921: 23-25, pl. v, figs. 20, 23-24. Type-locality: "British Guiana: Dunoon". Distr. - Trinidad, Guyana, Brazil (AM).

saucius Pereira, Foddai & Minelli 2000: 8-9, figs. 69-89. Type-locality: Brazil, AM, 02° 34' S, 60° 06' W.

sensibilis Pereira, Foddai & Minelli 2000: 9-11, figs. 90-112. Type-locality: Brazil, AM, 02° 34' S, 60° 06' W.

Genus *Taeniolinum* Pocock

Taeniolinum Pocock 1893: 469, 471. Type-species, *setosum* Pocock (mon.).

arborum Pereira, Minelli & Barbieri 1994: 163, 167-169, figs. 30-59. Type-locality: Brazil, AM, Rio Tarumã Mirim.

Family GEOPHILIDAE

Genus *Hyphydrophilus* Pereira, Minelli & Barbieri

Hyphydrophilus Pereira, Minelli & Barbieri 1994: 169. Type-species, *adisi* Pereira, Minelli & Barbieri (orig. des.).

adisi Pereira, Minelli & Barbieri 1994: 163, 169-171, figs. 60-91. Type-locality: Brazil, AM, Rio Tarumã Mirim.

projectus Pereira, Foddai & Minelli 2000: 11-13, figs. 113-140. Type-locality: Brazil, AM, 2° 30' S, 60° 10' W.

Genus ***Ribautia*** Brölemann

Ribautia Brölemann 1909a: 335-336. Type-species, *bouvieri* Brölemann (mon.).

bouvieri Brölemann 1909c: 421-425, figs. 19-26. Type-locality: “Brésil: Haut Carsevène”.

centralis (Silvestri) 1907: 256-257, figs. 81-86 (*Eurytion*). Type-locality: “Columbia: Inirida?”. Distr. - Colombia, Brazil (AM [Rio Tarumã Mirím]).

difficilis Pereira, Minelli & Barbieri 1995: 325, 333-334, figs. 123-152. Type-locality: Brazil, AM, Lago Janauarí.

ducalis Pereira, Minelli & Barbieri 1995: 325, 329, 331, figs. 46-87. Type-locality: Brazil, AM, Manaus [Reserva Ducke].

onycophaena Pereira, Foddai & Minelli 2000: 13-15, figs. 141-167. Type-locality: Brazil, AM, 02° 34' S, 60° 06' W.

proxima Pereira, Minelli & Barbieri 1995: 325, 331-333, figs. 88-122. Type-locality: Brazil, AM, Manaus [Reserva Ducke].

Genus ***Schizonampa*** Chamberlin

Schizonampa Chamberlin 1914: 214-215. Type-species, *manni* Chamberlin (orig. des.).

manni Chamberlin 1914: 215-217, pl. 6, figs. 1-7. Type-locality: “Brazil, State of Pará: Pará” [i. e., PA, Belém].

Family MACRONICOPHILIDAE

Genus ***Macronicophilus*** Silvestri

Macronicophilus Silvestri 1909: 267. Type-species, *ortonedae* Silvestri (orig. des.).

abbreviatus Pereira, Foddai & Minelli 2000: 16-18, figs. 168-197. Type-locality: Brazil, AM, Manaus, 03° 08' S, 60° 01' W.

unguiseta Pereira, Foddai & Minelli 2000: 18-19, figs. 205-237. Type-locality: Brazil, AM, Rio Tarumã Mirim.

Family MECISTOCEPHALIDAE

Genus ***Mecistocephalus*** Newport

Mecistocephalus Newport 1843: 178. Type-species, *punctifrons* Newport (subs. des., Chamberlin 1914).

guildingii Newport 1845: 429, 430, pl. xxiii, figs. 18-19. Type-locality: “In insula

Caribaea S.ti Vincentii". Distr. – United States (Florida), Panamá (Canal Zone), Cuba, Haiti, Guadeloupe, Jamaica, Martinique, St. Vincent, Brazil (AM [Manaus], RJ). *janeirensis* Verhoeff 1937: 230. Type-locality: Brazil, Rio de Janeiro.

Family ORYIDAE
Genus **Notiphilides** Latzel

Notiphilides Latzel 1880: 20. Type-species, *Notiphilus maximiliani* Humbert & Saussure (orig. des.).

grandis Brölemann [1905]: 71-73, pl. 1, figs. 8-11. Type-locality: Brazil, AM, Manaus.

Genus **Orphnaeus** Meinert

Orphnaeus Meinert 1870: 17-19. Type-species, *lividus* Meinert (subs. des., Crabill, 1968) = *brevilabiatius* (Newport).

Azygethus Chamberlin 1920: 32. Type-species, *atopus* Chamberlin (orig. des.) = *brevilabiatius* (Newport).

Andenophilus Verhoeff 1941: 62-64. Type-species, *striatus* Verhoeff (mon.) = *brasilianus* (Humbert & Saussure).

brasilianus (Humbert & Saussure) 1870: 205 (*Chomatobius*). Type-locality: "Brasilia". Distr. - Honduras, Nicaragua, Panama, Ecuador, Peru, Brazil (PA, RN, PE, RJ); South Africa, Madagascar.

brasilianus var. *nigropictus* Attems 1903: 201. Type-locality not fixed; described from Venezuela and Madagascar.

branneri Chamberlin 1914: 152. Type-locality: Brazil, RN, Natal.

striatus Verhoeff 1941: 64-65, figs. 91-96. Type-locality: "Südperu bei Sivia".

brevilabiatius (Newport), 1845: 436 (*Geophilus*). Type-locality: "In Orâ Tenasserim Peninsulae Indiae Ulteriori". Distr. - Guatemala, Nicaragua, Panama, Cuba, Bahama Is., Jamaica, Venezuela, Guyana, Brazil (AM [Manaus], PA, PE, RJ), Galapagos Is.; Borneo, Java, Sumatra, Celebes, Australia, Hawaii, Solomon Is., Samoa Is., Society Is., Japan, Formosa, Cameroon, Madagascar, Yemen, United States (Florida), New Caledonia.

bilineatus Peters 1855: 83. Type-locality: "Inhanbane, 23 1/5 lat. Austr.".

brasiliensis Meinert 1870: 20-21, pl. iii, fig. 12. Type-locality: Brazil, RJ, Rio de Janeiro.

lividus Meinert 1870: 19-20, pl. ii, figs. 6-11. Type-locality not fixed; described from Hawaii (Oahu) and Nicobares.

xanti Tömösvary 1885: 64. Type-locality: "Borneo".

phosphoreus, *auct.*

Family SCHENDYLIDAE
Genus *Pectiniunguis* Bollman

Pectiniunguis Bollman 1889: 212. Type-species, *americanus* Bollman (orig. des.).
Adenoschendyla Brölemann & Ribaut 1911a: 192. Type-species, *geayi* Brölemann & Ribaut (mon.).
Litoschedyla Chamberlin 1923: 391. Type-species, *Pectinunguis insulanus* Brölemann & Ribaut (orig. des.).

ascendens Pereira, Minelli & Barbieri 1994: 174-176, figs. 120-149. Type-locality: Brazil, AM, Rio Tarumã Mirim.

ducalis Pereira, Minelli & Barbieri 1995: 325, 338-340, figs. 188-225. Type-locality: Brazil, AM, Manaus [Reserva Ducke].

geayi (Brölemann & Ribaut) 1911b: 219-220 (*Adenoschendyla*). Type-locality: “Bas Carsévène”. Distr. - Brazil (AM [Rio Tarumã Mirim; Lago Janauari]), State of Pará, Bas Carsévène.

Genus *Schendylops* Cook

Schendylops Cook 1899: 305. Type-species, *Schendyla grandidieri* Saussure & Zehnter (orig. des.).

Schendylurus Silvestri 1907: 245-246. Type-species, *australis* Silvestri (mon.).

Schendylurus, subgen. *Plutoschendylurus* Brölemann & Ribaut 1912: 115. Type-species, *tropicus* Brölemann & Ribaut (subs. des., Hoffman & Pereira 1997).

Nesondyla Chamberlin 1950: 135. Type-species, *nealota* Chamberlin (orig. des.).

Koepckeilla Kraus 1954: 311-312. Type-species, *titicacaensis* Kraus (orig. des.).

amazonicus (Pereira, Minelli & Barbieri) 1994: 163, 176-178, figs. 150-177 (*Schendylurus*). Type-locality: Brazil, AM, Rio Tarumã Mirim.

bakeri (Chamberlin) 1914: 152, 196-198, pl. 4, figs. 5-8 (*Schendylurus*). Type-locality: Brazil, AM, Manaus.

continuus (Pereira, Minelli & Barbieri) 1995: 325, 340-342, figs. 229-265 (*Schendylurus*). Type-locality: Brazil, AM, Manaus [Reserva Ducke].

janauarius (Pereira, Minelli & Barbieri) 1995: 325, 342-343, figs. 266-293 (*Schendylurus*). Type-locality: Brazil, AM, Lago Janauari.

marchantariae (Pereira, Minelli & Barbieri) 1995: 325, 344-345, figs. 300-336 (*Schendylurus*). Type-locality: Brazil, AM, Rio Solimões, Ilha da Marchantaria, 03° 15' S, 59° 58' W. Distr. - Peru (Loreto [Iquitos]), Brazil (AM [Ilha da Marchantaria, Ilha de Curari]).

oligopus (Pereira, Minelli & Barbieri) 1995: 325, 345-347, figs. 337-371 (*Schendylurus*). Type-locality: Brazil, AM, Manaus [Reserva Ducke].

Acknowledgments. We are indebted to Nelson Papavero (Museu Paranense Emilio Goeldi, Belém, PA, Brasil) for his encouragement and help during the preparation of this paper.

Literature cited

- ADIS, J., A. MINELLI, J.W. DE MORAIS, L.A. PEREIRA, F. BARBIERI, & J.M.G. RODRIGUES. 1996. On abundance and phenology of Geophilomorpha (Chilopoda) from central Amazonian upland forests. *Ecotropica* 2: 165-175.
- ATTEMS, C. 1901. Dr. Brauer's Myriopoden-Ausbeute auf den Seychellen im Jahre 1895. *Zoologischer Jahrbücher, Systematik* 13: 133-171.
- ATTEMS, C. 1903. Synopsis der Geophiliden. *Zoologischer Jahrbücher, Systematik* 18: 155-302.
- BOLLMAN, C. H. 1889. Myriapoda in scientific results of explorations by the U.S. Fish. Commission Steamer 'Albatros'. In: C.H. Bollman. The Myriapoda of North America. *Proceedings of the United States National Museum, Washington* 12: 211-216.
- BRÖLEMANN, H. W. 1905. Myriapodes du Museu Paulista. II^e mémoire. *Revista do Museu Paulista* 6: 63-96.
- BRÖLEMANN, H. W. 1909a. A propos d'un système des Géophilomorphes. *Archives de Zoologie expérimentale et générale* 3 (5): 303-340.
- BRÖLEMANN, H. W. 1909b. Essai d'un catalogue des Myriapodes du Brésil. *Catálogos da Fauna Brasileira*. 2: xxx + 94 pp. Museu Paulista, São Paulo.
- BRÖLEMANN, H. W. 1909c. Quelques géophilides des collections du Muséum d'Histoire Naturelle. *Bulletin du Muséum National d'Histoire Naturelle* 7: 415-431.
- BRÖLEMANN, H. W. & H. RIBAUT. 1911a. Notes préliminaires sur les genres de Schendylina. *Bulletin de la Société entomologique de France* [1911](8): 191-193.
- BRÖLEMANN, H. W. & H. RIBAUT. 1911b. Diagnoses préliminaires d'espèces nouvelles de Schendylina [Myriap. Geophilomorpha]. *Bulletin de la Société Entomologique de France* [1911](10): 219-222.
- BRÖLEMANN, H. W. & H. RIBAUT. 1912. Essai d'une monographie des Schendylinae (Myriapodes, Géophilomorphes). *Nouvelles Archives du Muséum National d'Histoire Naturelle* 4(5): 53-183.
- BÜCHERL, W. 1940. Os chilopodos do Brasil. *Memorias do Instituto Butantan, São Paulo* 13: 43-362.
- BÜCHERL, W. 1941-1942. Catálogo dos quilópodos da zona neotrópica. *Memorias do Instituto Butantan, São Paulo* 15: 251-372.
- CHAMBERLIN, R. V. 1914. The Stanford Expedition to Brazil, 1911, John C. Branner, Director. The Chilopoda of Brazil. *Bulletin of the Museum of Comparative Zoology at Harvard College* 58(3): 151-221.
- CHAMBERLIN, R. V. 1920. The myriopod fauna of the Bermuda Islands, with notes on variation in *Scutigera*. *Annals of the Entomological Society of America* 13: 271-285.
- CHAMBERLIN, R. V. 1921. Results of the Bryant Walker Expeditions of the University of Michigan to Colombia, 1913 and British Guiana, 1914. *Occasional Papers of the Museum of Zoology, University of Michigan* 97: 1-28.
- CHAMBERLIN, R. V. 1923. On chilopods and diplopods from islands in the Gulf of California. *Proceedings of the California Academy of Natural Sciences, San Francisco* 4 (12): 389-407.

- CHAMBERLIN, R. V. 1950. Neotropical chilopods and diplopods in the Collection of the Department of Tropical Research, New York Zoological Society. *Zoologica, New York* 2(35): 133-144.
- CHAMBERLIN, R. V. 1957. Geophiloid chilopods taken in the Northern Andes in 1954-1955. *Proceedings of the Biological Society of Washington* 70: 21-30.
- COOK, O. F., 1899. The Geophiloidea of the Florida keys. *Proceedings of the Entomological Society of Washington* 4(3) (1896-1909): 302-312.
- CRABILL, R. E. Jr. 1968. Concerning the true identities of *Gosiphilus* and *Chomatobius* with redescription of the latter's type species (Chilopoda: Geophilomorpha: Himantariidae). *Entomological News* 79: 108-112.
- FODDAI, D. & A. MINELLI. 2000. Phylogeny of geophilomorph centipedes: old wisdom and new insights from morphology. *Fragmenta Faunistica, Warsaw* 43 (supplement), 61-71.
- FODDAI, D., A. MINELLI & L.A. PEREIRA. 2002. Geophilomorpha. In: J. Adis (ed.). *Amazonian Arachnida and Myriapoda*. Sofia-Moscow, Pensoft, pp. 459-474.
- FODDAI, D., L. A. PEREIRA & A. MINELLI. 2000. A catalogue of the geophilomorph centipedes (Chilopoda) from Central and South America including Mexico. *Amazoniana* 16(1/2): 59-185.
- FODDAI, D., L.A. PEREIRA & A. MINELLI. 2002. Geophilomorpha (Chilopoda). In: J. Llorente-Bousquets & J.J. Morrone (eds.). *Biodiversidad, taxonomía y biogeografía de artrópodos de México: Hacia una síntesis de su conocimiento*, vol. III. UNAM, México, D.F., pp. 417-427.
- HAFFER, J. 1982. General aspects of the refuge theory. In: G.T. Prance (ed.). *Biological diversification in the tropics*. New York. xvi+714 p. Columbia University Press, pp. 6-24.
- HUMBERT, A. & E. DE SAUSSURE. 1870. Myriapoda nova Americana. *Revue et Magasin de Zoologie pure et appliquée. Paris* 22 (2): 202-205.
- KRAUS, O. 1954. Myriapoden aus Peru. I. *Senckenbergiana* 34: 311-323.
- LATZEL, R. 1880. *Die Myriapoden der Österreichisch-Ungarischen Monarchie* 1: xv + 228 pp. J.C. Fischer, Wien.
- MEINERT, F. 1870. Myriapoda Musaei Hauniensis. - Bidrag til myriapodernes morfologi og systematik. I. Geophili. *Naturhistorisk Tidsskrift. Kjobenhavn* 7(3): 1-128.
- MINELLI, A., D. FODDAI, L.A. PEREIRA & J.G.E. LEWIS. 2000. The evolution of segmentation of centipede trunk and appendages. *Journal of Zoological Systematics and Evolutionary Research* 38: 103-117.
- MORAIS, J.W., J. ADIS, E. BERTI-FIHLLO, L.A. PEREIRA, A. MINELLI & F. BARBIERI. 1997. On abundance, phenology and natural history of Geophilomorpha from a mixedwater inundation forest in Central Amazonia (Chilopoda). *Entomologica Scandinavica Supplement* 51: 115-119.
- NELSON, B.W., C.A.C. FERREIRA, M.F. DA SILVA & M.L. KAWASAKI. 1990. Endemism centers, refugia and botanical collection density in Brazilian Amazonia. *Nature* 6277(345): 714-716.
- NEWPORT, G. 1843. On some new genera of the class Myriapoda. *Proceedings of the Zoological Society of London* 10: 177-181.
- NEWPORT, G. 1844-45. Monograph of the class Myriapoda order Chilopoda; with observations on the general arrangement of the Articulata. *Transactions of the Linnean Society of London* 9(19): 265-302; 349-439.
- PEREIRA, L. A., A. MINELLI & F. BARBIERI. 1994. New and little known geophilomorph centipedes from Amazonian inundation forests near Manaus, Brazil (Chilopoda, Geophilomorpha). *Amazoniana* 13(3-4): 163-204.

- PEREIRA, L. A., A. MINELLI & F. BARBIERI. 1995. Description of nine new centipede species from Amazonia and related matters on Neotropical geophilomorphs (Chilopoda: Geophilomorpha). *Amazoniana* 13(3-4): 325-416.
- PEREIRA, L. A., D. FODDAI & A. MINELLI, 1997. Zoogeographical aspects of Neotropical Geophilomorpha. *Entomologica Scandinavica Supplement* 51: 77-86.
- PEREIRA, L. A., D. FODDAI & A. MINELLI. 2000. New taxa of Neotropical Geophilomorpha (Chilopoda). *Amazoniana* 1/2(16): 1-57, 275 figs.
- PETERS, W.C.H. 1855. Über die Myriapoden im Allgemeinen und insbesondere über die in Mossambique beobachteten Arten dieser Familie. *Monatsber. königl. Akademische Wissenschaft Berlin* 75-85.
- POCOCK R.I. 1893. Contribution to our knowledge of the arthropod fauna of the West Indies. Part II. Chilopoda. *The Journal of the Linnean Society, Zoology, London* 24: 454-473.
- PRANCE, G.T. 1982. Forest refuges: evidence from woody Angiosperms. pp. 137-158. In: G.T. Prance (ed.). *Biological diversification in the tropics*. New York, xvi+714 p. Columbia University Press, pp.137-158.
- SILVESTRI, F. 1907. Neue und wenig bekannte Myriapoden des naturhistorischen Museums in Hamburg. *Mitteilungen aus des naturhistorischen Museum in Hamburg* 1906(24): 229-257.
- SILVESTRI, F., 1909. Descrizione preliminare di vari Artropodi, specialmente d' America. *Rendiconti della Accademia dei Lincei, Classe di Scienze Fisiche, Matematiche e Naturali* 18(5/6): 267-271.
- TÖMÖSVÁRY E. 1885. Myriopoda a Joahne Xanthus in Asia orientali collecta. Enumeravit specisque novas descripsit. *Természetráji Füzetek* 9: 63-72.
- VERHOEFF, K.W. 1937. Chilopoden aus Malacca nach den Objecten des Raffles Museum in Singapore. *Bulletin of the Raffles Museum, Singapore* 13: 198-270.
- VERHOEFF, K.W. 1941. Chilopoden und Diplopoden. In: E. Titschack (ed.). *Beiträge zur Fauna Perus* 1(2): 5-80.

Recibido: 8.VI.2004

Aceptado: 5.VII.2004