

TABLE S1. Growth rates and percentage of reduction in the growing rate of *Aspergillus* sp. in co-culture with brewer's yeasts

Treatment	Growth rate [cm/h] ^a	% Growth rate reduction ^b
<i>Af</i> + <i>Pk</i> MBELGA61	0.0146 ± 0.0007 ^{AB*}	57.2 ± 2.0
<i>Af</i> + <i>Sc</i> MBELGA62	0.0174 ± 0.0010 ^{B*}	49.0 ± 2.9
<i>Af</i> + starter M6	0.0133 ± 0.0018 ^{A*}	60.9 ± 5.4
<i>Af</i> control	0.0341 ± 0.0011	
<i>Ap</i> + <i>Pk</i> MBELGA61	0.0175 ± 0.0008 ^{B*}	54.9 ± 2.4
<i>Ap</i> + <i>Sc</i> MBELGA62	0.0178 ± 0.0002 ^{B*}	54.1 ± 0.5
<i>Ap</i> + starter M6	0.0135 ± 0.0008 ^{A*}	63.8 ± 4.3
<i>Ap</i> control	0.0387 ± 0.0003	

^aFungal growth rates were calculated as the slope of the linear section of the growth curves, located between 96 and 192 or 216 h of incubation (Armando et al., 2013). Results are expressed as the average of three replicas of each treatment ± standard deviation. ^bThe percentage was calculated regarding the positive control growth rate (100%). *Indicates significant differences with the respective positive control (P < 0.05). Different uppercase letters indicate significant differences between the growth rates of the treatments for each fungal species (P < 0.05). *Af*: *A. flavus* CMUNLP15. *Ap*: *A. parasiticus* CMUNLP7. *Pk* MBELGA61: *P. kudriavzevii* MBELGA61. *Sc* MBELGA62: *S. cerevisiae* MBELGA62.