



Referencias

REFERENCIAS

- Akinsanmi O.A., Backhouse D., Simpfendorfer S., Chakraborty S. (2006). Genetic diversity of Australian *Fusarium graminearum* and *F. pseudograminearum*. Plant Pathol. 55, 494-504.
- Akinsanmi O.A., Backhouse D., Simpfendorfer S., Chakraborty S. (2008). Mycelial compatibility reactions of australian *Fusarium graminearum* and *F. pseudograminearum* isolates compared with AFLP groupings. Plant Pathol. 57, 251-261.
- Alconada T.M. (1993). Enzimas de *Fusarium oxysporum f. sp. melonis* involucradas en la degradación de la pared celular de las plantas superiores. Tesis Doctoral FCE y N- UBA.
- Alconada T.M. and Martínez M.J. (1994). Purification and characterization of an extracellular endo-1,4- β -xylanase from *Fusarium oxysporum f. sp. melonis*. FEMS Microbiol Lett. 118, 305-310.
- Alconada T.M. and Martínez M.J. (1996). Purification and characterization of an β -glucosidase from *Fusarium oxysporum f. sp. melonis*. Lett Appl Microbiol. 22, 106-110.
- Alkorta I., Garbisu C., Llama M.J., Serra J.L. (1998). Industrial applications of pectic enzymes: a review. Process Biochem. 33, 21-28.
- Alvarez C.L., Azcarate M.P., Terminello L., Cabral D., Vaamonde G., Fernandez Pinto V. (2007) Genotypic diversity and mycotoxin production of *Fusarium graminearum* isolates from wheat cultivated in Argentina. Res Rev Biosci. 1, 172-180.
- Alvarez C.L., Somma S., Moretti A., Fernandez Pinto V. (2010). Aggressiveness of *Fusarium graminearum* sensu stricto Isolates in wheat kernels in Argentina. J Phytopathol. 158, 173-181.
- Annone J.G. (2003). Particularidades del control químico de la FET. Seminario: problemas asociados a la Fusariosis en trigo y estrategias para su prevención. Bolsa de Cereales de Buenos Aires 3-4 de julio de 2003.
- An H.J., Lurie S., Greve L.C., Rosenquist D., Kirmiz C., Labavitch J.M., Lebrilla C.B. (2005). Determination of pathogen-related enzyme action by mass spectrometry analysis of pectin breakdown products of plant cell walls. Anal. Biochem. 338, 71-82.

- Andersen A.L. (1948). The development of *Gibberella zeae* head blight of wheat. *Phytopathol.* 38, 599-611.
- Anderson J.B. and Stasovsky E. (1992). Molecular phylogeny of Northern Hemisphere species of *Armillaria*. *Mycologia* 84, 505-516.
- Annis, S. L. and Goodwin, P. H. (1997). Recent advances in the molecular genetics of plant cell wall-degrading enzymes produced by plant pathogenic fungi. *Eur J Plant Pathol* 103, 1-14.
- Aoki T. and O'Donnell K. (1999). Morphological and molecular characterisation of *Fusarium pseudograminearum* sp. nov., formally recognized as the Group 1 population of *F. graminearum*. *Mycologia* 91, 597- 609.
- Bai G.H. and Shaner G. (1994). Scab of wheat: prospects for control. *Plant Dis.* 78, 760-766.
- Bai G.H., Desjardins A.E., Lattner R.D. (2001). Deoxynivalenol nonproducing *Fusarium graminearum* causes initial infection, but does not cause disease spread in wheat spikes. *Mycopathologia* 153, 91-98.
- Bai G.H. and Shaner G.E. (2004). Management and resistance in wheat and barley to *Fusarium* head blight. *Annu Rev Phytopathol.* 42, 135-161.
- Barneix A.J. (2007). Physiology and biochemistry of source-regulated protein accumulation in the wheat grain. *J. Plant Physiol.* 164, 581- 590.
- Bateman G.L., Kwasna H., Ward E. (1996). Relationship among *Fusarium* spp. estimated by comparing restriction fragment length polymorphism in polymerase chain reaction amplified nuclear DNA. *Can J Microbiol.* 42, 1232-1240.
- Beardall J.M. and Miller J.D. (1994). Diseases in humans with mycotoxins as possible causes. In : Mycotoxins in Grain. Compounds Other Than aflatoxins. Miller J.D. and Trenholm H.L., eds. Eagan Press, St. Paul MN. Pages 487-540.
- Bennett J. and Klich M. (2003). Mycotoxins. *Clin Mycrobiol Rev.* 16, 497-516.
- Berto P., Comménil P., Belingheri L., Dehorter B. (1999). Occurrence of a lipase in spores of *Alternaria brassicicola* with a crucial role in the infection of cauliflower leaves. *FEMS Microbiol Lett.* 180, 183-189.
- Blinkovsky, A. M., Byun, T., Brown K, M. & Golightly, E. J. (1999). Purification, Characterization, and Heterologous Expression in *Fusarium venenatum* of a Novel Serine Carboxypeptidase from *Aspergillus oryzae* Applied and Environmental Microbiology, 3298- 3303.
- Blum H., Beier H., Gross H.J. (1987). Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels. *Electrophoresis* 8, 93-99.

Booth C. (1971). The genus *Fusarium*. Commonwealth Mycological Institute. Kew Surrey 35, 70-72.

Bowden R.L. and Leslie J.F. (1994). Diversity of *Gibberella zeae* (*Fusarium graminearum*) at small spatial scales. *Phytopathol.* 84, 1140

Bradford MM. (1976) A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye-binding. *Anal Biochem* (1976); 72:248-54.

Brennan J.M., Egan D., Cooke B.M., Doohan F.M. (2005) Effect of temperature on head blight of wheat caused by *Fusarium culmorum* and *F. graminearum*. *Plant Pathol.* 54, 156-160.

Broatch J. (2002). Alberta *Fusarium graminearum* Management Plan Agriculture Information Services Alberta (Canadá) [www.agric.gov.ab.ca/\\$departement/deptdocs.nsf/all](http://www.agric.gov.ab.ca/$departement/deptdocs.nsf/all). Verificado: 05/08/07

Bruker Optics. (1993). Sample Preparation, Measurement and Evaluation of the FT-IR Spectra of Microorganisms. Ettlingen, Germany: Bruker Optik GmbH.

Burgess L.W., Summerell B.A., Bullock S., Gott K.P., Backhouuse D. (1994). Laboratory Manual for *Fusarium* Research 3rd edn. University of Sydney, Sydney.

Bushnell W.R., Hazen B.E., Pritsch C. (2003). Histology and physiology of *Fusarium* head blight. In: Leonard K.J., Bushnell W.R., eds, *Fusarium Head Blight of Wheat and Barley*. APS Press, St. Paul, MN. Pages 44-83.

Busso C., Nobuyoshi Kaneshima E., de Assis Franco F., Boto Querol C., Alves de Castro-Prado M.A. (2007). Vegetative compatibility and molecular characterization of *Fusarium graminearum* isolates from the State of Paraná, Brazil. *Cienc Rur.* 37, 2-6.

Caprari, C., Bergmann, C., Miglieli, Q., Salvi, G., Albersheim, P., Darvill, A., Cervone, F. y De Lorenzo, G. (1993). *Fusarium moniliforme* secretes four endopolygalacturonases derived from a single gene product. *Physiol Mol Plant P* 43, 453-462.

Carpita N.C. and Gibeaut D.M. (1993). Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *Plant J.* 3, 1-30.

Carranza M.R., Moschini R.C., Kraan G., Bariffi J.H. (2007). Examination of meteorology-based predictions of *Fusarium* head blight of wheat grown at two locations in the southern Pampas region of Argentina. *Austral Plant Pathol.* 36, 1-4.

Carter J.P., Rezanoor H.N., Desjardins A.E., Nicholson P. (2000). Variation in *Fusarium graminearum* isolates from Nepal associated with their host of origin. Plant Pathol. 49, 452-460.

Carter J.P., Rezanoor H.N., Holden D., Desjardins A.E., Plattner R.D., Nicholson P. (2002). Variation in pathogenicity associated with the genetic diversity of *Fusarium graminearum*. Eur J Plant Pathol. 108, 573-583.

Chung W.H., Ishii H., Nishimura K., Ohshima M., Iwama T., Yoshimatsu H. (2008). Genetic analysis and PCR-based identification of major *Fusarium* species causing head blight on wheat in Japan. J Genet Plant Pathol. 74, 364-374.

Comménil P., Belingheri L., Dehorter B. (1998). Antilipase antibodies prevent infection of tomato leaves by *Botrytis cinerea*. Physiol Mol Plant Pathol. 52, 1-14.

Cumagun C.J.R., Bowden R.L., Jurgenson J.E., Leslie J.F., Miedaner T. (2004). Genetic mapping of pathogenicity and aggressiveness of *Gibberella zae* (*Fusarium graminearum*) toward wheat. Phytopathol. 94, 520-526.

Degan, F. D., Ribadeau-Dumas B., & K. Breddam. (1992). Purification and characterization of two serine carboxypeptidases from *Aspergillus niger* and their use in C-terminal sequencing of proteins and peptide synthesis. Applied Environmental and Microbiology, 58, 2144 - 2152.

Del Blanco I.A., Frohberg R.C., Stack R.W., Berzonsky W.A., Kianian S.F.(2003). Detection of QTL linked to Fusarium head blight resistance in Sumai 3-derived North Dakota bread wheat lines. Theoretical and Applied Genetics 106, 1027-1031

Degan, F. D., Ribadeau-Dumas B., & K. Breddam. (1992). Purification and characterization of two serine carboxypeptidases from *Aspergillus niger* and their use in C-terminal sequencing of proteins and peptide synthesis. Applied Environmental and Microbiology, 58, 2144 - 2152.

Del Ponte E.M., Fernandes J.M.C., Pierobom C.R., Bergstrom G.C. (2004). Giberela do trigo. Aspectos epidemiológicos e modelos de previsão. Fitopatologia Brasileira 29, 587-605.

Del Ponte E.M., Fernandes J.M.C., Pavan, W.A. (2005). A risk infection simulation model for *Fusarium* Head Blight of wheat. Fitopatologia Brasileira 30, 634-642.

Demeke T., Clear R.M, Patrick S.K., Gaba D. (2005). Species-specific PCR-based assays for the detection of *Fusarium* species and a comparison with the

whole seed agar plate method and trichothecene analysis. Int J Food Microbiol. 103, 271-284.

Desjardins A.E., Proctor R.H., Bai G., Shaner G., Hohon T.M. (1996). Reduced virulence of trichothecene- Nonproducing mutant of *G. zeae* in wheat field test. MPMI, 9, 775-781.

Detrixhe P., Chandelier A., Cavelier M., Buffet D., Oger R. (2003). Development of an agrometeorological model integrating leaf wetness duration estimation to assess the risk of head blight infection in wheat. Aspects of Applied Biology 68. Mycotoxins in food production systems.

De Lorenzo G., Castoria R., Bellicampi D., Cervone F. (1997). Fungal invasion enzymes and their inhibition. En The Mycota. Plant Relationships, Part B, pp. 61-83. Edited by: Carroll G.C. and Tudzinsky P. Berlin: Springer-Verlag

De Wolf E.D., Madden L.V., Lipps P.E. (2003). Risk assessment models for wheat *Fusarium* head blight epidemics based in within-season weather data. Phytopathol. 93, 428-435.

Díaz M.. (2006). Germoplasm Exchange in the Southern Cone of Latin America._The Global Fusarium Initiative for International Collaboration. Session 3: International scab nursery consortium, 103-108.

Di Pietro F., García-Maceira I., Méglecz E., Roncero M.I.G. (2001). A MAP kinase of the vascular wilt fungus is essential for root penetration and pathogenesis. Mol Microbiol. 39, 1140-1152.

Doohan F.M., Parry D.W., Jenkinson P., Nicholson P. (1998). The use of species-specific PCR-based assays to analyse *Fusarium* ear blight of wheat. Plant Pathol. 47, 109-205.

Dusabenyagasani M., Dostaler D., Hamelin R.C. (1999). Genetic diversity among *Fusarium graminearum* strains from Ontario and Quebec. Can J Plant Pathol. 21, 308-314.

Dyer R.B., Plattner R.D., Kendra D.F., Brown D.W. (2005). *Fusarium graminearum* TRII4 is required for high virulence and DON production on wheat but not for DON synthesis in vitro. J Agric Food Chem. 53, 9281-9287.

Edel V., Steinberg C., Avelange I., Lagurre G., Alabouvette C. (1995). Comparison of three molecular methods for the characterization of *Fusarium oxysporum* strains. Phytopathol. 85, 579-585.

Erukhimovitch V., Pavlov V., Talyshinsky M., Souprun Y., Huleihel M. (2005). FTIR microscopy assay method for identification of bacterial and fungal infections. *J Pharmaceut Biomed.* 37, 105-1108.

Fedak G., Cao. W., Xue A., Savard M., Gilbert J., Clarke J.and Somers D. (2006) Fusarium Head Blight Resistance From Wide Crosses in Bread Wheat and Durum. The Global Fusarium Initiative for International Collaboration. Session 3. International Scab Nursery Consortium. 20-23.

Feng J., Liu G., Selvaraj G., Hughes G.R., Wei Y. (2005). A secreted lipase encoded by LIP1 is necessary for efficient use of saturated triglyceride lipids in *Fusarium graminearum*. *Microbiol.* 151, 3911-3921.

Fischer G., Braun S., Thissen R., Dott W. (2006). FT-IR spectroscopy as a tool for rapid identification and intra-species characterization of airborne filamentous fungi. *J Microbiol Methods.*64(1):63-77

Franz E., Booij K., Vander Fels-Klerx I. (2009). Prediction of Deoxynivalenol content in Dutch winter wheat. *J Food Protect.* 72, 2170-2177.

Gagkaeva T.Y. and Yli-Mattila T. (2004). Genetic diversity of *Fusarium graminearum* in Europe and Asia. *Eur J Plant Pathol.* 110, 551-562.

Gale R.L, Chen L.F., Hernick C.A, Takamura K., Kistler H.C. (2002). Population analysis of *Fusarium graminearum* from wheat fields in eastern China. *Phytopathol.* 92, 1315-1322.

Galich de M.T.V. (1989). Importancia y difusión de la fusariosis del trigo en Argentina. En: M.M. Kohli, ed. Taller sobre Fusariosis de la espiga en América del Sur. México, D.F.: CIMMYT. Pages 7-26.

Galich de M.T.V. and Galich, A.N. (1994). Enfermedades del trigo en el área sur de Santa Fé y Córdoba. Información para extensión 5. EEA-INTA Marcos Juárez, Córdoba province, Argentina.

Galich A.N. and Galich M.T.V. de. (1996). Enfermedades del trigo en el área central norte de la región cerealera argentina. EEA INTA Marcos Juárez. Informe Técnico N°121. Pages 1-25.

Galich M.T.V de. (1997). *Fusarium Head Blight in Argentina*. En Dubin H.J., Gilchrist J., Reeves and Mc Nab A. eds. *Fusarium Head Scab: Global Status and Future Prospects*. México, D.F.: CIMMYT

Galich M.T. V. de (2004). Fusariosis de la espiga del trigo. Desarrollo de cultivares resistentes. IDIA XXI . INTA Año IV N°6. Pages 50-57.

Garcia-Maceira, F. I., Di Pietro, A. y Roncero, M. I. (2000). Cloning and disruption of *pgx4* encoding an in planta expressed exopolygalacturonase from *Fusarium oxysporum*. *Mol Plant Microbe Interact* 13, 359-365.

Garcia-Maceira, F. I., Di Pietro, A., Huertas-Gonzalez, M. D., Ruiz-Roldan, M. C. y Roncero, M. I. (2001). Molecular characterization of an endopolygalacturonase from *Fusarium oxysporum* expressed during early stages of infection. *Appl Environ Microbiol* 67, 2191-2196.

Gerlach W. and Nirenberg H. (1982). The genus *Fusarium* –a Pictorial Atlas. Mitt Biol Bundesanst. Ld – u Forstw. Berlin- Dahlem. 209. Pages 1-406.

Gilbert J., Abramson D., McCallum B., Clear R. (2001). Comparison of Canadian *Fusarium graminearum* isolates for aggressiveness, vegetative compatibility, and production of ergosterol and mycotoxins. *Mycopathologia* 153, 209-215.

Gilbert J. and Fernando W.G.D. (2004). Epidemiology and biological control of *Gibberella zae/Fusarium graminearum*. *Can J Plant Pathol.* 26, 464-472.

Gómez-Gómez E., Roncero M.I.G., Di Pietro A., Hera C. (2001). Molecular characterization of a novel endo-F-1,4-xylanase gene from the vascular wilt fungus *Fusarium oxysporum*. *Curr Genet.* 40, 268-275.

Goswami R.S. and Kistler H.C. (2004). Heading for disaster: *Fusarium graminearum* on cereal crops. *Mol. Plant Pathol.* 5, 515-525.

Guadet J., Julien J., Lafay J.F., Brygoo Y. (1989). Phylogeny of some *Fusarium* species, as determined by large-subunit rRNA sequence comparison. *Mol Biol Ecol.* 6, 227-242.

Guevara, M. A., M.T., G.-J. y Estévez, P. (1997). Multiple forms of pectic lyases and polygalacturonase from *Fusarium oxysporum* f.sp. *radicis lycopersici*: regulation of their synthesis by galacturonic acid. *Can J Microbiol* 43, 245-253.

Hare M.C., Parry D.W., Baker M.D. (1999). The relationship between wheat seed weight, infection by *Fusarium culmorum* or *Microdochium nivale*, germination and seedling disease. *Eur J Plant Pathol.* 105, 859-866.

Harris L.J., Desjardins A.E., Plattner R., Nicholson P., Butler G., Young J., Weston G., Proctor R.H., Hohn T.M. Possible Role of Trichothecene Mycotoxins in Virulence of *Fusarium graminearum* on Maize. *Plant Dis.* 83, 954-960.

Hatshc D.H., Phalip V., Petkovski E., Jeltsch J.M. (2006). *Fusarium graminearum* on plant cell wall: No fewer than 30 xylanase genes transcribed. *Biochem Biophys Res Co.* 345, 959-966.

- Hayashida D., Ohta K., Mo K. (1988). Xylanase of *Talaromyces byssochlamydoides*. Methods Enzymol. 160, 675-678.
- Hellweg M. (2003). Molecular biological and biochemical studies of proteolytic enzymes of the cereal pathogen *Fusarium graminearum*. Inaugural Dissertation Der Westfälischen Wilhelms-Universität Münster.
- Helm D. and Naumann D. (1995). Identification of some bacterial cell components by FT-IR spectroscopy. FEMS Microbiol Lett 126, 75-80.
- Helm D, Labischinski H, Naumann D. (1991)a. Elaboration of procedure for identification of bacteria using Fourier-Transform IR spectral libraries: a stepwise correlation approach. J Microbiol Meth 14:127-142.
- Helm, D., Labischinsky, H., Schallehn, G., and Naumann, D. (1991)b. Classificaton and identification of bacteria by Fourier-transform infrared spectroscopy. J.Gen.Microbiol. 137, 69-79.
- Hestbjerg H., Nielsen K.F., Thrane U., Elmholz S. (2002). Production of trichothecenes and other secondary metabolites by *Fusarium culmorum* and *Fusarium equiseti* on common laboratory media and a soil organic matter agar: An ecological interpretation. J Agricul Food Chem. 50, 7593-7599.
- Hinojo M.J., Llorens A., Mateo R., Patiño B., Gonzalea M.T., Jimenez M. (2004). Utility of the PCR- RFLP of the IGS of the rDNA for characterizing *Gibberella fujikuroi* isolates. Syst Appl Microbiol. 27, 681-688.
- Hooker D.C., Schaafsma A.W., Tamburinc-Illincic L. (2002). Using weather variables pre and post-heading to predicting deoxinivalenol content in winter wheat. Plant Dis. 86, 611-619.
- Huang, Q. y Allen, C. (2000). Polygalacturonases are required for rapid colonization and full virulence of *Ralstonia solanacearum*. Physiol Mol Plant P 57, 77-83.
- Jarvis M. C., Forsyth W., Duncan H. (1988). A Survey of the Pectic Content of Nonlignified Monocot Cell Walls. J. Plant Physiol., 88, 309-314.
- Jayani R.S, Saxena S., Gupta R. (2005). Microbial pectinolytic enzymes: A review. Process Biochem. 40, 2931-2944.
- Jenczmionka N.J., Maier F.J., Losch A.P., Schafer F.J. (2003). Mating, conidiation and pathogenicity of *Fusarium graminearum*, the main causal agent of the head-blight disease of wheat, are regulated by the MAP kinase gpmk1. Curr Genet. 43, 87-95.

Jenczmionka N.J. and Schäfer W. (2005). The Gpmk1 MAP kinase of *Fusarium graminearum* regulates the induction of specific secreted enzymes. Curr Genet. 47, 29-36.

Kang Z. and Buchenauer H. (2000a). Ultrastructural and cytochemical studies on cellulose, xylan and pectin degradation in wheat spikes infected by *Fusarium culmorum*. J Phytopatol. 148, 263-275.

Kang Z. and Buchenauer H. (2000b). Cytology and ultrastructure of the infection of wheat spikes by *Fusarium culmorum*. Mycol Res. 104, 1083-1093.

Kang Z., Zingen-Sell I., Buchenauer H. (2005). Infection of wheat spikes by *Fusarium avenaceum* and alterations of a cell wall components in the infected tissue. Eur J Plant Pathol. 111, 19-28.

Kang Z., Buchenauer H., Huang L., Han Q. and Zhang H. (2007). Cytological and immunocytochemical studies on responses of wheat spikes of the resistant Chinese cv. Sumai 3 and the susceptible cv.Xiaoyan 22 to infection by *Fusarium graminearum*. European Journal of Plant Pathology. 120, (4), 383-396.

Kashyap D.R., Vohra P.K., Chopra S., Tewari R. (2001). Applications of pectinases in the commercial sector: a review. Biores Technol. 77, 215-227.

Kieliszewski M.J. and Lamport T.A. (1994). Extensin: repetitive motifs, functional sites, post-translational codes, and phylogeny. The Plant J. 5, 157-172.

Kikot G.E., Hours R.A., Alconada T.M. (2009). Contribution of cell wall degrading enzymes to pathogenesis of *Fusarium graminearum*: a review. J Basic Microbiol. 49, 231-241.

Kikot G., Hours R., Alconada T. (2010). Extracellular enzymes of *Fusarium graminearum* isolates. Brazilian Arch Biol Technol. 53, 779-783.

Kikot G., Moschini R., Consolo F., Rojo R., Hours R., Salerno G.,Gasoni L., Arambarri M.A., Alconada T. M. (2011). Occurrence of *Fusarium graminearum* from wheat in relation to climatic models in Argentina Pampas Region. Mycopathologia 171, 139-149.

Koopmann B., Karlovsky P., Wolf G. (1994). Differentiation between *Fusarium culmorum* and *Fusarium graminearum* by RFLP and with species-specific DNA probes. In: Schots A., Dewey F.M., Oliver R., eds Modern Assays for Plant Pathogenic Fungi, Identification, Detection and Quantification. Wallingford, UK: CAB International. Pages 37-46.

Krause R.A. and Massie L.B. (1975). Predictive systems: Modern approaches to disease control. Annu Rev Phytopathol 13, 31-47.

Kruger W.M., Pritsch C., Chao S., Muehlbauer G.J. (2002). Functional and comparative bioinformatic analysis of expressed genes from wheat spikes infected with *Fusarium graminearum*. Mol Plant Microbe Int. 15, 445-455.

Láday M., Juhász Á., Mulé G., Moretti A., Logrieco A. (2004). Mitochondrial DNA diversity and lineage determination of European isolates of *Fusarium graminearum* (*Gibberella zaeae*). Eur J Plant Pathol. 110, 545-550.

Lang C. and Dörnenburg H. (2000). Perspectives in the biological function and the technological application of polygalacturonases. Appl Microbiol Biotechnol. 53, 366-375.

Lazzari F.A. (2000). Control integrado de plagas, manejo de hongos e insectos. Granos y Post-cosecha Latinoamericana VI nº XXIII.

Lemmens M., Haim K., Lew H., Ruckenbauer P. (2004) The effect of nitrogen fertiliser on *Fusarium* head blight development and deoxynivalenol contamination in wheat. J Phytopathol. 152, 1-8.

Leslie J.F., Anderson L.L., Bowden R.L., Lee Y.W. (2007). Inter13 and intra-specific genetic variation in *Fusarium*. Int J Food Microbiol. 119, 25-32.

Leslie J.F. and Summerell B.A. (2006). The *Fusarium* laboratory manual. First Edition Backwell Publishing.

Lev S. and Horwitz B.A. (2003). A mitogen-activated protein kinase pathway modulates the expression of two cellulase genes in *Cochlobolus heterostrophus* during plant infection. Plant Cell. 5, 835-844.

Llorens A., Hinojo M., Mateo R., Gonzalez M.T., Valle Alagarr F.M., Jimenez M. (2006). Characterization of *Fusarium* spp. isolates by PCR- RFLP analysis of the intergenic spacer región of the rRNA gene. Int J Food Microbiol 103, 297-306.

Lori G.A. (1985).Género *Fusarium* en Argentina II. Identificación de cultivos en la zona de La Plata. Revista Argentina de Microbiología, 17, 61-67.

Lori G.A., Sisterna, M.N., Haldukowski, M., Rizzo I. (2004). *Fusarium graminearum* and deoxynivalenol contamination in the durum wheat area of Argentina. Microbiol Research. 158, 29-35.

Lori G.A., Sisterna M.N., Sarandón S.J., Rizzo I., Chidichimo H. (2009). *Fusarium* head blight in wheat: impact of tillage and other agronomic practices under natural infection. Crop Protect. 28, 495-502.

MacCallum B.D., Tekauz A., Gilbert J. (2001). Vegetative compatibility among *Fusarium graminearum* (*Gibberella zaeae*) isolates from barley spikes in Southern Manitoba. Can J Plant Pathol. 23, 83-87.

Maquelin, K., L. P. Choo-Smith, C. Kirschner, N. A. Ngo Thi, D. Naumann, G. Puppels. (2002). Vibrational spectroscopic studies of microorganisms. In J. M. Chalmers and P. R. Griffiths (ed.), *Handbook of vibrational spectroscopy*. John Wiley & Sons Ltd., Chichester, United Kingdom. Pages 1-27

Martínez M.J., Alconada T.M., Guillén F., Vázquez C., Reyes F. (1991). Pectic activities from *Fusarium oxysporum f.sp. melonis*. Purification and characterization of an exopolysaccharidase. *FEMS Microbiol Lett.* 81, 145-150.

Martínez M.I. (2009). Efecto del ambiente sobre la intensidad de la Fusarosis de la espiga de trigo y el contenido de Deoxinivalenol en grano. Tesis Magister de la Universidad de Buenos Aires, área: Producción Vegetal.

Mayorga M.E. and Gold S.E. (1999). A Map kinase encode by the ubc3 gene of *Ustilago maydis*. *Fungal Genetics Biol.* 34, 485-497.

McMullen M., Jones R., Gallemberg D. (1997). Scab of wheat and barley: a re-emerging disease of devastating impact. *Plant Dis.* 81, 1340-1348.

Mentewab A., Rezanoor H.N., Gosman N., Worland A.J., Nicholson P. (2000). Chromosomal location of *Fusarium* head blight resistance genes and analysis of the relationship between resistance to head blight and brown foot rot. *Plant Breed* 119, 15-20.

MEROPS <http://merops.sanger.ac.uk/>

Mesterházy A., Bartók T., Mirocha C.G., Komoróczy R. (1999). Nature of wheat resistance to *Fusarium* head blight and the role of deoxynivalenol for breeding. *Plant Breeding* 118, 97-110.

Miedaner T. and Schilling A.G. (1996). Genetic variation of aggressiveness in individual field populations of *Fusarium graminearum* and *Fusarium culmorum* tested on young plants of winter rye. *Eur J Plant Pathol.* 102, 823-830.

Miedaner T., Schilling A.G., Geiger H.H. (2001). Molecular genetic diversity and variation for aggressiveness in populations of *Fusarium graminearum* and *Fusarium culmorum* sampled from wheat field in different countries. *J Phytopathol.* 149, 641-648.

Mirete Castaneda S. (1999). Análisis mediante la técnica PCR-RFLP de la región espaciadora inter génica (IGS) del rDNA en el género *Fusarium*. Graduation Thesis. Universidad Complutense de Madrid, Spain.

Mishra P.K., Fox R.T.V., Culham A. (2002). Restriction analysis of PCR amplified nrDNA regions revealed intraspecific variation within populations of *F. culmorum*. *FEMS Microbiol Lett.* 215, 291-296.

Mishra P.K., Fox R.T.V., Culham A. (2003). Inter-simple sequence repeat and gressiveness analyses revealed high genetic diversity, recombination and long-range dispersal in *Fusarium culmorum*. *Annals Appl Biol.* 143, 291-301.

Mishra P.K., Tewari J.P., Clear R.M., Turkington T.K. (2004). Molecular genetic variation and geographical structuring in *Fusarium graminearum*. *Annals Appl Biol.* 145, 299-307.

Mishra P.K., Tewari J.P., Clear R.M., Turkington T.K. (2006). Genetic diversity and recombination within populations of *Fusarium pseudograminearum* from western Canada. *Int Microbiol.* 9, 65-68.

Moeller E.M., Chelkowski J., Geiger H.H. (1999). Species-specific PCR assays for the fungal pathogens *F. moniliforme* and *F. subglutinans* and their application to diagnose maize ear rot. *J Phytopathol.* 147, 497-508.

Monds R.D., Crome M.G., Lauren D.R., di Menna M., Marshall J. (2005). *Fusarium graminearum*, *F. cortaderiae* and *F. pseudograminearum* in New Zealand: molecular phylogenetic analysis, mycotoxin chemotypes and co-existence of species. *Mycol Res.* 109, 410-420.

Moon J.H., Lee Y.H., Lee Y.W. (1999). Vegetative compatibility groups in *Fusarium graminearum* isolates from corn and barley in Korea. *Plant Pathol J.* 15, 53-56.

Moschini R.C., Fortugno C. (1996). Predicting wheat head blight incidence using models based on meteorological factors in Pergamino, Argentina. *European J Plant Pathol.* 102, 211-218.

Moschini R.C. and Carmona M. (1998). Fusariosis en trigo. Nuevo enfoque para su control para el área de Balcarce. *Revista ALEA Informa.* Año 1 Nº 8. Pages 16-20.

Moschini R.C., Pioli R., Carmona M. and Sacchi O. (2001). Empirical predictions of wheat head blight in the northern Argentinean Pampas region. *Crop Sci.* 41, 1541-1545.

Moschini R.C., Galich M.T.V. de, Annone J.G., Polidoro O. (2002). Enfoque Fundamental-Empírico para estimar la evolución del Índice de *Fusarium* en trigo. *Revista RIA,* 31, 39-53.

Moschini R.C; Carranza M.R., Carmona M. (2004). Meteorological-based predictions of wheat head blight epidemic in the southern argentinean pampas region. *Cereal Res Commun.* 32, 45-52.

Moschini R.C., Castellarín J., González M., Ferraguti F., Martínez M. (2008). Validación de los valores estimados de la Fusariosis de la espiga de trigo por sistemas de pronóstico basados en variables meteorológicas, en Oliveros (campaña 2007/08). VII Congreso Nacional de Trigo

Murray M.G. and Thompson W.F. (1980). Rapid isolation of high molecular weight plant DNA. Nucleic Acids Res. 8, 4321-4325.

Naef A. and Defago, G. (2006). Population structure of plant-pathogenic *Fusarium* species in overwintered stalk residues from Bt-transformed and non-transformed maize crops. Europ J Plant Pathol, 116, 129-143.

Naumann D, Helm D, Labischinski H. (1991). Microbiological characterizations by FT-IR spectroscopy. Nature 351:81-82.

Naumann D. (2000).in: R. A. Meyers (ed), Encyclopedia of Analytical Chemistry, John Wiley & Sons Ltd, eds, Chichester, pp. 102-131.

Naumann, D. (2001). FT-IR and FT-NIR Raman Spectroscopy in Biomedical Research. Gremllich, Hans-Ulrich and Yang, Bing. [24], 323-377 USA, Marcel Dekker, Inc. Practical Spectroscopy.

Naumann A., Navarro-González M., Peddireddi S., Kües U., Polle A. (2008) Fourier transform infrared microscopy and imaging: Detection of fungi in wood. Fungal Genetics and Biology 42, 829–835.

Nelson P.E., Toussoun T.A., Marasas W.F.O. (1983). *Fusarium* Species: An illustrated Manual for Identification. Pennsylvania State University Press, University Park, Pennsylvania.

Nelson N. (1944). A phtometric adaptation of the Somogyi method for determination of glucose. J Biol Chem. 153, 376-380.

Neuhoff V., Arnold N., Taube D., Ehrhardt W. (1988). An improved procedure for staining of proteins following separation in polyacrylamide gels in described which utilizes the colloidal properties of Coomassie Brilliant Blue G- 250 and R-250. Electrophoresis 9, 255-262.

Nichols PD, Henson JM, Guckert JB, Nivens DE, White DC. (1995). Fourier transform-infrared spectroscopic methods for microbial ecology: analysis of bacteria, bacteria-polymer mixtures and biofilms. J Microbiol Methods 4:79-94.

Nicholson P., Jenkinson P., Rezanoor H.N., Parry D.W. (1993). Restriction fragment length polymorphism analysis of variation in *Fusarium* species causing ear blight of cereals. Plant Pathol. 42, 905-914.

Nicholson P., Simpson D.R., Weston G., Rezanoor H.N., Lees A.K., Parry D.W., Joyce D. (1998). Detection and quantification of *Fusarium culmorum* and *Fusarium graminearum* in cereal using PCR assays. *Physiol Mol Plant Pathol.* 53, 17-37.

Nie M. and Li W.H. (1979). Mathematical model for studing genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA* 70, 5269-5273.

Nie M., Zhang W.Q., Xiao M., Luo J.L., Bao K., Chen J.K., Li B. (2007a). FT-IR Spectroscopy and artificial neural network identification of *Fusarium* species. *J Phytopathol.* 115, 364-367.

Nie M., Luo J., Xiao M., Chen J., Bao K., Zhang W., Chen J., Li B. (2007b). Structural differences between *Fusarium* strains investigated by FTIR spectroscopy. *Biochem.* 72, 61-67.

Niessen L. and Vogel R.F. (1998). Group specific PCR- detection of potential trichothecene producing *Fusarium* species in pure cultures and cereal samples. *Syst Appl Microbiol.* 21, 618-631.

Nightingale M.J., Marchylo B.A., Clear R.M., Dexter J.E., Preston K.R. (1999). *Fusarium* head blight: Effect of fungal proteases on wheat storage proteins. *Cereal Chem.* 76, 150-158.

Niture S.K., Kumar A.R., Pant A. (2006). Role of glucose in production and repression of polygalacturonase and pectate lyase from phytopathogenic fungus *Fusarium moniliforme* NCIM 1276. *World J Microb Biotech.* 22, 893-899.

O'Donnell K., Cigelnik E., Casper H.H. (1998). Molecular, phylogenetic, morphological, and mycotoxin data support reidentification of the Quorn mycoprotein fungus as *Fusarium venenatum*. *Fungal Genet Biol.* 23, 57-67.

O'Donnell K., Kistler H.C., Tacke B.K., Casper H.H. (2000). Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proc Natl Acad Sci. USA.* 97, 7905-7910.

O'Donnell K., Ward T.J., Geiser D.M., Kistler H.C., Aoki T. (2004). Genealogical concordance between the mating type locus and seven other nuclear genes supports formal recognition of nine phylogenetically distinct species within the *Fusarium graminearum* clade. *Fungal Genet Biol.* 41, 600-623.

Oeser B., Heidrich P.M., Müller U., Tudzynski P., Tenberge K.B. (2002). Polygalacturonase is a pathogenicity factor in the *Claviceps purpurea*/rye interaction. *Fungal Genet Biol.* 36, 176-186.

- Osborne L.E. and Stein J.M. (2007). Epidemiology of *Fusarium* head blight on small-grain cereals. *Int J Food Microbiol.* 119, 103-108.
- Ouellet T., Siefert K.A. (1993). Genetic characterisation of *Fusarium* strains using RAPD and PCR amplification. *Phytopathol.* 83, 1003-1007.
- Panda T., Nair S.R., Kumar P. (2004). Regulation of synthesis of the pectolytic enzymes of *Aspergillus niger*. *Enzyme Microb Tech.* 34, 466-473.
- Patiño B., Mirete S., Vazquez C., Jimenez M., Rodriguez M.T., Gonzales M.T. (2006). Characterization of *F. verticillioides* strains by PCR- RFLP analysis of the IGS of the rDNA. *J Sci Food Agric.* 86, 420-435.
- Parry D.W., Jenkinson P., McLeod L. (1995). "Fusarium ear blight (scab) in small grain cereals - a review." *Plant Pathol.* 44, 207-238.
- Parry D.W. and Nicholson P. (1996). Development of a PCR assay to detect *Fusarium poae* in wheat. *Plant Pathol.* 45, 383-391.
- Pekkarinen A.I., Mannonen L., Jones B.L., Niku-Paavola M.L. (2000). Production of proteases by *Fusarium* species grown on barley grains and in media containing cereal proteins. *J Cereal Sci.* 31, 253-261.
- Pekkarinen A.I., Jones B.L., Niku-Paavola M.L. (2002). Purification and properties of an alkaline proteinase of *Fusarium culmorum*. *Eur J Biochem.* 269, 798-807.
- Pekkarinen A.I. (2003). The serine proteinases of *Fusarium* grown on cereal proteins and in barley grain and their inhibition by barley proteins. Academic dissertation, University of Wisconsin, Madison, USA.
- Pekkarinen, A.I., Longstaff C. & Jones B.L. (2007). Kinetics of the Inhibition of *Fusarium* Serine Proteinases by Barley (*Hordeum vulgare* L.) Inhibitors. *Journal of Agricultural and Food Chemistry,* 55, 2736-2742.
- Peraica M., Radic B., Lucic A., Pavlovic M. (1999). Toxic effects of mycotoxins in humans. *Bulletin of World Health Organ.* 77, 754-766.
- Phalip V., Delande F., Carapito Ch., Goubet F., Hatsch D., Leize Wagner E., Dupree P., Van Dorsselaer A., Jetsch J.M. (2005). Diversity of the exoproteome of *Fusarium graminearum* grown on plant cell wall. *Curr Genet.* 48, 366-379.
- Powell A.L.T., van Kan J.A.L., ten Have A., Visser J., Greve L.C., Bennett A.B., Labavitch J.M. (2000). Transgenic expression of pear PGIP in tomato limits colonization. *Mol Plant Microbe Interact.* 13, 942-950.

Prandini A., Sigolo S., Filippi L., Battilani P., Piva G. (2009). Review of predictive models for *Fusarium* head blight and related mycotoxin contamination in wheat. *Food Chem Toxicol.* 47, 927-931.

Pritsch C., Muehlbauer G.J., Bushnell W.R., Somers D.A., Vance C.P. (2000). Fungal development and induction of defense response genes during early infection of wheat spikes by *Fusarium graminearum*. *Mol Plant Microbiol Interact.* 13, 159-169.

Proctor R.H., Honh T.M., McCormick S.P. (1995). Reduced virulence of *G. zeae* caused by disruption of a trichothecene toxin biosynthetic gene. *Mol Plant Microbe Interact.* 8, 593-601.

Qu B., Li H.P., Zhang J.B., Xu Y.B., Huang T., Wu A.B., Zhao C.S., Carter J., Nicholson P., Liao Y.C. (2008). Geographic distribution and genetic diversity of *Fusarium graminearum* and *F. asiaticum* on wheat spikes throughout China. *Plant Pathol.* 57, 15-24.

Quiroga E. N., Sgariglia M. A., Molina C. F., Sampietro, D. A., Soberón J. R., Vattuone M. A. (2009) Purification and characterization of an exo-polygalacturonase from *Pycnoporus sanguineus*. *Mycological research.* 113 (12), 1404- 1410.

Ramírez M.L., Reynoso M.M., Farnochi M.C., Chulze S. (2006). Vegetative compatibility and mycotoxin chemotypes among *Fusarium graminearum* (*Gibberella zeae*) isolates from wheat in Argentina. *Europ J Plant Pathol.* 115, 139-148.

Rebuffo, C., J. Schmitt, M. Wenning, F. von Stetten, and S. Scherer. (2006) Reliable and rapid identification of *Listeria monocytogenes* and *Listeria* species by artificial neural network-based Fourier transform infrared spectroscopy. *Appl. Environ. Microbiol.* 72:994–1000.

Reis E.M. and Carmona M. (2002). Fusariosis del trigo. Biología, epidemiología y estrategias para su manejo. Pages 25.

Ribichich K. and Vegetti A.C. (2001). Fusariosis de la espiga de trigo: evaluación de caracteres exomorfológicos asociados a la resistencia. *Revista de la Facultad de Agronomía, La Plata.* 104, 121-127.

Riou Ch.; Freyssinet, G.; Fevre, M. (1992). Purification and characterization of extracellular pectinolytic enzymes produced by *Sclerotinia sclerotiorum*. *Appl. Environ. Microbiol.*, 58(2), 578- 583.

Rombouts F.M. and Pilnik W. (1980). In: *Economic microbiology*. Rose H. (ed.) Academic press. 5: 227- 282.

Roncero M.I.G., Hera C., Ruiz-Rubio M., García-Maceira F.I., Madrid M.P., Caracuel Z., Calero F., Delgado-Jarana J., Roldán-Rodriguez R., Martínez-Rocha A.L. (2003). *Fusarium* as a model for studying virulence in soilborne plant pathogens. *Physiol. Mol Plant Pathol.* 62, 87-98.

Rossi V., Giousue S., Pattori E., Spanna F., Del Vechio A. (2003). A model estimating the risk of *Fusarium* head blight on wheat. *EPPO Bulletin.* 33, 421-425.

Ruiz-Roldán M.C., Maier F.J., Shäfer W. (2001). PTK1, a mitogen-activated-protein kinase gene, is required for conidiation, appressorium formation, and pathogenicity of on barley. *Mol Plant Microbe Interact.* 14, 116-125.

Rohlf F.J. (1998). NTSYS-pc. Numerical taxonomy and multivariate analysis system. Setauket, NY: Software.

Saharan M.S., Kumar J., Sharma A.K., Nagarajan S. (2004). Proc. Indian Natl.Sci.Acad. B70 N°3. Pages 255-268.

Saharan,M.S., Naef A., Kumar J., Tiwari R. (2007). Characterization of variability among isolates of *Fusarium graminearum* associated with head scab of wheat using DNA markers. *Curr Sci.* 92, 230-235.

Saharan M.S. and Naef A. (2008). Detection of genetic variation among Indian wheat head scab pathogens (*Fusarium* spp./isolates) with microsatellite markers. *Crop Protect.* 27, 1148-1154.

Salman A., Tsror L., Pomerantz A., Morehc R., Mordechai S., Huleihel M. (2010). FTIR spectroscopy for detection and identification of fungal phytopathogens. *Spectroscopy.* 0, 1-7

Salman A., Tsror L., Pomerantz A., Lapidet A., Morehc R., Mordechai S., Huleihel M. (2011). Distinction of *Fusarium oxysporum* fungal isolates using FTIR-ATR spectroscopy and advanced statistical methods. *Analyst,* 136, 988- 995.

Salman A., Tsror L., Pomerantz A., Morehc R., Mordechai S., Huleihel M. (2012). Identification of fungal phytopathogens using ATR-FTIR spectroscopy and advanced statistical methods. *J of Biomedical Optics,* 17 (1), 1-8.

Samuels G.J., Niremberg H.I., Seifert K.A. (2001). Perithecial species of *Fusarium*, En: Summerell B.A., Leslie J.F., Backhouse D., Bryden W.L., Burgess L.W.(Eds). *Fusarium: Paul E. Nelson Memorial Symposium.* APS Press, St. Paul, Minnesota. Pages 1-14.

Schaafsma A.W., Tamburic-Ilinic Miller J.D., Hooker D.C. (2001). Agronomic consideration for reducing DON in wheat grain. *Can.J Plant Pathol.* 23, 279-285.

Schilling A.G., Moller E.M., Geiger H.H. (1994). RAPDs of *Fusarium culmorum* and *F. graminearum*. Application for genotyping and species identification. In: Schots A, Dewey, F.M. and Oliver, R.P. (Ed.), Modern Assays for Plant Pathogenic Fungi CAS International, Oxford, UK. Pages 47-56.

Schilling A.G., Möller E.M., Geiger H.G. (1996). Polymerase chain reaction-based assays for species-specific detection of *Fusarium culmorum*, *F. graminearum* and *F. avenaceum*. *Phytopathol.* 86, 515-22.

Schilling G.A., Miedaner T., Geiger H.H. (1997). Molecular variation and genetic structure in field populations of *Fusarium* species causing head blight in wheat. *Cereal Res Commun.* 25, 549-554.

Schmitt, J., and T. Udelhoven. (2001). Use of artificial neural networks in biomedical diagnosis. In H.-U. Gremlich and B. Yang (ed.), Infrared and Raman spectroscopy of biological material. Marcel Dekker, Inc., New York, NY. , p. 379–419

Schwarz P.B., Schwarz J.G., Zhou A., Prom L.K., Steffenson B.J. (2001). Effect of *Fusarium graminearum* and *F. poae* infection on barley and malt quality. *Montsschr Brauwiss.* 54, 55-63.

Schwarz P.B., Jones B.L., Steffenson B.J. (2002). Enzymes associated with *Fusarium* infection of barley. *J Amer Soc Brewing Chem.* 60, 130-134.

Scott-Craig, J. S., Cheng, Y. Q., Cervone, F., De Lorenzo, G., Pitkin, J. W., Walton, J. D. (1998). Targeted mutants of *Cochliobolus carbonum* lacking the two major extracellular polygalacturonases. *Appl Environ Microbiol* 64, 1497-1503.

Seong K., Hou Z., Tracy M., Kistler H.C., Xu J.R. (2005). Random insertional mutagenesis identifies genes associated with virulence in the wheat scab fungus *Fusarium graminearum*. *Phytopathol.* 95, 744-750.

Shieh M., Brown R.L., Whitehead M.P., Carey J.W., Cotty P.L., Cleveland T.E., Dean R.A. (1997). Molecular genetic evidence for the involvement of a specific polygalacturonase P2c, in the invasion and spread of *Aspergillus flavus* in cotton bolls. *Appl Environ Microbiol.* 63, 3548-3552.

Shimokawa T., Kakegawa K., Ishii T. (2002). Feruloyl esterases from suspension-cultured rice cells. *Bull. FFPRI* 1, 225-230.

Sneath and Sokal (1973). Sneath PHA & Sokal RR (1973) Numerical Taxonomy. Freeman, San Francisco.

Singh Jayani R., Saxena S., Gupta R. (2005). Microbial pectinolytic enzymes: A review. *Process Biochem.* 40, 2931-2944.

Smith B.J. Methods in Molecular Biology. 1984. Walker J.M.E. (ed.). Clifton, New Jersey. Pages 41-56

Snijders C.H.A. (1990). Systemic fungal growth of *Fusarium culmorum* in stems of winter wheat. J Phytopathol. 129, 133-140.

Snijders C.H.A.. (1990). Genetic variation for resistance to Fusarium head blight in bread wheat. Euphytica, 50 (2), 171-179

Snijders C.H.A., Krechting C.F. (1992). Inhibition of deoxynivalenol translocation and fungal colonization in *Fusarium* head blight resistant wheat. Can J Bot. 70, 1570-1576.

Snijders C.H.A.. (2004). Resistance in wheat to *Fusarium* infection and trichothecene formation. Toxicol Lett. 153, 37-46.

Sologuk S. (2005). Diseases Can Bewitch Durum Millers *Fusarium* head blight, ergot, and black point are the most common wheat-related fungi. Milling Journal. Pages 44-45. www.northern-crops.com/technical/durumdisease.pdf. Verificado 20/07/07.

Stack R.W. (2000). Return of an Old Problem: Fusarium Head Blight of Small Grains. www.apsnet.org/education/feature/FHB. Verificado 03/07/07

Ten Have A., Mulder W., Visser J.N., van Kan A.L. (1998). The endopolygalacturonase gene Bcpg1 is required for full virulence of *Botrytis cinerea*. Mol Plant Microbe Interact. 11, 1009-1016.

Ten Have A., Tenberge K.B., Benen J..A.E., Tudzynski P., Visser J., van Kan A.L. (2002). The contribution of cell wall degrading enzymes to pathogenesis of fungal plant pathogens. In: The Mycota XI Agricultural Application. Kempken F. (Ed). Springer Verlag, Berlín Heidelberg. Pages 341- 358.

Teramoto Y., Saigusa N., Yoshida Y., Ueda S., Yosizawa K. (1994) Produccion and characteristics of red rice sake. J. Inst. Brew 100, 3- 6

Thakur A., Pahwa R., Singh S., Gupta R.. (2010). Production, Purification, and Characterization of Polygalacturonase from *Mucor circinelloides* ITCC 6025. Enzime research. 1-7.

Tomassini, A., Sella, L., Raiola, A., D'Ovidio, R. and Favaron, F. (2009). Characterization and expression of *Fusarium graminearum* endo-polygalacturonases in vitro and during wheat infection. Plant Pathology 58 (3), 556–564.

Torres E. (1989). Assessing the importance of *Fusarium* Head Blight in wheat
En Kohli M.M. (1989). Taller sobre Fusariosis de la espiga en America del Sur
Science. Pages 144.

Tóth B., Mesterházy Á., Horváth Z., Bartók T., Varga M., Varga J. (2005).
Genetic variability of central European isolates of the *Fusarium graminearum*
species complex. Eur J Plant Pathol. 113, 35-45.

Trail F., Xu J.R., San Miguel P., Halgren R.G., Kistler H.C. (2003). Analysis of
expressed sequence tags from Gibberella zeae (anamorph *Fusarium graminearum*).
Fungal Genet Biol. 28, 187-197.

Trail F. (2009). For Blighted Waves of Grain: *Fusarium graminearum* in the
Postgenomics Era. Plant Physiol. 149, 103-110.

Turner A.S., Lees A.K., Rezanoor H.N., Nicholson P. (1998). Refinement of
PCR-detection of *Fusarium avenaceum* and evidence from DNA marker studies for
phonetic relatedness to *Fusarium tricinctum*. Plant Pathol. 47, 274-288.

Valette-Collet O., Cimerman A., Reignault P., Levis C., Boccara M. (2003).
Disruption of *Botrytis cinerea* pectin methylesterase gene Bcpme1 reduces on
several host plants. Mol Plant Microbe Interact. 16, 360-367.

Van Ginkel M., Van Der Schaar W., Zhuping Y., Rajaram S.. (1996).
Inheritance of Resistance to Scab in Two Wheat Cultivars from Brazil and China. –
The American Phytopathological Society. Plant Disease. 80, (8),: 863-867.

Vogelsgang S., Widmer F., Jenny E., Enkerli J. (2009). Characterisation of
novel *Fusarium graminearum* microsatellite markers in different *Fusarium* species
from various countries. Eur J Plant Pathol. 123, 477-482.

Voigt Ch.A., Schafer W., Salomon S. (2005). A secreted lipase of *Fusarium*
graminearum is a virulence factor required for infection of cereals. The Plant J. 42,
364-375.

Walker S., Leath S., Hagler W.M., Murphy J.P. (2001). Variation among
isolates of *Fusarium graminearum* associated with *Fusarium* Head Blight in North
Carolina. Plant Dis. 85, 404-410.

Wanyoike W.M., Kang Z., Buchenauer H. (2002). Importance of cell wall
degrading enzymes produced by *Fusarium graminearum* during infection of wheat
head. Eur J Plant Pathol. 108, 803-810.

Ward T., Bielawski J., Kistler H., Sullivan E., O'Donnell K. (2002). Ancestral
polymorphism and adaptive evolution in the trichothecene mycotoxin gene cluster of
phytopathogenic *Fusarium*. Proc Natl Acad Sci USA. 99, 9278-9283.

Williams K.J., Dennis J.J., Smyl C., Wallwork H. (2002). The application of species-specific assay based on the polymerase chain reaction to analyze *Fusarium* crown rot of durum wheat. *Austral Plant Pathol.* 31, 119-129.

Windels C. (2000). Economical and social impacts of *Fusarium* Head Blight changing farms and rural communities in the northern great plains. *Phytopathol.* 90, 17-21.

Wood T.M. and García-Campayo V. (1990). Enzymology of cellulose degradation. *Biodegradation.* 1, 147-161.

Xu J.R., Staiger C.J., Hamer J.E. (1998). Inactivation of the mitogen activated protein kinase Mps 1 from the rice blast fungud prevent penetration of the host cell but allows activation of plant defense. *Proceedings of the National Academy of Sciense USA.* 95, 12713-12718.

Xu D.H., Juan H.F, Nohda M. and Ban T.. (2001). QTLs Mapping of Type I and Type II Resistance to FHB in Wheat. *National Fusarium Head Blight Forum. Biotechnology,* 40-42.

Xu X. (2003). Effects of environmental conditions on the development of *Fusarium* ear blight. *Eur J Plant Pathol* 109, 683-689.

Yakoby N., Beno-Moualem D., Keen N.T., Dinnor A., Pines O., Prusky D. (2001). *Colletotrichum gloeosporioides* pELB is an important virulence factor in avocado fruit-fungus interaction. *Mol Plant Microbe Interact.* 14, 988-955.

Yang Z.P, Gilbert J., Somers D.J., Fedak G., Procnier J.D. and McKenzie I.H.. (2003). Marker assisted selection of *Fusarium* head blight resistance genes in two doubled haploid populations of wheat. *Molecular Breeding* 12 pp: 309-317

Yli-Mattila T., Paavanen-Huhtala S., Bulat S.A., Alekhina I.A., Nirenberg H.I., (2002). Molecular, morphological and phylogenetic analysis of the *Fusarium avenaceum/F. arthrosporioides/F. tricinctum* species complex-a polyphasic approach. *Mycological Res.* 106, 655-669.

Yli-Mattila T., Paavanen-Huhtala S., Parikka P., Konstantinova P.,Gagkaeva T.Y. (2004). Molecular and morphological diversity of *Fusarium* species in Finland and northwestern Russia. *Eur J Plant Pathol.* 110, 573-585.

Yoder W.T. and Christianson L.M. (1998). Species-specific primers resolve members of *Fusarium* section *Fusarium*. Re-evaluation of the taxonomic status of the edible Quorn fungus. *Fungal Genet Biol.* 23, 68-80.

Yoshimura Y., Nakamichi K., Kobayashi S., Nishimura T., Ikenaga M., Sato N., Sato M., Susuki T., Takeuchi T., and Yanagisawa A. (2006) Progress in

Improving Fusarium Head Blight Resistant Wheat in Hokkaido, Japan. The Global Fusarium Initiative for International Collaboration. Session 3. International Scab Nursery Consortium. 82.

Zhang W. and Yang R. (2001). Application of engieneering strain in preparation of acetic acid with orange juice. Zhongguo Niangzao, 3, 22- 26.

Zeller K.A., Bowden R.L., Leslie J.F. (2003). Diversity of epidemic populations of *Gibberella zeae* from small quadrats in Kansas and North Dakota. *Phytopathology*. 93, 874-880.

Zietkiewicz E., Rafalski A., Labuda D. (1994). Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics*. 20, 176-183.

PRODUCCIÓN CIENTÍFICA

Los resultados obtenidos en este trabajo formaron parte de las siguientes publicaciones:

- . Kikot Gisele, Hours Roque and Alconada Teresa. (2009). Contribution of cell wall degrading enzymes to pathogenesis of *Fusarium graminearum*: A review. Journal of Basic Microbiology 49(3): 231-41.
- . Kikot Gisele, Hours Roque and Alconada Teresa. (2010). Extracellular enzymes of *Fusarium graminearum* isolates. Brazilian Archives of Biology and Technology 53(4): 779-783.
- . Alconada Magliano Teresa M. and Kikot Gisele E. (2010) "Microscopy techniques to analyse plant-pathogen interaction: *Fusarium graminearum* as pathogen of wheat", in "Microscopy: Science, Technology, Applications and Education" Eds Antonio Méndez Vilas and Jesús Díaz Alvarez. Formatex Research Center. Vo. 2 ISBN (13): 978-84-614-6190-5. pp 1108-1114.
- . Kikot Gisele, Moschini Ricardo, Consolo Fabiana, Rojo Rodrigo, Hours Roque, Salerno Graciela, Gasoni Laura, Arambarri Angélica M., and Alconada Teresa M. (2011). Occurrence of *Fusarium graminearum* from wheat in relation to climatic models in Argentina Pampas Region. Mycopathologia. 171:139-149.
- . Kikot Gisele, Ortega Leonel, Rojas Natalia, Hours Roque, Lopez Laura, Alberione Enrique, Alconada Teresa. Behavior of different wheat genotypes against a *Fusarium graminearum* isolate with high polygalacturonase activity. Enviado a Phytopathology.