



Capítulo 5

BIBLIOGRAFÍA

- Agaba M, Kemp SJ. 1994. PCR-RFLP typing of the bovine myoglobin gene. *Anim Genet* 25: 187-9.
- Alin P, Danielson UH, Mannervik B. 1985. 4-hydroxyalk-2-enals are substrates for glutathione transferase. *FEBS Lett* 179: 267-270.
- Allais S, Levéziel H, Payet-Duprat N, Hocquette JF, Lepetit J, Rousset S, Denoyelle C, Bernard-Capel C, Journaux L, Bonnot A, Renard G. 2010. The two mutations, Q204X and nt821, of the myostatin gene affect carcass and meat quality in young heterozygous bulls of French beef breeds. *J Anim Sci* 88: 446-54.
- Anderson PT, Bergen WG, Merkel RA, Enright WJ, Zinn SA, Refsal KR, Hawkins DR. 1988. The relationship between composition of gain and circulating hormones in growing beef bulls fed three dietary crude protein levels. *J Anim Sci* 66: 3059-3067.
- Andersson - Eklund L, Danell B, Rendel I. 1991. Association between blood group blood protein polymorphisms and breeding values for production traits in Swedish Red and White Dary Bulls. *Anim Genet* 24: 361-376.
- Andrade PC, Grossi DA, Paz CCP, Alencar MM, Regitano LCA and Munari DP. 2008. Association of an insulin-like growth factor 1 gene microsatellite with phenotypic variation and estimated breeding values of growth traits in Canchim cattle. *Anim Genet* 39: 480-485
- Andrés S, Silva A, Soares-Pereira AL, Martins C, Bruno-Soares AM, Murray I. 2008. The use of visible and near infrared reflectance spectroscopy to predict beef *M. longissimus thoracis et lumborum* quality attributes. *Meat Sci* 78: 217-224.
- Armstrong E, Postiglioni A, Martínez A, Rincón G, Vega-Pla JL. 2006. Microsatellite analysis of a sample of Uruguayan Creole bulls (*Bos taurus*). *Genet Mol Biol* 29: 267-272.
- Barendse W, Armitage SM, Kossarek LM, Shalom A, Kirkpatrick BW, Ryan AM, Clayton D, Li L, Neibergs HL, Zhang N, Grosse WM, Weiss J, Creighton P, McCarthy F, Ron M, Teale AJ, Fries R, McGraw RA, Moore SS, Georges M, Soller M, Womack JE, Hetzel DJS. 1994. A genetic linkage map of the bovine genome. *Nat Genet* 6: 227-35.
- Bassam BJ, Caetano-Anolles G, Gresshoff PM. 1991. Fast and sensitive silver staining of DNA in polyacrilamide gels. *Ann Biochem* 19: 680-683.

- Bauman DE. Bovine somatotropin and lactation: from basic science to commercial application. 1999. *Domest Anim Endocrinol* 17: 101–116.
- Beauchemin VR, Thomas MG, Franke DE, Silver GA. 2006. Evaluation of DNA polymorphisms involving growth hormone relative to growth and carcass characteristics in Brahman steers. *Genet Mol Res* 5: 438-447.
- Behrends JM, Mikel WB, Armstrong CL, Newman MC. 2003. Color stability of semitendinosus, semimembranosus, and biceps femoris steaks packaged in a high-oxygen modified atmosphere. *J Anim Sci* 8: 2230-8.
- Bekhit Ael-D, Geesink GH, Ilian MA, Morton JD, Bickerstaffe R. 2003. The effects of natural antioxidants on oxidative processes and mettmyoglobin reducing activity in beef patties. *Food Chem* 81: 175–187.
- Bines JA, Hart IC. 1978. Hormonal regulation of the partition of energy between milk and body tissue in adult cattle. *Proc Nutr Soc* 37: 281-7.
- Bishop MD, Kappes SM, Keele JW, Stone RT, Sunden SL, Hawkins GA, Toldo SS, Fries R, Grosz MD, Yoo J, Beattie CW. 1994. A genetic linkage map for cattle. *Genetics* 136: 619–39.
- Blott S, Kim JJ, Moisio S, Schmidt-Küntzel A, Cornet A, Berzi P, Cambisano N, Ford C, Grisart B, Johnson D, Karim L, Simon P, Snell R, Spelman R, Wong J, Vilkki J, Georges M, Farnir F, Coppieters W. 2003. Molecular Dissection of a Quantitative Trait Locus: A Phenylalanine-to-Tyrosine Substitution in the Transmembrane Domain of the Bovine Growth Hormone Receptor Is Associated With a Major Effect on Milk Yield and Composition. *Genetics* 163: 253–266.
- Boles JA, Bowman JG, Surber LM, Boss DL. 2004. Effects of barley variety fed to steers on carcass characteristics and color of meat. *J Anim Sci*: 82: 2087-91.
- Brameld JM, Butterly PJ, Dawson JM and Harper JMM. 1998. Nutritional and hormonal control of skeletal-muscle cell growth and differentiation. *Proc Nutr Soc* 57: 207-217.
- Casas E, Shackelford SD, Keele JW, Stone RT, Kappes SM, Koohmaraie M. 2000. Quantitative Trait Loci Affecting Growth and Carcass Composition of Cattle Segregating Alternate Forms of Myostatin. *J Anim Sci* 78: 560-569.

- Casas E, Keele JW, Shackelford SD, Koohmaraie M, Stone RT. 2003. Identification of quantitative trait loci for growth and carcass composition in cattle. *Anim Genet* 35: 2-6.
- Ceelen HJ. 1995. Bovine somatotropin and cow health--what are the facts? *Can Vet J* 36:25-7.
- Chan WK, Faustman C, Velasquez-Pereira J, McDowell LR, Batra TR. 1998. Effects of alpha-tocopherol on metmyoglobin formation and reduction in beef from cattle fed soybean or cottonseed meal diets. *J Anim Sci* 76: 1421-6.
- Chan WKM, Faustman C, Decker EA. 1997. Oxymyoglobin oxidation as affected by oxidation products of phosphatidylcholine liposomes. *J Food Sci* 62: 709-712.
- Chen C. 2000. Growth hormone secretagogue actions on the pituitary gland: multiple receptors for multiple ligands? *Clin Exp Pharmacol Physiol* 27: 323-9.
- Chen J, Anderson JB, DeWeese-Scott C, Fedorova ND, Geer LY, He S, Hurwitz DI, Jackson JD, Jacobs AR, Lanczycki CJ, Liebert CA, Liu C, Madej T, Marchler-Bauer A, Marchler GH, Mazumder R, Nikolskaya AN, Rao BS, Panchenko AR, Shoemaker BA, Simonyan V, Song JS, Thiessen PA, Vasudevan S, Wang Y, Yamashita RA, Yin JJ, Bryant SH. 2003. MMDB: Entrez's 3D-structure database. *Nucl Ac Res* 31: 474-477.
- Chikuni K, Fukumoto Y, Tanabe R, Muroya S, Ozawa S. 1997. A simple method for genotyping the bovine growth hormone gene. *Anim Genet* 28: 230–232.
- Chillard Y. 1989. Long-term effects of recombinant bovine somatotropin (rBST) on dairy cow performances: a review. En: *Use of Somatotropin in Livestock Production*. Editores: Sejrsen K, Vestergaard M, Neimann-Sorensen A. Editorial: Elsevier Applied Science (New York) 61–87.
- CIE (Commission International de l'Eclairage) 1986. *Colorimetry* 2 ed. Vienna. s.p.
- Cogan JD, Phillips JA. 2001. III Inherited Defects in Growth Hormone Synthesis and Action (Capítulo 162). En: *The Metabolic & Molecular Bases of Inherited Disease* 8TH ED. Editor: Scriver CR. Editorial: McGraw-Hill.

- Cowan CM, Dentine MR, Ax RL, Schuler LA. 1989. Restriction fragment length polymorphisms associated with growth hormone and prolactin genes in Holstein bulls: evidence for a novel growth hormone allele. *Anim Genet* 20:157-65.
- Curi RA, Oliveira HN, Silveira AC, Lopes CR. 2005. Effects of polymorphic microsatellites in the regulatory region of IGF1 and GHR on growth and carcass traits in beef cattle. *Anim Genet* 36: 58-62.
- Curi RA, Palmieri DA, Sugisawa L, Oliveira HN, Silveira AC, Lopes CR. 2006. Growth and carcass traits associated with GH1/Alu I and POU1F1/Hinf I gene polymorphisms in Zebu and crossbred beef cattle. *Genet Mol Biol* 29: 56-61.
- De Donato M, Gallagher Jr DS, Lehn C, Gill C, Taylor JF. 2003. Molecular cytogenetic assignment of genes to bovine chromosome 5. *Genet Mol Res* 2: 260-70.
- Dekkers JC. 2004. Commercial application of marker- and gene-assisted selection in livestock: strategies and lessons. *J Anim Sci* 82: E313-328.
- Di Stasio L, Destefanis G, Brugia Paglia A, Albera A, Rolando A. 2005. Polymorphism of the GHR gene in cattle and relationships with meat production and quality. *Anim Genet* 36: 138–140.
- Di Stasio L, Sartore S, Albera A. 2002. Lack of association of GH1 and POU1F1 gene variants with meat production traits in Piemontese cattle. *Anim Genet* 33: 61-64.
- Dolf G, Glowatzki ML, Gaillard C. 1991. Searching for genetic markers for hereditary diseases in cattle by means of DNA fingerprinting. *Electrophoresis* 12:109-12.
- Dunne PG, O'Mara FP, Monahan FJ, Moloney AP. 2006. Changes in colour characteristics and pigmentation of subcutaneous adipose tissue and M. longissimus dorsi of heifers fed grass, grass silage or concentrate-based diets. *Meat Sci* 74: 231–241.
- Erlich HA. 1989. PCR technology: principles and applications for DNA amplification. Editorial: Stockton Press (Nueva York) 246 pp.
- Etherton TD. 2004. Somatotropic function: the somatomedin hypothesis revisited. *J Anim Sci* 82: E239-244.
- Etherton TD, Bauman DE. 1998. Biology of somatotropin in growth and lactation of domestic animals. *Physiol Rev* 78: 745-61.
- Faustman C, Cassens RG. 1990. The biochemical basis for discoloration in fresh meat: a review. *J Muscle Foods* 1: 217– 243.

Faustman C, Cassens RG. 1991. The effect of cattle breed and muscle type on discoloration and various biochemical parameters in fresh beef. *J Anim Sci* 69:184-93.

Faustman C, Chan WK, Schaefer DM, Havens A. 1998. Beef Color Update: The Role for Vitamin E. *J Anim Sci* 76:1019-26.

Florini JR, Ewton DZ, Roof SL. 1991. Insulin-like growth factor-I stimulates terminal myogenic differentiation by induction of myogenin gene expression. *Mol Endocrinol* 5: 718-724.

Fontanesi L, Scotti E, Tazzoli M, Beretti F, Dall'Olio S, Davoli R, Russo V. 2007. Investigation of allele frequencies of the growth hormone receptor (GHR) F279Y mutation in dairy and dual purpose cattle breeds. *Ital J Anim Sci* 6: 415-420.

Ge W, Davis ME, Hines HC, Irvin KM, Simmen RC. 2001. Association of a genetic marker with blood serum insulin-like growth factor-I concentration and growth traits in Angus cattle. *J Anim Sci* 79: 1757–1762.

Ge W, Davis ME, Hines HC, Irvin KM, Simmen RCM. 2003. Association of single nucleotide polymorphisms in the growth hormone and growth hormone receptor genes with blood serum insulin-like growth factor I concentration and growth traits in Angus cattle. *J Anim Sci* 81: 641–648.

Gelfi C, De Palma S, Ripamonti M, Eberini I, Wait R, Bajracharya A, Marconi C, Schneider A, Hoppeler H, Cerretelli P. 2004. New aspects of altitude adaptation in Tibetans: A proteomic approach. *FASEB J* 18: 612–614.

Georges M, Larthrop M, Dietz AB, Lefort A, Libert F, Mishra A, Nielsen D, Sargeant LS, Steele MR, Zhao X, Leipold H, Womack E. 1993. Microsatellite mapping of the gene causing weaver disease in cattle will allow the study of an associated QTL. *Proc Natl Acad Sci USA* 90: 1058-1062.

Giovambattista G, Ripoli MV, Peral García P, Bouzat, JL. 2001. Indigenous domestic breeds as reservoirs of genetic diversity: The Argentine Creole Cattle. *Anim Genet* 32: 240-247.

Gordon DF, Quick DP, Erwin CR, Donelson JE, Maurer RA. 1983. Nucleotide sequence of the bovine growth hormone chromosomal gene. *Mol Cell Endocrinol* 33: 81-95.

Graml R, Olbrich-Bludau A, Schwab M, Schallenberger E, Schams D, Pirchner F. 1994. Relationship between plasma hormone and metabolite levels and breeding values of bulls. *J Anim Breed Genet* 112: 313–326.

- Grochowska R, Sørensen P, Zwierzchowski L, Snochowski M, Løvendahl P. 2001. Genetic variation in stimulated GH release and in IGF-I of young dairy cattle and their associations with the leucine/valine polymorphism in the GH gene. *J Anim Sci* 79: 450–476.
- Hale CS, Herring WO, Shibuya H, Lucy MC, Lubahn DB, Keisler DH, Johnson GS. 2000. Decreased growth in Angus steers with a short TG-microsatellite allele in the P1 promoter of growth hormone receptor gene. *J Anim Sci* 78: 2099–104.
- Hallerman EM, Nave A, Kashi Y, Holzer Z, Soller, Beckmann JS. 1987. Restriction fragment length polymorphisms in dairy and beef cattle at the growth hormone and prolactin loci. *Anim Genet* 18: 213–22.
- Heaton MP, Harhay GP, Bennett GL, Stone RT, Grosse WM, Casas E, Keele JW, Smith TP, Chitko-McKown CG, Laegreid WW. 2002. Selection and use of SNP markers for animal identification and paternity analysis in U.S. beef cattle. *Mamm Genome* 13:272-81.
- Hediger R, Johnson SE, Barendse W, Drinkwater RD, Moore SS, Hetzel J. 1990. Assignment of the growth hormone gene locus to 19q26-qter in cattle and 11q25-qter in sheep by in situ hybridization. *Genomics* 8: 171–4.
- Henderson CR, Quaas RL. 1976. Multiple trait evaluation using relatives' records. *J Anim Sci* 43: 1188-1197.
- Høj S, Fredholm M, Larsen NJ, Nielsen VH. 1993. Growth hormone gene polymorphism associated with selection for milk fat production in lines of cattle. *Anim Genet* 24: 91–6.
- Hunter RS, Harold RW. 1987. The measurement of appearance (2nd ed.). Editorial: Wiley (New York).
- Ihara N, Takasuga A, Mizoshita K, Takeda H, Sugimoto M, Mizoguchi Y, Hirano T, Itoh T, Watanabe T, Reed KM, Snelling WM, Kappes SM, Beattie CW, Bennett GL, Sugimoto Y. 2004. A comprehensive genetic map of the cattle genome based on 3802 microsatellites. *Genome Res* 14: 1987-98.
- Ishikawa T, Esterbauer H, Sies H. 1986. Role of cardiac glutathione transferase and of the glutathione S-conjugate export system in biotransformation of 4-hydroxynonenal in the heart. *J Biol Chem* 261: 1576-1581.

Katoh K, Kouno S, Okazaki A, Suzuki K, Obara Y. 2008. Interaction of GH polymorphism with body weight and endocrine functions in Japanese black calves. Domestic Anim Endoc 34: 25–30

Kawano K, Uda K, Otsuki R, Suzuki T. 2004. Preparation of artificial 2-, 3-, 4- and 8-domain myoglobins and comparison of their autoxidation rates. FEBS Letters 574: 203–207.

Kemenes PA, Correia de Almeida Regitano L, Jordão de Magalhães Rosa A, Packer IU, Razook AG, Andrade de Figueiredo L, Silva NA, Etchegaray MAL, Lehmann Coutinho L. 1999. k-casein, b-lactoglobulin and growth hormone allele frequencies and genetic distances in Nelore, Gyr, Guzerá, Caracu, Charolais, Canchim and Santa Gertrudis cattle. Genet Mol Biol 22, 4, 539-541.

Kirkpatrick M, Heckman N. 1989. A quantitative genetic model for growth, shape, reaction norms, and other infinitedimensional characters. J Math Biol 27: 429-450.

Kitagawa H, Kitoh K, Ito T, Ohba Y, Nishii N, Katoh K, Obara Y, Motoi Y, Sasaki Y. 2001. Serum Growth Hormone and Insulin-Like Growth Factor-1 Concentrations in Japanese Black Cattle with Growth Retardation. J Vet Med Sci 63: 167-70.

Koshkohi AE, Pitchford WS, Kruk ZA, Morris CA, Cullen NG, Crawford AM, Bottema CDK. 2005. QTL for meat colour and pH in *Bos Taurus* cattle. Proc. Assoc. Advmt. Anim. Breed. Genet. 16: 111-114

Kossarek LM, Finlay O, Grosse WM, Su X, McGraw R. 1994. Five bovine dinucleotide repeat polymorphisms: RM026, RM029, RM032, RM033 and RM038. Anim Genet 25: 296-97.

Kuimelis RG, Livak KJ, Mullah B, Andrus A. 1997. Structural analogues of TaqMan probes for real-time quantitative PCR. Nucleic Acids Symp Ser 37:255-6.

Ledward DA. 1985. Post-slaughter influences on the formation of metmyoglobin in beef muscles. Meat Sci 15: 149.

Li C, Basarab J, Snelling WM, Benkel B, Murdoch B, Hansen C, Moore SS. 2004. Assessment of Positional Candidate Genes myf5 and igf1 for Growth on Bovine Chromosome 5 in Commercial Lines of *Bos Taurus*. J Anim Sci 82: 1-7.

Li C, Basarab J, Snelling WM, Benkel B, Murdoch B, Moore SS. 2002. The identification of common haplotypes on bovine chromosome 5 within commercial lines of Bos taurus and their associations with growth traits. *J Anim Sci* 80: 1187-94.

Li MH, Adamowicz T, Switonski M, Ammosov I, Ivanova Z, Kiselyova T, Popov R, Kantanen J. 2006. Analysis of population differentiation in North Eurasian cattle (Bos taurus) using single nucleotide polymorphisms in three genes associated with production traits. *Anim Genet* 37: 390–392.

Li N, Stephens M. 2003. Modelling linkage disequilibrium, and identifying recombination hotspots using snp data. *Genetics* 165: 2213-2233.

Lirón JP. 2004. Tesis de Doctorado: "Estudio de la Variabilidad Genética de Razas Bovinas Criollas, Europeas e Indicas de Argentina y Bolivia. Su aplicación en la Identificación Individual, Paternidades y Trazabilidad". Fac Cs Veterinarias, UNLP, La Plata – Argentina.

Lo Bello M, Nuccetelli M, Caccuri AM, Stella L, Parker MW, Rossjohn J, McKinstry WJ, Mozzi AF, Federici G, Polizio F, Pedersen JZ, Ricci G. 2001. Human glutathione transferase P1-1 and nitric oxide carriers; a new role for an old enzyme. *J Biol Chem* 276: 42138–42145.

Machado MBB, Alencar MM, Pereira AP, Oliveira HN. 2003. QTL affecting body weight in a candidate region of cattle chromosome 5. *Genet Mol Biol* 3: 259-265.

Mancini RA, Hunt MC. 2005. Current research in meat color. *Meat Sci* 71: 100–121.

Mangiacapra FJ, Roof SL, Ewton DZ, Florini JR. 1992. Paradoxical decrease in Myf-5 messenger RNA levels during induction of myogenic differentiation by insulin-like growth factors. *Mol Endocrinol* 6: 2038-2044.

McClure MC, Morsci NS, Schnabel RD, Kim JW, Yao P, Rolf MM, McKay SD, Gregg SJ, Chapple RH, Northcutt SL, Taylor JF. 2010. A genome scan for quantitative trait loci influencing carcass, post-natal growth and reproductive traits in commercial Angus cattle. *Anim Genet* (On line).

Melucci LM, Birchmeier AN, Cappa EP, Cantet RJC. 2009. Bayesian analysis of selection for higher weaning weight while maintaining birth weight in beef cattle. *J Anim Sci* 87: 3089-96.

Melucci LM, Miquel MC, Molinuevo HA. 1983. Índices de selección para crecimiento en bovinos para carne. *Producción Animal* 10:417-426.

Melucci LM. 1995. Estimación de un índice genético restricto para crecimiento predestete en bovinos para carne. Resúmenes 1º Jornada Argentino-Chilena de Genética. XXVI Congreso Argentino de Genética y XXVIII Reunión Anual de la Sociedad de Genética de Chile. Bariloche (Argentina).

Meyer K. 1997. Estimates of genetic parameters for weaning weight of beef cattle accounting for direct-maternal environmental covariances. *Livest Prod Sci* 52: 187- 199.

Miquel MC, Villarreal E, Mezzadra C, Melucci L, Soria L, Corva P, Schor A. 2009. The association of CAPN1 316 marker genotypes with growth and meat quality traits of steers finished on pasture. *Genet Mol Biol* 32:491-496.

Mizoshita K, Takano A, Watanabe T, Takasuga A, Sugimoto Y. 2005. Identification of a 1.1-Mb region for a carcass weight QTL on bovine Chromosome 14. *Mamm Genome* 16: 532-7.

Montoya AE, Cerón-Muñoz MF, Moreno MA, Martínez E, Corrales JD, Tirado JF, Calvo SJ. 2010. Genetic characterization of the Hartón del Valle, Angus, Brangus, Holstein, and Senepol cattle breeds in Colombia, using ten microsatellite markers. *Rev Colomb Cienc Pecu* 23: 283-291.

Moody DE, Pomp D, Barendse W, Womack JE. 1995. Short communication: assignment of the growth hormone receptor gene to bovine chromosome 20 using linkage analysis and somatic cell mapping. *Anim Genet* 26:341– 4.

Moody DE, Pomp D, Newman S, MacNeil MD. 1996. Characterization of DNA polymorphisms in three populations of Hereford cattle and their associations with growth and maternal EPD in line 1 Herefords. *J Anim Sci* 74: 1784-1793.

Mrode RA. 2005. Linear Models for the Prediction of Animal Breeding Values. 2nd edn. Editorial: CABI Publishing Company (Cambridge).

Navarro I, Leibush B, Moon TW, Plisetskaya EM, Banos N, Mendez E, Planas JV, Gutierrez J. 1999. Insulin, insulin-like growth factor- I (IGF-I), and glucagon: the evolution of their receptors. *Comp Biochem Physiol B* 122: 137–153.

Nelson DL, Cox MM. 2000. Glycolysis and the catabolism of hexoses. En: Lehninger Principle of Biochemistry 3a edición. Editorial: Worth Publishers (New York): 560.

- Novoa MA, Usaquén W. 2010. Population genetic analysis of the Brahman cattle (*Bos indicus*) in Colombia with microsatellite markers. *J Anim Breed Genet* 127: 161–168.
- O'Keefe M, Hood DE. 1982. Biochemical factors influencing metmyoglobin formation in beef from muscles of differing colour stability. *Meat Sci* 7: 209–228.
- Orita M, Suzuki Y, Sekiya K, Hayashi. 1989b. A rapid and sensitive detection of point mutations and genetic polymorphism using polymorphism chain reaction. *Genomics* 5: 874–879.
- Orita MH, Wahana I, Kanazawa H, Sekna T. 1989a. Detection of polymorphism of human DNA by gel electrophoresis as single strand conformation polymorphism. *Proc Natl Acad Sci USA* 86: 2766–2770.
- Otto G, Roehe R, Looft H, Thoelking L, Knap PW, Rothschild MF, Plastow GS, Kalm E. 2007. Associations of DNA markers with meat quality traits in pigs with emphasis on drip loss. *Meat Sci* 75: 185–195.
- Ouali A, Herrera-Mendez CH, Coulis G, Becila S, Boudjellal A, Aubry L, Sentandreu MA. 2006. Revisiting the conversion of muscle into meat and the underlying mechanisms. *Meat Sci* 74: 44–58.
- Papaleo Mazzucco J, Melucci LM, Villarreal EL, Mezzadra CA, Soria L, Corva P, Motter MM, Schor A, Miquel MC. 2010. Effect of ageing and μ -calpain markers on meat quality from Brangus steers finished on pasture. *Meat Sci* 86: 878–882.
- Pastore A, Federici G, Bertini E, Piemonte F. 2003. Analysis of glutathione: implication in redox and detoxification. *Clin Chim Acta* 333: 19–39.
- Pereira AP, Alencar MM, Oliveira HN, Regitano LCA. 2005. Association of GH and IGF-I polymorphism with growth traits in a synthetic beef cattle breed. *Genet Mol Biol* 28: 230–236.
- Priolo A, Micol D, Agabriel J. 2001. Effects of grass feeding systems on ruminant meat colour and flavour. A review. *Anim Res* 50: 185–200.
- Radko A, Zyga A, Zabek T, Slota E. 2005. Genetic variability among Polish Red, Hereford and Holstein-Friesian cattle raised in Poland based on analysis of microsatellite DNA sequences. *J Appl Genet* 46: 89–91.
- Reardon W, Mullen AM, Sweeney T, Hamill RM. 2010. Association of polymorphisms in candidate genes with colour, water-holding capacity, and

composition traits in bovine *M. longissimus* and *M. semimembranosus*. Meat Sci 86: 270-5.

Regitano LCA, Azevedo JL, Vencovsky R, Packer IU, Barbosa PF, Rosa AJM, Silva NA, Etchegaray MAL, Coutinho LL. 1999. Selection for breed-specific growth hormone and IGF-I alleles in synthetic beef cattle cross, Canchim. Genet Mol Biol 22: 531-537.

Reis C, Navas D, Pereira M, Cravador A. 2001. Growth hormone AluI polymorphism analysis in eight portuguese bovine breeds. Arch Zoot 50: 189-190.

Renaville R, Hammadi M, Portetelle D. 2002. Role of the somatotropic axis in the mammalian metabolism. Domest Anim Endocrinol 23: 351-60.

Renerre M, Mazuel JP. 1985. Relations entre méthodes de mesures instrumentales et sensorielles de la couleur de la viande. Sci Aliments 5: 541-557.

Ripoli MV, Rogberg-Muñoz A, Francisco EI, Villegas-Castagnasso EE, Lirón JP, Peral-García P, Giovambattista G. 2010. History and selection imprinting on genetic relationships among bovine breeds analyzed through five genes related with marbling. Res Vete Sci, *in press*.

Rogberg Muñoz A, Ripoli MV, Villegas-Castagnasso EE, Kienast ME, Furnus CC, Peral-García P, Giovambattista G. 2006. Genetic Variability Study in 19 Bovine Breeds of Five Candidate Genes Related with Marbling (DGAT1, TG, GH, LP AND SCD). Proceedings of the 30th International Conference on Animal Genetics ISAG 2006 (ISBN 85-85584-02-5).

Ronaghi M, Uhlen M, Nyrén P. 1998. A sequencing method based on real-time pyrophosphate. Science 281: 363-5.

Ronaghi M. 2001. Pyrosequencing sheds light on DNA sequencing. Genome Res 11: 3-11.

Rousset F. 2008. GENEPOL '007: a complete re-implementation of the GENEPOL software for Windows and Linux. Mol Ecol Res 8: 103-6.

Rudnicki MA, Schnegelsberg PNJ, Stead RH, Braun T, Arnold HH, Jaenisch R. 1993. MyoD or Myf-5 is required for the formation of skeletal muscle. Cell 75: 1351-1359.

Schulman NF, Sahana G, Lund MS, Viitala SM, Vilkki JH. 2008. Quantitative trait loci for fertility traits in Finnish Ayrshire cattle. Genet Sel Evol 40: 195-214.

- Scollan N, Hocquette JF, Nuernberg K, Dannenberger D, Richardson I, Moloney A. 2006. Innovations in beef production systems that enhance the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Sci* 74: 17-33. International Congress 52nd ICoMST, Dublin, Ireland.
- Seyfert M, Mancini RA, Hunt MC, Tang J, Faustman C, Garcia M. 2006. Color stability, reducing activity, and cytochrome c oxidase activity of five bovine muscles. *J Agric Food Chem* 54: 8919-25.
- Sherman EL, Nkrumah JD, Murdoch BM, Li C, Wang Z, Fu A, Moore SS. 2007. Polymorphisms and haplotypes in the bovine NPY, GHR, GHRL, IGF2, UCP2, and UCP3 genes and their associations with measures of growth, performance, feed efficiency and carcass merit in beef cattle. *J Anim Sci* (On line).
- Sherman GB, Kachman SD, Hungerford LL, Rupp GP, Fox CP, Brown MD, Feuz BM, Holm TR. 2004. Impact of candidate sire number and sire relatedness on DNA polymorphism-based measures of exclusion probability and probability of unambiguous parentage. *Anim Genet* 35: 220–226.
- Smith GC, Belk KE, Sofos JN, Tatum JD, Williams SN. 2000. Economic implications of improved color stability in beef. En: Antioxidants in muscle foods: Nutritional strategies to improve quality. Editores: Decker EA, Faustman C, Lopez-Bote CJ. Editorial: Wiley Interscience (New York): 397–426.
- Sobrino B, Brion M, Carracedo A. 2005. SNPs in forensic genetics: a review on SNP typing methodologies. *Forensic Sci Int* 154: 181 – 194.
- Steffen P, Eggen A, Dietz Ab, Womack Je, Stranzinger G, Fries R. 1993. Isolation and mapping of polymorphic microsatellites in cattle. *Anim Genet* 24: 1221-1224.
- Stevanović J, Stanimirović Z, Dimitrijević V, Stojić V, Fratrić N, Lazarević M. 2009. Microsatellite DNA polymorphism and its usefulness for pedigree verification in Simmental cattle from Serbia. *Acta Vet (Beogr)* 59: 621-631.
- Stone RT, Casas E, Smith TP, Keele JW, Harhay G, Bennett GL, Koohmaraie M, Wheeler TL, Shackelford SD, Snelling WM. 2005. Identification of genetic markers for fat deposition and meat tenderness on bovine chromosome 5: development of a low-density single nucleotide polymorphism map. *J Anim Sci* 83: 2280-8.

- Suman SP, Faustman C, Stamer SL, Liebler DC. 2007. Proteomics of lipid oxidation-induced oxidation of porcine and bovine oxymyoglobins. *Proteomics* 7: 628–640.
- Suzuki R, Kemp SJ, Teale AJ. 1993. Polymerase chain reaction analysis of mitochondrial DNA polymorphism in N'Dama and Zebu cattle. *Anim Genet* 24: 339–343.
- Tang J, Faustman C, Hoagland TA. 2004. Krzywicki Revisited: Equations for Spectrophotometric Determination of Myoglobin Redox Forms in Aqueous Meat Extracts *J Food Sci* 69:717–720.
- Tang J, Faustman C, Lee S, Hoagland TA. 2003. Effect of Glutathione on Oxymyoglobin Oxidation. *J Agric Food Chem* 51: 1691-1695.
- Taylor J, Coutinho L, Herring K, GallagherJr D, Brenneman R, Burney N, Sanders J, Turner J, Smith S, Miller R, Savell J, Davis S. 1998. Candidate gene analysis of GH1 for effects on growth and carcass composition of cattle. *Anim Genet* 29: 194-201.
- Thallman RM, Bennett GL, Keele JW, Kappes SM. 2001a. Efficient computation of genotype probabilities for loci with many alleles: I. Allelic peeling. *J Anim Sci* 79: 26–33.
- Thallman RM, Bennett GL, Keele JW, Kappes SM. 2001b. Efficient computation of genotype probabilities for loci with many alleles: II. Iterative method for large, complex pedigrees. *J Anim Sci* 79: 34–44.
- Theilmann JL, Skow LC, Baker JF, Womack JE. 1989. Restriction fragment length polymorphisms for growth hormone, prolactine, osteonectin, cristallin, fibronectin and 21-steroid hydroxilase in cattle. *Anim Genet* 20: 257 - 266.
- Thomas MG, Enns RM, Shirley KL, Garcia MD, Garrett AJ, Silver GA. 2007. Associations of DNA polymorphisms in growth hormone and its transcriptional regulators with growth and carcass traits in two populations of Brangus bulls. *Genet Mol Res* 6: 222-37.
- Tollefson SE, Lajara R, McCusker RH, Clemmons DR, Rotwein P. 1989. Insulin-like Growth Factors (IGF) in Muscle Development. *J Biol Chem* 264: 13810-13817.
- Unanian MM, Chaves Barreto C, Ribeiro de Freitas A, Torres Cordeiro CM, Josahkian LA. 2000. Associação do Polimorfismo do Gene do Hormônio de

Crescimento com a Caraterística Peso em Bovinos da Raça Nelore. Rev Bras Zootec 29: 1380-1386.

Van de Goor LHP, Koskinen MT, van Haeringen WA. 2009. Population studies of 16 bovine STR loci for forensic purposes. Int J Legal Med (On line).

Varvio SL, Iso-Touru T, Kantanen J, Viitala S, Tapiola I, Mäki-Tanila A, Zerabruk M, Vilkki J. 2008. Molecular anatomy of the cytoplasmic domain of bovine growth hormone receptor, a quantitative trait locus. Proc R Soc B 275: 1525-1534.

Vasconcellos LP, Tambasco-Talhari D, Pereira AP, Coutinho LL, Regitano LC. 2003. Genetic characterization of Aberdeen Angus cattle using molecular markers. Genet Mol Biol 26: 133-137.

Vestergaard M, Oksberg N, Henckel P. 2000. Influence of feeding intensity, grazing and finishing feeding on muscle fibre characteristics and meat colour of semitendinosus, longissimus dorsi and supraspinatus muscles of young bulls. Meat Sci 54: 177-185.

Viitala S, Szyda J, Blott S, Schulman N, Lidauer M, Mäki-Tanila A, Georges M, Vilkki J. 2006. The role of the bovine growth hormone receptor and prolactin receptor genes in milk, fat and protein production in Finnish Ayrshire dairy cattle. Genetics 173: 2151-64.

Villegas-Castagnasso EE, Kienast M, Ripoli MV, Prando AJ, Francisco EI, Rogberg Muñoz A, Sorarrain N, Baldo A, Peral-García P, Giovambattista G. 2006. Comparación de la Variabilidad Genética de los Genes DGTA1, TG, LP, SCD y GH en Diferentes Tipos de Bovinos de la Raza Aberdeen Angus de Argentina. XXXV Congreso Argentino de Genética, San Luís. BAG XVIII (supplement): 161.

Walshe BE, Sheehan EM, Delahunty CM, Morrissey PA, Kerry JP. 2006. Composition, sensory and shelf life stability analyses of Longissimus dorsi muscle from steers reared under organic and conventional production systems. Meat Sci 73: 319-325.

Wang L, Marincola FM. 2003. Applying PyrosequencingTM to HLA-Typing. Vendor Forum (American Society for Histocompatibility and Immunogenetics). 27: 16-18. Pagina visitada el 15 de mayo de 2010. http://www.ashihla.org/docs/newsletter/ASHI_Quarterly/27_1_2003/Pyrosequencing-Vendor_Web.pdf

Wang YD, Wood WI. 1995. Amino acids of the human growth hormone receptor that are required for proliferation and Jak-Stat signaling. Mol Endocrinol 9: 303–11.

Waters SM, McCabe MS, Howard DJ, Giblin L, Magee DA, Machugh DE, Berry DP. 2010. Associations between newly discovered polymorphisms in the Bos tauri growth hormone receptor gene and performance traits in Holstein-Friesian dairy cattle. Anim Genet (On line).

Weller JI. 2007. Current and future developments in patents for quantitative trait loci in dairy cattle. Recent Pat DNA Gene Seq 1: 69-76.

White SN, Casas E, Allan MF, Keele JW, Snelling WM, Wheeler TL, Shackelford SD, Koohmaraie M, Smith TP. 2007. Evaluation in beef cattle of six deoxyribonucleic acid markers developed for dairy traits reveals an osteopontin polymorphism associated with postweaning growth. J Anim Sci 85: 1-10.

Yan Z, Serrano AL, Schiaffino S, Bassel-Duby S, Williams RS. 2001. Regulatory Elements Governing Transcription in Specialized Myofiber Subtypes. J Biol Chem 276: 17361–17366.

Yao J, Aggrey SE, Zadworny D, Hayes JF, Kühnlein U. 1996. Sequence variations in the bovine growth hormone gene characterized by single-strand conformation polymorphism (SSCP) analysis and their association with milk production traits in Holsteins. Genetics 144: 1809-1816.

Yuan ZR, Xu SZ. 2010. Novel SNPs of the bovine CACNA2D1 gene and their association with carcass and meat quality traits. Mol Biol Rep (On line).

