

Semblanzas Ictiológicas
Mariano González Castro



Hugo L. López
y
Justina Ponte Gómez

Indizada en la base de datos ASEA C.S.A.
2013

Semblanzas Ictiológicas

Mariano González Castro



Hugo L. López y Justina Ponte Gómez

ProBiota
División Zoología Vertebrados
Museo de La Plata
FCNyM, UNLP

Octubre, 2013

Imagen de Tapa

Amplificando genes en el Laboratorio de Referencia Barcode, Museo Argentino de Ciencias Naturales Bernardino Rivadavia, Buenos Aires, 2011

El tiempo acaso no exista. Es posible que no pase y sólo pasemos nosotros.

Tulio Carella

Cinco minutos bastan para soñar toda una vida, así de relativo es el tiempo.

Mario Benedetti

Semblanzas Ictiológicas

A través de esta serie intentaremos conocer diferentes facetas personales de los integrantes de nuestra “comunidad”.

El cuestionario, además de su principal objetivo, con sus respuestas quizás nos ayude a encontrar entre nosotros puntos en común que vayan más allá de nuestros temas de trabajo y sea un aporte a futuros estudios históricos.

Esperamos que esta iniciativa pueda ser otro nexo entre los ictiólogos de la región, ya que consideramos que el resultado general trascendería nuestras fronteras.

Hugo L. López

Nombre y apellido completos: Mariano González Castro

Lugar de nacimiento: Mar del Plata, Buenos Aires.

Lugar, provincia y país de residencia: Mar del Plata, Buenos Aires. Argentina.

Título máximo, Facultad y Universidad: Doctor en Ciencias Biológicas, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata (UNMdP).

Posición laboral: Investigador Adjunto CONICET.

Lugar de trabajo: Laboratorio de Biotaxonomía Morfológica y Molecular de Peces (BIMOPE), Instituto de Investigaciones Marinas y Costeras, IIMyC-CONICET, UNMdP.

Especialidad o línea de trabajo: Biología reproductiva; morfometría geométrica y taxonomía (morfológica y molecular) de peces.

Correo electrónico: gocastro@mdp.edu.ar

Cuestionario

- **Un libro:** *El mundo perdido* de Sir. Arthur Conan Doyle.
- **Una película:** *La sociedad de los poetas muertos*.
- **Un CD:** *Cuando los ángeles lloran*; Intérprete: Maná.
- **Un artista:** Sting.
- **Un deporte:** padel.
- **Un color:** verde.
- **Una comida:** corvina negra a la parrilla.
- **Un animal:** perro.
- **Una palabra:** voluntad.
- **Un número:** 26.
- **Una imagen:** el amanecer en el mar.
- **Un lugar:** La Habana, Cuba.
- **Una estación del año:** otoño.
- **Un nombre:** Mariano.
- **Un hombre:** Martin Luther King.
- **Una mujer:** María teresa de Calcuta.
- **Un personaje de ficción:** Luke Skywalker de Star Wars.
- **Un superhéroe:** Batman.



Mariano González Castro, centro, Jefe Científico de la Campaña Costera Litoral Bonaerense, Buque Oceanográfico Puerto Deseado, CONICET, Santa Cruz, 2012

Journal of the Marine Biological Association of the United Kingdom, 2009, 89(3), 593–604. © 2009 Marine Biological Association of the United Kingdom
doi:10.1017/S0025315409005002 Printed in the United Kingdom

Fish composition in a south-western Atlantic temperate coastal lagoon: spatial – temporal variation and relationships with environmental variables

M. GONZÁLEZ CASTRO^{1,2}, J.M. DÍAZ DE ASTARLOA^{1,3,4}, M.B. COUSSEAU¹, D.E. FIGUEROA¹, S.M. DELPIANI^{1,5}, D.O. BRUNO¹, J.M. GUZZONI¹, G.E. BLASINA^{1,3} AND M.Y. DELI ANTONI^{1,3}

¹Laboratorio de Ictiología, Departamento de Ciencias Marinas, Universidad Nacional de Mar del Plata (UNMDP), Argentina, ²Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC), ³Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), ⁴Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), ⁵Museo del Mar, Mar del Plata, Argentina

*Mar Chiquita, an irregularly shaped brackish-water coastal lagoon, is located in the Buenos Aires province of Argentina and considered since 1996 by the Coordination Council of the Man and Biosphere Program of UNESCO as a World Reserve of Biosphere. The present paper aims to study both the spatial and temporal variation of fish composition in this coastal lagoon and the influence of some environmental variables on the relative abundance of the main fish species. Monthly sampling surveys over a two-year period in three different areas were conducted, using a beach-seine net and three monofilament-gill nets with different mesh size. Twenty-eight species belonging to four bio-ecological categories were identified, five of them are new records for Mar Chiquita fish community. The correspondence analysis showed strong relationships between high salinity range and the abundance of *Brevoortia aurea*, *Gynoscion gusatucupa* and *Pomatomus saltatrix*. Conversely, low salinity range corresponded to high abundance of *Mugil platanus* and *Odontesthes argentinensis*. High temperatures were corresponded with abundance of *Micropogonias furnieri* and *Brevoortia aurea*. In contrast, high abundance of both *Odontesthes argentinensis* and *Oligosarcus jenynsii* were corresponded to low temperatures. *Brevoortia aurea*, *Mugil platanus* and *Odontesthes argentinensis* were the most abundant species, representing more than 80% of the total capture. The group of estuarine-nondependent-marine fish presented the highest species richness. Estuarine-dependent-marine species presented for both juveniles and adults specimens the highest abundance values.*

Keywords: fish composition, species diversity, coastal lagoons, salinity gradients, environmental factors, multivariate analysis, South America, Argentina, Buenos Aires, Mar Chiquita

Submitted 24 July 2008; accepted 21 September 2008; first published online 24 March 2009

INTRODUCTION

Coastal lagoons are shallow estuarine environments where salt and freshwater interact, usually oriented parallel to the coast, separated from the ocean by some type of sedimentary barrier, and connected to the open sea by one or more restricted inlets (Isla, 1995). Because of permanent exposure to the effects of the tides, seasonal amount of freshwater entering the estuarine environment and oceanic storms that may push in more salt water, coastal lagoons are extremely rigorous ecosystems, with highly fluctuating physical (temperature, salinity, turbidity and dissolved oxygen) and biological (recruitment, predation and competition) factors (Whitfield, 1999; Hemingway & Elliott, 2002).

Coastal lagoons are considered highly productive, even more so than the open sea (Day *et al.*, 1981). The functional roles of these habitats to fish have been extensively

investigated worldwide, in temperate, subtropical and tropical areas (e.g. Whitfield, 1999; Elliott & Hemingway, 2002; James *et al.*, 2007), with a particular focus on their nursery function. These habitats also have long been recognized as important feeding areas for fish, and the role of coastal lagoons to commercial fish as a spawning area is relatively well understood (Elliott & Hemingway, 2002). This importance is related to the fact that coastal lagoon fish communities are seasonally dominated by high densities of juvenile stages of many marine species (Day *et al.*, 1981), potentially enhancing their early growth and survival (Elliott & Hemingway, 2002).

Fish fauna occurring in these habitats is a mixture of tolerant species from both marine and freshwater environments, species migrating from one environment to another, and a small number of resident species (Moyle & Cech, 2004). Therefore, these ecosystems are known as sites of low diversity but high abundance of a few dominant species (Veiga *et al.*, 2006).

The abundance of individual species and the composition, abundance, and diversity of the total fish fauna have been widely studied in both tropical (Araújo & Costa de Azevedo, 2001; Kuo *et al.*, 2001) and temperate lagoons (Vieira & Musick, 1994; Whitfield, 1999; Gordo & Cabral, 2001).

Corresponding author:
M.G. Castro
Email: gcastro@mdp.edu.ar



Journal of Fish Biology (2011) 78, 1336–1358

doi:10.1111/j.1095-8649.2011.02937.x, available online at wileyonlinelibrary.com

Morphological, morphometric, meristic and osteological evidence for two species of hake (Actinopterygii: Gadiformes: *Merluccius*) in Argentinean waters

J. M. DÍAZ DE ASTARLOA*†‡, S. I. BEZZI§, M. GONZÁLEZ CASTRO*†, E. MABRAGAÑA*†, D. HERNÁNDEZ§, S. M. DELPIANI*†, D. E. FIGUEROA*, M. B. COUSSEAU*, M. Y. DELI ANTONI*† AND L. TRINGALI§

*Departamento de Ciencias Marinas, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350, B7602AYL Mar del Plata, República Argentina, †Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Rivadavia 1917, Buenos Aires, República Argentina and ‡Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), Paseo Victoria Ocampo N° 1, Escolera Norte, B7602HSA Mar del Plata, República Argentina

(Received 17 February 2009, Accepted 4 February 2011)

Morphologically, both classic and landmark-based morphometry and meristic analyses of 241 specimens of *Merluccius*, along with the re-examination of six paratype specimens of *Merluccius hubbsi*, the holotype and three paratypes of *Merluccius patagonicus* and the syntype of *Merluccius australis* revealed the presence of only two species of *Merluccius* in Argentinean waters. Internal structures (kyonandibula, urotyal and sagitta otolith) of *M. hubbsi* were compared to those reported for *M. patagonicus* and were shown to have identical morphology. Type specimens of *M. patagonicus* showed a complete overlap in morphometric and meristic characters with *M. hubbsi*, whereas *M. australis* had a greater number of second dorsal and anal-fin rays, and more lateral-line scales. In addition, *M. australis* had a smaller eye and longer snout than *M. hubbsi* and *M. patagonicus*. The results indicate that there is no evidence for a third hake species in Argentinean waters. *Merluccius patagonicus* is a synonym of *M. hubbsi*.

© 2011 The Authors

Journal of Fish Biology © 2011 The Fisheries Society of the British Isles

Key words: landmarks; *M. australis*; *M. hubbsi*; *M. patagonicus*; Patagonia.

INTRODUCTION

Species of hake (*Merluccius* spp.) are widely distributed throughout temperate–cool waters of the continental shelf and shelf-break, in the Mediterranean Sea, eastern and western Atlantic Ocean, eastern and south-western Pacific Ocean, and the south-west Indian Ocean (Lloris *et al.*, 2003) where they constitute important fisheries resources. In Argentina, *Merluccius* spp. is the most important fish resource from an economic perspective. *Merluccius* spp. catches represented 63% of the total catch of the Argentine fleet at the end of the 1980s (Bezzi *et al.*, 1995). Declared landed commercial catches were 600 000 t per annum over 1995–1997. By 2007, landed

‡Author to whom correspondence should be addressed. Tel.: +54 223 4751 107; email: astarfoa@mdp.edu.ar

Studies on reproduction of the mullet *Mugil platanus* Günther, 1880 (Actinopterygii, Mugilidae) from the Mar Chiquita coastal lagoon, Argentina: Similarities and differences with related species

M. GONZÁLEZ-CASTRO^{1,2,4*}, G. J. MACCHI^{3,4}, & M. B. COUSSEAU¹

¹Laboratorio de Ictiología, Departamento de Ciencias Marinas, Universidad Nacional de Mar del Plata, Argentina, ²Universidad Autónoma Metropolitana-Iztapalapa Departamento de Hidrobiología, México, D.F México, ³Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), Mar del Plata, Argentina, and ⁴Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

(Received 12 July 2010; accepted 6 December 2010)

Abstract

The aim of the present study is to provide the first results on the reproductive biology of *Mugil platanus*, from the Mar Chiquita coastal lagoon (Argentina). To address this topic, we employed mainly female gonads, at both microscopic and macroscopic levels. The following stages of oocyte development were observed: (A) oogonias, (B) primary growth oocyte, (C) cortical alveolus stage, (D) yolked oocytes, and (E) atretic follicle. Also five of the seven stages of gonad maturity were macro and microscopically stated. The potential fecundity ranged between 1,002,026 and 2,548,769 yolked oocytes, with a mean of around 1,800,000 oocytes. The length at first maturity (L_{50}) was 450.6 total length (TL) and 367.7 standard length (SL) for females, 436.3 TL and 354.9 SL for males, showing a late sexual maturity for both sexes. Seasonal changes in gonadosomatic index (GSI) of females were observed. The mean GSI was ≤ 1 during six months of the year (January, June–October). Two modes in the ovarian maturation were observed, the most important in April–May and a secondary one in November–December. Both modes, correlated with the CPUE values, are indicative of the mullet migration from the coastal lagoon towards the sea for spawning. Differences and similarities have been found between *Mugil platanus* and other mugilid species (*Mugil cephalus*, *Mugil curema*, *Liza aurata*, *Liza abu*, *Liza argentea*, *Myxus elongatus*, and *Valamugil cumesius*), from the reproductive point of view. The length at first maturity obtained for *M. platanus* is higher than in other mugilid species and when a relationship between maximum length and L_{50} is established the percentage corresponding to *Mugil platanus* is 75%, while the others range between 50 and 61%.

Keywords: Fish reproduction, gonad histology, fecundity, Mugilidae, Mugil platanus

Introduction

Fish species of Mugilidae occur in both coastal marine and brackish waters from all tropical and temperate seas (Nelson 2006). Conservative morphological features make identification of mullet species difficult. Recently, in order to clarify the controversy regarding the taxonomic status of the striped mullet *Mugil platanus*, in relation to the cosmopolite *Mugil cephalus*, a comprehensive analysis using sequences of the mitochondrial gene cytochrome *b*, landmark-based morphometry and meristic data have been performed (González Castro et al. 2008). The demonstrated discontinuity in geographic distribution and the mitochondrial DNA, morphometric and meristic

analysis allow these authors to consider *Mugil cephalus* and *Mugil platanus* as valid allopatric nominal species.

The striped mullet *Mugil platanus* is the only mugilid of permanent presence in Argentina, with a wide distribution along the coast, coastal lagoons and some fresh water beds (Cousseau et al. 2005; González Castro 2007). However, the white mullet *Mugil curema* Valenciennes, 1836 has been occasionally captured (González Castro et al. 2006).

Mugil platanus is commercially exploited not only in the south of Brazil (Vieira & Scalabrin 1991), but also in Samborombón Bay, Argentina (González Castro 2007). It is a gonochoristic and ovuliparous species, which shows high fecundity and migratory

*Correspondence: M. González-Castro, Laboratorio de Ictiología, Departamento de Ciencias Marinas, Universidad Nacional de Mar del Plata, Argentina. Tel: 54 92234751107. Email: gocastro@mdp.edu.ar

Zoological Studies 51(8): 1515-1528 (2012)

*Zoological
Studies*

Assessment of Lineal Versus Landmark-Based Morphometry for Discriminating Species of Mugilidae (Actinopterygii)

Mariano González-Castro^{1,2,3,*}, Ana Laura Ibáñez², Sandra Heras⁴, María Inés Roldán⁴, and María Berta Cousseau^{1,5}

¹Grupo de Biotaxonomía Morfológica y Molecular de Peces, IIMyC-Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Univ. Nacional de Mar del Plata, Mar del Plata, Argentina

²Univ. Autónoma Metropolitana-Iztapalapa Departamento de Hidrobiología. Av. San Rafael Atlixco 186, Col. Vicentina. México, D.F. 09340, México

³Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET, Argentina

⁴Laboratori d'Ictiologia Genètica, Univ. de Girona, Campus Montilivi, Girona 17071, Spain

⁵Laboratorio de Ictiología, Departamento de Ciencias Marinas, Univ. Nacional de Mar del Plata, Mar del Plata, Argentina

(Accepted December 19, 2012)

Mariano González-Castro, Ana Laura Ibáñez, Sandra Heras, María Inés Roldán, and María Berta Cousseau (2012) Assessment of lineal versus landmark-based morphometry for discriminating species of Mugilidae (Actinopterygii). *Zoological Studies* 51(8): 1515-1528. Meristic and different morphometric approaches were employed to assess the discrimination of 7 species of Mugilidae fishes (*Mugil cephalus*, *M. iliza*, *M. curema*, *M. hospes*, *Liza aurata*, *L. ramada*, and *Chelon labrosus*), but also to contribute to a better understanding of body-shape differences among this valuable species group. Three types of variables and their corresponding morphometric approaches were employed: 1) linear morphometrics measurements (LMMs); 2) interlandmark distances (IIDs); and 3) coordinate data (landmarks). Before the analyses, data exhibiting allometric growth were normalized. Data analysis included a one-way ANOVA (meristic data), a principal component analysis (PCA), and a cross-validated discriminant analysis (DA). The ANOVA showed significant differences in both lateral and transverse series scales. The PCA based on LMMs allowed the characterization of 6 groups, although some overlap between them was detected. The DA correctly classified 68.4% of the fishes according to their LMMs. The centroids of the 6 groups were separated for both the 1st and 2nd discriminant functions. The morphometric analysis based on IIDs yielded the best discrimination rates of the 3 approaches employed (96% for the DA). In the geometric morphometric analysis, the DA correctly classified 83.8% of the fishes according to their body shape. Although 6 groups were defined, some overlap among samples was detected. *Mugil hospes* was the best defined and most isolated species as observed in both the PCA and DA. Interestingly, the 3 morphometrics approaches employed separated *M. curema* specimens in 2 groups (Argentinean and Mexican samples). Moreover, European and Mexican samples of *M. cephalus* plotted separately in the PCA of the LMM- and IID-based approaches. These shape differences among *M. curema* of Argentina/Mexico and *M. cephalus* of Europe/Mexico reinforce the current hypothesis of a species complex, or even undescribed species as previously suggested by the authors.
<http://zoostud.sinica.edu.tw/Journals/51.8/1515.pdf>

Key words: Mugilidae, Landmarks, Meristic characters, Morphometry, Multivariate analysis.

Members of the Mugilidae, called mullet, are ray-finned fishes that usually inhabit coastal marine

and brackish waters in tropical and temperate seas (Thomson 1997, Nelson 2006). This family

*To whom correspondence and reprint requests should be addressed. Tel: 54-223-5456643. E-mail: gcastro@mdp.edu.ar



Mariano González Castro, Profesor Titular de Ictiología (2009-2010) en la Estación Hidrobiológica, Facultad de Hidrobiología de la Universidad Autónoma Metropolitana, México DF, 2009



En el río Las Estacas, a cargo de las prácticas de la asignatura Ictiología (Necton I) de la Universidad Autónoma Metropolitana, México DF, 2010

ProBiota

Serie Técnica y Didáctica **21 - Colección Semblanzas Etnológicas** **Archivos Editados**

- 01 – *Pedro Carriquiriborde*. Hugo L. López y Justina Ponte Gómez.
- 02 – *Pablo Agustín Tedesco*. Hugo L. López y Justina Ponte Gómez.
- 03 – *Leonardo Ariel Venerus*. Hugo L. López y Justina Ponte Gómez.
- 04 – *Alejandra Vanina Volpedo*. Hugo L. López y Justina Ponte Gómez.
- 05 – *Cecilia Yanina Di Prinzio*. Hugo L. López y Justina Ponte Gómez.
- 06 – *Juan Martín Díaz de Astarloa*. Hugo L. López y Justina Ponte Gómez.
- 07 – *Alejandro Arturo Dománico*. Hugo L. López y Justina Ponte Gómez.
- 08 – *Matías Pandolfi*. Hugo L. López y Justina Ponte Gómez.
- 09 – *Leandro Andrés Miranda*. Hugo L. López y Justina Ponte Gómez.
- 10 – *Daniel Mario del Barco*. Hugo L. López y Justina Ponte Gómez.
- 11 – *Daniel Enrique Figueroa*. Hugo L. López y Justina Ponte Gómez.
- 12 – *Luis Alberto Espínola*. Hugo L. López y Justina Ponte Gómez.
- 13 – *Ricardo Jorge Casaux*. Hugo L. López y Justina Ponte Gómez.
- 14 – *Manuel Fabián Grosman*. Hugo L. López y Justina Ponte Gómez.
- 15 – *Andrea Cecilia Hued*. Hugo L. López y Justina Ponte Gómez.
- 16 – *Miguel Angel Casalnuovo*. Hugo L. López y Justina Ponte Gómez.
- 17 – *Patricia Raquel Araya*. Hugo L. López y Justina Ponte Gómez.
- 18 – *Delia Fabiana Cancino*. Hugo L. López y Justina Ponte Gómez.
- 19 – *Diego Oscar Nadalin*. Hugo L. López y Justina Ponte Gómez.

Esta publicación debe citarse:

López, H. L. & J. Ponte Gómez. 2013. Semblanzas Ictiológicas: *Mariano González Castro*. *ProBiota*, FCNyM, UNLP, La Plata, Argentina, *Serie Técnica y Didáctica* 21(20): 1-13. ISSN 1515-9329.

ProBiota

(Programa para el estudio y uso sustentable de la biota austral)

Museo de La Plata
Facultad de Ciencias Naturales y Museo, UNLP
Paseo del Bosque s/n, 1900 La Plata, Argentina

Directores

Dr. Hugo L. López

hlopez@fcnym.unlp.edu.ar

Dr. Jorge V. Crisci

crisci@fcnym.unlp.edu.ar

Versión electrónica, diseño y composición

Justina Ponte Gómez

División Zoología Vertebrados

Museo de La Plata

FCNyM, UNLP

jpg_47@yahoo.com.mx

<http://ictiologiaargentina.blogspot.com/>

<http://raulringuelet.blogspot.com.ar/>

<http://aquacomm.fcla.edu>

<http://sedici.unlp.edu.ar/>

Indizada en la base de datos ASFA C.S.A.