

ProBiota, FCNyM, UNLP
ISSN 1515-9329

Serie Técnica y Didáctica n° 24(01)

Semblanzas Ictiológicas Iberoamericanas
Franco Andrés Teixeira de Mello Ramírez



Hugo L. López
y
Justina Ponte Gómez

Indizada en la base de datos ASEFA C.S.A.
2014



“El tiempo es invención o no es nada en absoluto”. Henri Bergson

“El tiempo es olvido y es memoria”. Jorge. L. Borges

A través de esta nueva serie tratamos de conocer diferentes aspectos personales de los integrantes de la comunidad ictiológica iberoamericana.

Esta iniciativa, comparte el espíritu y objetivo de las semblanzas nacionales buscando informalmente, otro punto de unión en la “comunidad de ictiólogos iberoamericanos”.

Quizás esté equivocado en mi apreciación, pero creo que vale la pena este intento, ya que, con la colaboración generosa e insoslayable de los integrantes de este “universo”, señalaremos un registro en el tiempo de la *Ictiología Neotropical*.

Hugo L. López

Imagen de Tapa

Franco Teixeira de Mello - Pesca eléctrica en el arroyo Mattrup, Dinamarca, 2010

Imagen de fondo

Porque en realidad nuestro norte es el sur, dibujo de Joaquín Torres García

Semblanzas Ictiológicas Iberoamericanas
Franco Andrés Teixeira de Mello Ramírez



Instalación de experimentos de exclusión de peces en el arroyo Gudena, Dinamarca, 2009

Hugo L. López y Justina Ponte Gómez

ProBiota
División Zoología Vertebrados
Museo de La Plata
FCNyM, UNLP

Marzo, 2014

Nombre y apellido completos: Franco Andrés Teixeira de Mello Ramírez
Lugar y fecha de nacimiento: Tacuarembó, Uruguay
Lugar, provincia y país de residencia: Maldonado, Uruguay
Título máximo, Facultad y Universidad: Doctor en Biología, Opción Ecología, Facultad de Ciencias, Universidad de la República
Posición laboral: Asistente Grado 2- Dedicación total
Lugar de trabajo: Centro Universitario Regional Este- Maldonado
Especialidad o línea de trabajo: Ecología de sistemas acuáticos- Biomonitoreo
Correo electrónico: frantei@fcien.edu.uy

Cuestionario

- **Un libro:** Las enseñanzas de Don Juan de Carlos Castaneda
- **Una película:** Nada es para siempre (A river run through it)
- **Un tema musical:** Cancioncilla amorosa, Eduardo Mateo
- **Un artista:** Eduardo Mateo
- **Un deporte:** pesca
- **Un color:** rojo
- **Una comida:** asado
- **Un animal:** pez
- **Una palabra:** libertad
- **Un número:** 7
- **Una imagen:** mi hermano (1982-2011)
- **Un lugar:** Tacuarembó
- **Una estación del año:** verano
- **Un nombre:** lara
- **Un hombre:** mi padre
- **Una mujer:** Dani
- **Un personaje de ficción:** Homero Simpson
- **Un superhéroe:** mi madre

*Tararira - Tararira
que arisca y sabia que estás
anzuelo que cae al agua
mojarra que te llevás
pica pica Tararira
plata viva del juncal
mientras no se corte el hilo
junto al agua me hallarás...*

Osiris Rodríguez Castillo



Noche en Las Cañas
Franco Teixeira de Mello con su esposa Dani, río Uruguay, República Oriental del Uruguay, 2009



Franco Teixeira de Mello con su esposa Dani y su hija Lara , Pan de Açúcar, Río de Janeiro, Brasil, 2013

Journal of Fish Biology (2006) **69**, 1770–1778

doi:10.1111/j.1095-8649.2006.01245.x, available online at <http://www.blackwell-synergy.com>

Ontogenetic allometric coefficient changes: implications of diet shift and morphometric traits in *Hoplias malabaricus* (Bloch) (Characiforme, Erythrinidae)

F. TEIXEIRA DE MELLO*†‡§, C. IGLESIAS*†, A. I. BORTHAGARAY*†, N. MAZZEO*, J. VILCHES*, D. LARREA* AND R. BALLABIO*†

*Aquatic Systems Research and Recovery Group, Ecology Department, Facultad de Ciencias, Universidad de la República, Iguaú 4225, CP. 11400 Montevideo, Uruguay. †Basic and Applied Ecology Research Group, NGO Investigación y desarrollo (I+D), Iguaú 4225, CP. 11400 Montevideo, Uruguay and ‡Ecotoxicology and Environmental Chemistry Group, Facultad de Ciencias, Universidad de la República, Iguaú 4225, CP. 11400 Montevideo, Uruguay

(Received 11 November 2005, Accepted 21 July 2006)

This study evaluated the relationship between body size and digestive tract characteristics of the important predatory freshwater fish *Hoplias malabaricus*, which is widely distributed in South America. The allometric coefficients were calculated for the mass and standard length (L_t) relationships for two different L_t groups: (1) between 20 and 100 mm (characterized as insectivores) and (2) > 100 mm (characterized as piscivores). Differential growth measured from the allometric coefficient, b , between the insectivore ($b < 3$) and the piscivore ($b > 3$) groups was detected. Anterior intestine length and pyloric caeca zone length showed significant differences between groups. Two complementary hypotheses were developed to explain the differential growth: (1) *H. malabaricus* has a digestive tract adapted to a piscivorous diet, which is independent of its ontogenetic stage of development, and (2) the negative allometry observed in group 1 individuals agrees with a general behavioural strategy, allowing individuals to grow in L_t during a shorter period of time.

© 2006 The Authors

Journal compilation © 2006 The Fisheries Society of the British Isles

Key words: allometric coefficient; diet shift; *Hoplias malabaricus*; ontogenetic changes; South America.

INTRODUCTION

The lack of basic biological knowledge about neotropical fish communities and the implications for the development of management and conservation measures have been deplored by many researchers (Menni *et al.*, 1984; Buti, 1995; Buti & Miquelarena, 1995; Vari & Malabarba, 1998; Lizama & Ambrósio, 2002). Analyses of mass–length relationships contribute to reducing this lack of knowledge.

The allometric coefficient of the mass–length relationships is influenced by, among other factors, the quantity and quality of the ingested food. Many

§Author to whom correspondence should be addressed. Tel: +598 2 5258618 (ext. 162); fax: +598 2 5258617; email: frantoi@fcien.edu.uy

Substantial differences in littoral fish community structure and dynamics in subtropical and temperate shallow lakes

FRANCO TEIXEIRA-DE MELLO^{a,c,d}, MARIANA MEERHOFF^{a,b}, ZEYNEP PEKCAN-HEKIM^e AND ERIK JEPPESEN^{b,***}

^aGrupo de Ecología y Rehabilitación de Sistemas Acuáticos, Departamento de Ecología, Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay

^bGrupo de Investigación en Ecotoxicología y Química Ambiental, Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay

^cAsociación Investigación y Desarrollo I+D, Montevideo, Uruguay

^dNational Environmental Research Institute, University of Aarhus, Silkeborg, Denmark

^eDepartment of Biological and Environmental Sciences/Aquatic Sciences, University of Helsinki, Helsinki, Finland

^{**}Department of Biological Sciences, University of Aarhus, Aarhus, Denmark

SUMMARY

1. Fish play a key role in the functioning of temperate shallow lakes by affecting nutrient exchange among habitats as well as lake trophic structure and dynamics. These processes are, in turn, strongly influenced by the abundance of submerged macrophytes, because piscivorous fish are often abundant at high macrophyte density. Whether this applies to warmer climates as well is virtually unknown.

2. To compare fish community structure and dynamics in plant beds between subtropical and temperate shallow lakes we conducted experiments with artificial submerged and free-floating plant beds in a set of 10 shallow lakes in Uruguay (30°–35°S) and Denmark (55°–57°N), paired along a gradient of limnological characteristics.

3. The differences between regions were more pronounced than differences attributable to trophic state. The subtropical littoral fish communities were characterised by higher species richness, higher densities, higher biomass, higher trophic diversity (with predominance of omnivores and lack of true piscivores) and smaller body size than in the comparable temperate lakes. On average, fish densities were 93 ind. m⁻² (±10 SE) in the subtropical and 10 ind. m⁻² (±2 SE) in the temperate lakes. We found a twofold higher total fish biomass per unit of total phosphorus in the subtropical than in the temperate lakes, and as fish size is smaller in the former, the implication is that more energy reaches the littoral zone fish community of the warmer lakes.

4. Plant architecture affected the spatial distribution of fish within each climate zone. Thus, in the temperate zone fish exhibited higher densities among the artificial free-floating plants while subtropical fish were denser in the artificial submerged plant beds. These patterns appeared in most lakes, regardless of water turbidity or trophic state.

5. The subtropical littoral fish communities resembled the fish communities typically occurring in temperate eutrophic and hypertrophic lakes. Our results add to the growing evidence that climate warming may lead to more complex and omnivory-dominated food webs and higher density and dominance of smaller-sized fish. This type of community

Correspondence: Mariana Meerhoff and Franco Teixeira-de Mello, Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay. E-mail: merluz@fcien.edu.uy; frantei@fcien.edu.uy

Hydrobiologia (2012) 684:143–160
 DOI 10.1007/s10750-011-0979-7

PRIMARY RESEARCH PAPER

Community structure of fish in lowland streams differ substantially between subtropical and temperate climates

Franco Teixeira-de Mello · Mariana Meerhoff · Annette Baatrup-Pedersen · Thomas Maigaard · Peter B. Kristensen · Torben K. Andersen · Juan M. Clemente · Claudia Fosalba · Esben A. Kristensen · Malvina Masdeu · Tenna Riis · Néstor Mazzeo · Erik Jeppesen

Received: 8 July 2011 / Revised: 28 November 2011 / Accepted: 3 December 2011 / Published online: 4 January 2012
 © Springer Science+Business Media B.V. 2011

Abstract Fish are important in the structuring of other communities and may have large effects on the functioning of aquatic ecosystems. The structure of fish communities, in turn, seems to differ with climate. We compared the characteristics of fish assemblages in lowland streams located in two contrasting climates (cold-temperate Europe and subtropical South America) by use of published and unpublished data on streams of similar depth, width, and slope (n total = 91 streams). We also selected a subset of seven comparable little-affected streams in the two contrasting

climates: temperate (Denmark, 55°–57°N, Dk) and subtropical (Uruguay, 30°–35°S, Uy) and compared the fish community structures in relation to environmental characteristics. We then analysed a series of potential explanatory factors behind the patterns observed, in particular the effect of ambient temperature, by comparing temperature-corrected community metabolism. Significantly higher species richness, higher densities, lower biomass, smaller mean body size, and lower mean weight of fish were observed for the subtropical streams than for the temperate streams, both in the literature review and in the subset of streams. Several characteristics of fish assemblages in streams may be explained by direct and indirect effects of temperature. Accordingly, fish in subtropical systems had a

Electronic supplementary material The online version of this article (doi:10.1007/s10750-011-0979-7) contains supplementary material, which is available to authorized users.

Handling editor: Sidinei Magela Thomaz

F. Teixeira-de Mello (✉) · M. Meerhoff · J. M. Clemente · C. Fosalba · M. Masdeu · N. Mazzeo
 Departamento de Ecología & Evolución, CURE-Facultad de Ciencias, Universidad de la República, Maldonado, Uruguay
 e-mail: frantei@fcien.edu.uy

F. Teixeira-de Mello · M. Meerhoff · J. M. Clemente · C. Fosalba
 Asociación Civil Investigación y Desarrollo I+D, Iguaú 4225, CP 11400 Montevideo, Uruguay

M. Meerhoff · A. Baatrup-Pedersen · T. Maigaard · P. B. Kristensen · T. K. Andersen · E. A. Kristensen · E. Jeppesen
 Department of Bioscience, Aarhus University, Vejløvej 25, 8600 Silkeborg, Denmark

T. Maigaard · P. B. Kristensen · T. K. Andersen · T. Riis
 Department of Bioscience, Aarhus University, Ole Worms Allé, Building 135, 8000 Aarhus, Denmark

J. M. Clemente
 Departamento de Medio Ambiente, Laboratorio Tecnológico del Uruguay, Av Italia 6201, Montevideo, Uruguay

E. Jeppesen
 Greenland Climate Research Centre (GCRC), Greenland Institute of Natural Resources, Kivioq 2, P.O. Box 570, 3900 Narsik, Greenland

E. Jeppesen
 Sino-Danish Centre for Education and Research (SDC), Beijing, China

Environ Monit Assess (2014) 186:1665–1677
 DOI 10.1007/s10661-013-3484-9

Monitoring fish communities in wadeable lowland streams: comparing the efficiency of electrofishing methods at contrasting fish assemblages

Franco Teixeira-de Mello · Esben A. Kristensen · Mariana Meerhoff ·
 Iván González-Bergonzoni · Annette Baatrup-Pedersen · Carlos Iglesias ·
 Peter B. Kristensen · Néstor Mazzeo · Erik Jeppesen

Received: 17 February 2013 / Accepted: 30 September 2013 / Published online: 5 November 2013
 © Springer Science+Business Media Dordrecht 2013

Abstract Electrofishing is considered a reliable tool to assess the assemblages and biodiversity of fish in wadeable streams. The most widely used electrofishing techniques (point [P], single-pass [S-P], and multiple-pass [M-P]) vary as to the effort needed for sample collection, and this may potentially influence the degree of accuracy. Moreover, little is known about the comparability of the methods and their specific performance in streams with different fish assemblages. The aim of this investigation was to validate (using M-P sampling as reference) the use of P and S-P electrofishing techniques to accurately assess the richness, density and size distribution of fishes in small streams at both regional and global scale independently of fish assemblages and geographical region. We sampled 50-m-long reaches in a total of 33 lowland stream reaches that were

located in different climatic and biogeographical regions (Uruguay and Denmark) and hosted different fish assemblages. Subtropical fish communities exhibited higher richness (Uy: 12–32, Dk: 1–9) and densities (Uy: 1.3–5.2, DK: 0.1–4.9 ind. m⁻²) than temperate streams. We applied both "global models" using the entire database (33 sites) and "local models" including the same number of sites but using the climatic region as a model variable. Regression analyses revealed that the P, S-P and M-P methods all provided an adequate picture of the species composition and size distribution, and transfer equations for comparison between methods are thus not required. Conversely, richness was better predicted by S-P and by P techniques for regional and global models, respectively. Transfer equations obtained for abundance revealed that the P and S-P models can accurately transform catch data

F. Teixeira-de Mello · M. Meerhoff ·
 I. González-Bergonzoni · C. Iglesias · N. Mazzeo
 Departamento de Ecología & Evolución,
 CURE-Facultad de Ciencias, Universidad de la República,
 Tacuarembó s/n, Maldonado CP 20000, Uruguay

F. Teixeira-de Mello · M. Meerhoff ·
 I. González-Bergonzoni · C. Iglesias
 Asociación Civil Investigación y Desarrollo I+D,
 Iguaí 4225, CP 11400 Montevideo, Uruguay

E. A. Kristensen · M. Meerhoff · I. González-Bergonzoni ·
 A. Baatrup-Pedersen · P. B. Kristensen · E. Jeppesen
 Department of Bioscience, Aarhus University,
 Vejløvej 25, 8600 Silkeborg, Denmark

I. González-Bergonzoni · E. Jeppesen
 Sino-Danish Centre for Education and Research (SDC),
 Beijing, China

E. Jeppesen
 Greenland Climate Research Centre (GCRC),
 Greenland Institute of Natural Resources,
 Kivioq 2, P.O. Box 570 3900, Narsak, Greenland

F. Teixeira-de Mello (✉)
 Ecología y Rehabilitación de Sistemas Acuáticos,
 CURE-Facultad de Ciencias, Universidad de la República,
 Tacuarembó s/n, Maldonado CP 20000, Uruguay
 e-mail: franstei@fcien.edu.uy



Llegando a la laguna Sinha Mariana, Pantanal, cerca de Rondonopolis, Mato Grosso do Sul, Brasil, 2013

Esta publicación debe citarse:

López, H. L. & J. Ponte Gómez. 2014. Semblanzas Ictiológicas Iberoamericanas: *Franco Andrés Teixeira de Mello Ramírez*, FCNyM, UNLP, La Plata, Argentina, *Serie Técnica y Didáctica* 24(01): 1-12. ISSN 1515-9329.

ProBiota

(Programa para el estudio y uso sustentable de la biota austral)

Museo de La Plata
Facultad de Ciencias Naturales y Museo, UNLP
Paseo del Bosque s/n, 1900 La Plata, Argentina

Directores

Dr. Hugo L. López

hlopez@fcnym.unlp.edu.ar

Dr. Jorge V. Crisci

crisci@fcnym.unlp.edu.ar

Versión electrónica, diseño y composición

Justina Ponte Gómez

División Zoología Vertebrados

Museo de La Plata

FCNyM, UNLP

jpg_47@yahoo.com.mx

<http://ictiologiaargentina.blogspot.com/>

<http://raulringuelet.blogspot.com.ar/>

<http://aquacomm.fcla.edu>

<http://sedici.unlp.edu.ar/>

Indizada en la base de datos ASFA C.S.A.