

FRESHWATER BIOLOGICAL ASSOCIATION



FRESHWATER FORUM



1995

VOLUME 5

NUMBER 1

FRESHWATER FORUM

VOLUME 5: NUMBER 1 1995

Published by the Freshwater Biological Association
The Ferry House, Far Sawrey, Ambleside, Cumbria LA22 0LP

© Freshwater Biological Association 1995

ISSN 0961-4664

FISHES OF THE LAGUNAS ENCADENADAS (PROVINCE OF BUENOS AIRES, ARGENTINA), A WETLAND OF INTERNATIONAL IMPORTANCE

AMALIA M. MIQUELARENA AND HUGO L. LÓPEZ

(Dr A. M. Miquelarena (CONICET) and Dr H. L. López (CIC), Instituto de Limnología "Dr Raúl A. Ringuelet", C.C. 712, 1900 La Plata, Argentina, and Departamento Científico Zoología Vertebrados, Museo de la Plata, Paseo del Bosque s/n, 1900 La Plata, Argentina.)

Introduction

The Lagunas Encadenadas (36° 30' – 37° 30' S, 61° 00' – 63° 30' W) form a large endorheic wetland, i.e. the drainage lies in an enclosed basin. The wetland is situated in the south-west of the Province of Buenos Aires and has a total area of nearly 58,000 hectares (Fig. 1). The system comprises a chain of five main lagoons (Alsina, Chochicó, Del Monte, Del Venado, and Epecuen) (Fig. 2), distinguished by a well defined east–west gradient in salinity from fresh water to saline.

From a hydrological viewpoint these waterbodies are part of the "diagonal group" of a larger lacustrine basin (Frenguelli 1956) along the northeastern slope of the Sierra de la Ventana (Mazza 1962; Ringuelet 1962). Periodic floods affect the zone, often with critical consequences, as in 1986 when the town Villa Epecuen was completely covered by water. As a result of flooding, in recent years there has been an enlargement of the inundated zone: the modified landscape includes land/inland water ecotones biologically unique due to the biodiversity they support (López et al. 1993; Eerden & Iedema 1994). The chain of lagoons form the most southerly wetland in South America having this particular combination of hydrological, physico-chemical and biological characters. Although it is an isolated system, a few years ago a canal for water-level control in dry periods was constructed. This connects the lagoons to the Salado River drainage in the north, though it is temporarily closed now because of the recent inundations in the 1980s.

Concerning the regional biogeographical aspect, this wetland is within the Pampasian dominion of the Guayano–Brazilian subregion (Ringuelet 1961, 1981). In addition, it may be regarded as a transition zone between the Pampasian and Sub-Andean dominions, also influenced by Patagonian elements. Ringuelet (1975) noted that the Lagunas Encadenadas are inhabited by fish species from the meridional pampean plains, in the south of the Salado River basin. Additional information about the fish fauna from this region is provided by Mac Donagh (1934), Ringuelet et al.

(1967), Gallardo (1970), Ringuelet (1975), Lüling (1981), Menni et al. (1988), and López et al. (1993).

The fishes of the Lagunas Encadenadas

During two recent collection trips to the Lagunas Encadenadas, the second author recorded thirteen species of fish; the material was caught at thirteen sampling localities. From the collected material and available data in the literature, we can make the following points.

The fish fauna of this wetland comprises eighteen species from eleven families and six orders (Table 1). The list also includes three species reported by Lüling (1981) that we did not confirm (marked with double asterisks in Table 1).




FIG. 1. Lagunas Encadenadas del Oeste within Argentina and the Province of Buenos Aires.

Four species, including one exotic, are new records for these waterbodies (marked with a single asterisk in Table 1).

With the exception of the cypriniform *Cyprinus carpio*, all of the other species belong to the Guayano-Brazilian subregion.

Fish species diversity decreases from east to west, i.e. from the most freshwater lagoon (Alsina, with 15 species) to the most saline one (Epecuen, 1 species). A similar diversity pattern has been found in zooplankton species (López et al. 1993).

Lüling (1981) and other authors have already noted the remarkably high salinity at Epecuen, accounting for the presence of the crustacean *Artemia salina*, and noting the absence of any fish species. Our new record of *Jenynsia lineata*, found along the shore line of Epecuen, is supported by the remarkable euryhalinity of this species but also may reflect recent changes in physico-chemical conditions of the lagoon, resulting from the recent overall alteration to the system of lagoons.




FIG. 2. Llagunas Encadenadas showing the range of extension of water during the period 1981 to 1993 (modified from Eerden & Iedema 1994).

Table 1. Fish species from the Lagunas Encadenadas del Oeste (Buenos Aires, Argentina).

	Alsina	Cochico	Del Monte	Del Venado	Epecuen
* <i>Cyprinus carpio</i>	X				
<i>Oligosarcus jenynsi</i>	X	X	X	X	
<i>Astyanax cf. eigenmanniorum</i>	X	X	X	X	
** <i>Astyanax taeniatus</i>	X	X			
<i>Bryconamericus iheringi</i>	X	X		X	
<i>Cheirodon cf. interruptus</i>	X	X		X	
** <i>Cheirodon cf. galusdae</i>	X	X			
<i>Cyphocharax boga</i>	X	X	X	X	
** <i>Galeocharax humeralis</i>	X	X			
* <i>Hoplias malabaricus</i>	X				
<i>Corydoras paleatus</i>	X	X	X		
* <i>Loricariichthys anus</i>		X			
* <i>Parapimelodus valenciennis</i>	X				
<i>Rhamdia sapo</i>	X	X	X		
<i>Cnesterodon decemmaculatus</i>				X	
<i>Jenynsia lineata</i>		X	X	X	X
<i>Odontesthes bonariensis</i>	X	X	X	X	
<i>Cichlasoma facetum</i>	X	X			
Number of species:	15	14	7	8	1

The presence of the exotic *Cyprinus carpio* shows the extent of the advance of this species southwards, and the progressive occupation of the waterbodies of the Province of Buenos Aires (López et al. 1994).

The records of *Hoplias malabaricus*, *Loricariichthys anus* and *Parapimelodus valenciennis* extend the distribution area of three species of Brazilian biogeographical lineages. Therefore, they have to be listed together with the highest widespread neotropical taxa (i.e. *Rhamdia*, *Astyanax* and *Jenynsia*) that reach the most southern and western limits of the Guayano–Brazilian subregion. These new reports should be considered in association with the artificial canal that now connects the lagoon system to the Salado River basin, enhancing the advance of fish populations. Their ultimate adaptation to this particular lentic environment depends on the ecological fidelity and resistance capability of each species (Ringuelet 1975).

A wetland of international importance

In agreement with Eerden & Iedema (1994) we believe that the Lagunas Encadenadas form a wetland of international importance. Moreover, because of its land/inland water ecotone character, the study of these waterbodies should be integrated into the Ecotone Programme, within the Man and the Biosphere Programme (MAB-UNESCO) (Décamps 1990).

Acknowledgements

The authors thank Adriana Aquino for the English translation and suggestions, and Roberto Menni for commenting on the manuscript.

References

- Décamps, H., Fournier, F., Naiman, R. J. & Petersen Jr., R. C. (1990). An International Research Effort on Land/Inland Water Ecotones in Landscape Management and Restoration 1990-1996. *Ambio*, **19**, 175-176.
- Eerden van, M. R. & Iedema, C. W. (1994). *The Lagunas Encadenadas del Oeste. Pearls of The Pampas. Ecological Aspects of Flood Control Alternatives for the Lagunas Encadenadas*. Ministry of Transport, Public Works and Watermanagement, Directorate Flevoland The Netherlands, Prov. of Buenos Aires, Argentina. 53 pp.
- Frenquelli, J. (1956). Rasgos generales de la hidrografía de la provincia de Buenos Aires. *LEMIT, Ser. II, No. 62*, 19 pp.
- Gallardo, J. M. (1970). Estudio ecológico sobre anfibios y reptiles del sudoeste de la provincia de Buenos Aires. *Rev. Mus. Arg. Cs. Nat. "Bernardino Rivadavia", Zool.*, **10**, 27-63.
- López, H. L., Padin, O. H. & Iwaszkiw, J. M. (1993). *Biología Pesquera de las Lagunas Encadenadas del Sudoeste, Provincia de Buenos Aires*. Inf. Técnico, IATASA, Buenos Aires. 64 pp., t. and figs.
- López, H. L., Protogino, L. C. & Aquino, A. E. (1994). *Ichtiofauna de la Argentina (Provincias de Santiago del Estero, Catamarca, Córdoba, San Luis, La Pampa y Buenos Aires)*. Dto. Cient. Zool. Vertebrados, Museo de La Plata (Ictiología), 23 pp., La Plata.
- Lüling, K. H. (1981). Wissenschaftliche ergebnisse des forschung saufenathaltes Dr. K. H. Lüling in Argentinien 1975/76. I. Ichthyologische und gewasserkundliche beobachtungen und untersuchungen an der seenplate von Guamini (Sudl. Prov. Buenos Aires, Argentinien). *Zool. Beitr.*, **27**, 1-24.
- Mac Donagh, E. J. (1934). Nuevos conceptos sobre la distribución geográfica de los peces argentinos basados en expediciones del Museo de La Plata. *Rev. Mus. La Plata*, **24**, 21-170.

- Mazza, G. (1962). Recursos hidráulicos superficiales. *Serie Evaluación de los Recursos Naturales de la Argentina (Primera Etapa), Consejo Federal de Inversiones*, 4 (1), 1-459.
- Menni, R. C., López, H. L. & Arámburu, R. H. (1988). Ictiofauna de Sierra de la Ventana y Chasicó (Provincia de Buenos Aires, Argentina). Zoogeografía y parámetros ambientales. *An. Mus. Hist. Nat. Valparaíso*, 19, 75-84.
- Ringuelet, R. A. (1961). Rasgos fundamentales de la zoogeografía Argentina. *Physis*, 22(63), 151-170.
- Ringuelet, R. A. (1962). *Ecología Acuática Continental*. Manuales EUDEBA, IX, Buenos Aires, 138 pp.
- Ringuelet, R. A. (1975). Zoogeografía y ecología de los peces de aguas continentales de la Argentina y consideraciones sobre las áreas ictiológicas de América del Sur. *Ecosur.*, 2(3), 1-122.
- Ringuelet, R. A. (1981). El ecotono faunístico subtropical pampásico y sus cambios históricos. *Symposia, VI Jorn. Arg. Zool., La Plata*, 75-80.
- Ringuelet, R. A., Arámburu, R. H. & de Arámburu, A. A. (1967). Los Peces Argentinos de Agua Dulce. *Com. Inv. Cient., Prov. Bs. As.* 602 pp.

Versión Electrónica

Justina Ponte Gómez

División Zoología Vertebrados

FCNyM

UNLP

Jpg_47@yahoo.com.mx