

## First report of *Rhabdias* sp. infecting *Leptodactylus macrosternum* from the Caatinga domain, Neotropical region

Diêgo Alves Teles<sup>1</sup>, Cícero Leonardo de Moraes Pinto<sup>2</sup>, Adonias Aphoena Martins Teixeira<sup>1</sup>, João Antonio de Araujo Filho<sup>1</sup>

<sup>1</sup> Programa de Pós-Graduação em Ciências Biológicas (Zoologia), Laboratório/Coleção de Herpetologia, Universidade Federal da Paraíba – UFPB , Cidade Universitária, Campus I, CEP 58059-900, João Pessoa, Paraíba, Brazil.

<sup>2</sup> Laboratorio de Zoologia/Parasitologia, Universidade Regional do Cariri –URCA, Campus Pimenta, CEP 63100-000, Crato, Ceará, Brazil.

Recibida: 27 Marzo 2018

### ABSTRACT

Revisada: 16 Mayo 2018

On 27 March 2011, 20 specimens of *Leptodactylus macrosternum* were collected in the municipality of Aiuba at the Ecological Station of Aiuba, state of Ceará, Brazil. Of the 20 frogs examined, four were infected with lung nematodes. Five specimens of *Rhabdias* sp. are found. To our knowledge, *L. macrosternum* is a new host for *Rhabdias* sp.

Aceptada: 29 Mayo 2018

Editor Asociado: C. Borteiro

doi: 10.31017/CdH.2018.(2018-09)

Key Words: Parasitism; Semi-arid region; Nematodes.

*Leptodactylus macrosternum* Miranda-Ribeiro, 1926, occurs in the northeast of Brazil (Frost, 2018). It is a medium-sized frog which features nocturnal activity and diet consisting mainly arthropods (Teles *et al.*, 2018), included consumption of crustaceans (Teles *et al.*, 2014.). Here, we report the nematode infection in the respiratory tracts of *L. macrosternum* in an area of Caatinga in Aiuba, Ceará, northeastern Brazil. Twenty specimens of *L. macrosternum* (15 males; mean SVL=  $63.67 \pm 7.71$  mm; range = 84.13 – 52.43 mm; five females; mean SVL=  $72.97 \pm 10.40$  mm; range 87.28 – 58.99 mm) were collected in the municipality of Aiuba at the Ecological Station of Aiuba, Brazil ( $6.573476^{\circ}$ S;  $40.123564^{\circ}$ W, datum SAD69; 466 m a.s.l.), on 27 March 2011 and examined for endoparasites. Each specimen was dissected by a longitudinal incision and respiratory and digestive tracts were removed and analyzed separately (stomach, small intestine and large intestine) and body cavity. Helminths were preserved in 70% alcohol and subsequently mounted on temporary slides using Hoyer's medium, and identified beneath a light microscope. Of the 20 frogs examined, four were infected with lung nematodes. The nematodes were identified (according to Vicente *et al.* 1990 and Kuzmin *et al.* 2015) as: *Rhabdias* sp. (Fig. 1), and later deposited in the parasitological collection of the Universidade Regional do Cariri (URCA-P: 518). The prevalence and intensity of infection were

calculated according to Bush *et al.* (1997). Overall prevalence was 20% (4/20) and the intensity of infection was 1.25. The prevalence in males was 20% (3/15) and intensity of infection 1.33. On the other hand, prevalence in females was 20% (1/5) and intensity of infection 1. The nematodes genus *Rhabdias* Stiles & Hassal, 1905 can be found infecting lungs of anuran amphibians and less commonly in lizards, snakes and salamanders (Kuzmin *et al.*, 2016; Teles *et al.*, 2014). In South America, there are 19 species of *Rhabdias* reported to affect the respiratory tract of amphibians (Kuzmin *et al.*, 2016; Campião *et al.*,



**Figure 1.** Specimen of *Rhabdias* sp. found in the lungs of *Leptodactylus macrosternum*.

2014). There are records of *Rhabdias* sp. infections in Bufonidae, Brachycephalidae, Hylidae, and Leptodactylidae in Argentina and Brazil (Campião *et al.*, 2014) and most recently in *Proceratophrys aridus* (Odontophrynididae), Caatinga domain, in Brazil (Teles *et al.*, 2017). There are still few studies on the effects of infections caused by nematodes of the genus *Rhabdias* on amphibians hosts. However, in an experimental study conducted by Goater and Ward (1992) they observed that in juvenile anurans of the *Bufo bufo* species they had their body growth and their survival reduced due to the infection caused by *Rhabdias bufonis*. The host specimens examined did not exhibit evident malformations or behavioral changes. To our knowledge, *L. macrosternum* is a new host for *Rhabdias* sp.

### Acknowledgements

We thank CNPq for providing a scholarship to DAT and JAAF and CAPES for the scholarship granted to AAMT. ICMBio granted permission to collect under license System Authorization and Information on Biodiversity, no. 27542-2, process no. 96683918.

### Literature cited

- Bush, A. O.; Lafferty, K.D.; Lotz, J.M. & Shostak, A.W. 1997. Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* 83: 575-583.
- Campião, K.M.; Morais, D.H.; Dias, O.T.; Aguiar, A.; Toledo, G.M.; Tavares, L.E.R. & Silva, J.R. 2014. Checklist of Helminth parasites of Amphibians from South America. *Zootaxa* 3843: 001-093.
- Frost, D.R. 2018 Amphibian Species of the World: an Online Reference. Version 6.0. American Museum of Natural History, New York, USA. Available at: <<http://research.amnh.org/vz/herpetology/amphibia/Amphibia/Anura/Leptodactylusmacrosternum>>. Last accessed: 20 may 2018.
- Goater, G.P. & Ward, P.I. 1992. Negative effects of *Rhabdias bufonis* (Nematoda) on the growth and survival of toads (*Bufo bufo*). *Oecologia* 89: 161-165.
- Kuzmin, Y.; du Preez, L.H. & Junker, K. 2015. Some nematodes of the genus *Rhabdias* Stiles et Hassall, 1905 (Nematoda: Rhabdiasidae) parasitising amphibians in French Guiana. *Folia Parasitologica* 62: 031.
- Kuzmin Y.; Melo F. T. V.; Silva Filho H. F. & Santos J. N. 2016. Two new species of *Rhabdias* Stiles et Hassall, 1905 (Nematoda: Rhabdiasidae) from anuran amphibians in Para, Brazil. *Folia Parasitologica* 63: 015.
- Teles, D.A.; Teixeira, A.P.M.; Araujo Filho, J.A.; Cabral, M.E.S.; Sales, R.M.A. & Dias, D.Q. 2014. *Leptodactylus macrosternum*. Diet. *Herpetological Review* 45: 304.
- Teles, D.A.; Cabral, M.E.S.; Araujo-Filho, J.A.; Dias, D.Q.; Ávila, R.W. & Almeida, W.O. 2014. Helminths of *Leptodactylus vastus* (Anura: Leptodactylidae) in area of Caatinga, Brazil. *Herpetology Notes* 7: 355-356.
- Teles, D.A.; Rodrigues, J.K.; Teixeira, A.A.M., Araujo-Filho, J.A.; Sousa, J.G.G. & Ribeiro, S.C. 2018. Diet of *Leptodactylus macrosternum* (Miranda-Ribeiro 1926) (Anura: Leptodactylidae) in the Caatinga domain, Northeastern Brazil, Neotropical Region. *Herpetology Notes* 11: 223-226.
- Teles, D.A.; Brito, S.V.; Araujo Filho, J.A., Teixeira, A.A.M., Ribeiro, S.C., Mesquita, D.O. & Almeida, W.O. 2017. Nematode parasites of *Proceratophrys aridus* (Anura: Odontophrynididae), an endemic frog of the Caatinga domain of the Neotropical Region in Brazil. *Herpetology Notes* 10: 525-527.
- Vicente, J.J., Rodrigues, H.O., Gomes, D.C. & Pinto, R.M. 1990. Nematóides do Brasil. Parte II: Nematóides de anfíbios. *Revista Brasileira de Zoología* 7: 549-626.