



First record of *Litomosoides pardinasi* (Nematoda: Onchocercidae) in native and exotic rodents from Chile

Primer registro de *Litomosoides pardinasi* (Nematoda: Onchocercidae) en roedores nativos y exóticos de Chile

Carlos Landaeta-Aqueveque¹, Juliana Notarnicola^{2✉}, Juana P. Correa³, Andrea Yáñez-Meza⁴, AnaLía Henríquez⁵, Pedro E. Cattán³, Carezza Botto-Mahan⁴ and Fernando Torres-Pérez⁶

¹Departamento de Patología y Medicina Preventiva, Facultad de Ciencias Veterinarias, Universidad de Concepción. Vicente Méndez 595, Chillán, Chile.

²Centro de Estudios Parasitológicos y de Vectores, CCT La Plata, Consejo Nacional de Investigaciones Científicas y Técnicas. Calle 2, Núm. 584, La Plata (1900), Buenos Aires, Argentina.

³Departamento de Ciencias Biológicas Animales, Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile. Santa Rosa 11735, Santiago Chile.

⁴Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile. Las Palmeras 3425, Ñuñoa, Santiago Chile.

⁵Facultad de Recursos Naturales y Medicina Veterinaria, Universidad Santo Tomás. Manuel Rodríguez 060, Temuco, Chile.

⁶Instituto de Biología, Pontificia Universidad Católica de Valparaíso. Valparaíso 2373223, Chile.

✉ julinota@cepave.edu.ar

Abstract. We report the first record of *Litomosoides pardinasi* in native and exotic rodents from Chile. *Litomosoides pardinasi*, described in the Argentine Patagonia parasitizing *Phyllotis xanthopygus* and *Oligoryzomys longicaudatus*, was found in Chile parasitizing the peritoneal and thoracic cavities of *O. longicaudatus* (prevalence (P)= 18.9%, mean intensity (MI)= 57.3) and *Phyllotis darwini* (P= 4.2%, MI= 10), and in the peritoneal cavity of *Rattus rattus* (P= 1.24%; MI= 22.5). Total P in native rodents from Chile was significantly lower than in Argentina, while the total MI was higher. Prevalence and MI of *L. pardinasi* in *O. longicaudatus* from Chile and Argentina showed non-significant differences; prevalence in *P. darwini* from Chile was significantly lower than in *P. xanthopygus* from Argentina and than in Chilean *O. longicaudatus*. Our results, together with those from Argentina, support the hypothesis that *L. pardinasi* is well established in *O. longicaudatus*, but seems to be recently acquired by *P. darwini* and the exotic *R. rattus*. Considering the known host distribution of *Litomosoides* species among the sigmodontines, our results also support the hypothesis that *L. pardinasi* first colonized the Oryzomyini tribe and later, by different phenomena of host-switching, colonized the Phyllotini tribe and the exotic *R. rattus*.

Key words: host-switching, nematodes, filarioid, sigmodontine rodents, *Oligoryzomys*, *Phyllotis*, *Rattus*.

Resumen. *Litomosoides pardinasi* es registrada por primera vez en roedores nativos y exóticos de Chile. La especie *Litomosoides pardinasi*, parásita de *Phyllotis xanthopygus* y *Oligoryzomys longicaudatus* en la Patagonia Argentina, fue encontrada en Chile parasitando la cavidad torácica y abdominal de *O. longicaudatus* (prevalencia (P)= 18.9%, intensidad media (MI)= 57.3) y de *Phyllotis darwini* (P= 4.2%, MI= 10), y la cavidad abdominal del roedor exótico *Rattus rattus* (P= 1.24%; MI= 22.5). La P total en roedores nativos de Chile fue significativamente menor que la P de Argentina, en tanto la MI total fue significativamente mayor. La P y MI de *L. pardinasi* en *O. longicaudatus* de Chile y Argentina no mostraron diferencias significativas; la P en *P. darwini* de Chile fue significativamente menor que la de *P. xanthopygus* de Argentina y que la de *O. longicaudatus* de Chile. Considerando la distribución de las especies de *Litomosoides* entre los sigmodontinos, estos resultados apoyan la hipótesis que *L. pardinasi* está bien establecida en *O. longicaudatus* y que posiblemente haya sido recientemente adquirida por *P. darwini* y *R. rattus*. Así, *L. pardinasi* colonizó primero a la Tribu Oryzomyini y posteriormente por un fenómeno de cambio de hospedero a la Tribu Phyllotini y a *R. rattus*.

Palabras clave: cambio de hospedero, nemátodos, filarias, roedores sigmodontinos, *Oligoryzomys*, *Phyllotis*, *Rattus*.

Introduction

Worms of the genus *Litomosoides* Chandler, 1931 are filarioid parasites of the thoracic and abdominal cavities of bats, marsupials, and rodents of the families Ctenomyidae, Echimyidae, Sciuridae, and Cricetidae (Esslinger, 1973; Bain et al., 1989; Notarnicola et al., 2002, 2010). They occur in Neotropical and southern Nearctic regions. Most of the *Litomosoides* species occur in South America, including Venezuela (6 species), Colombia (10 species), Peru (3 species), Bolivia (5 species), Argentina (8 species), Brazil (11 species), and Uruguay (1 species) (Travassos, 1919; Esslinger, 1973; Bain et al., 1980, 1989; Brant and Gardner, 1997; Moraes Neto et al., 1997; Notarnicola et al., 2000, 2002; Notarnicola and Navone, 2002, 2011; Guerrero et al., 2002, 2003, 2011).

All the records of *Litomosoides* species have been reported in native host species, with the exception of some specimens of *Litomosoides sigmodontis* Chandler, 1931 found in one individual of the exotic *Rattus norvegicus* Berkenhout, 1769 –originally mentioned as *Mus decumanus*– from 131 specimens trapped in Caracas, Venezuela (Vogel and Gabaldon, 1932).

Among the species of this genus, *Litomosoides pardinasi* Notarnicola and Navone, 2011 is the southernmost worldwide record described in cricetid rodents from Neuquén, Argentina (Notarnicola and Navone, 2011). These authors found *L. pardinasi* in *Phyllotis xanthopygus* Waterhouse, 1837 with a prevalence of 25% and a mean intensity of 7.2, and in *Oligoryzomys longicaudatus* Bennett, 1832 with 33% and 6, respectively.

In Chile, several studies have reported helminths in rodents (e.g., Babero and Cattán, 1975; Durette-Desset et al., 1976; Landaeta-Aqueveque et al., 2007a, 2007b), but none have reported filarioid species. Filarioids in Chile include *Dipetalonema reconditum* Grassi, 1980, *D. dracunculoides* Cobbold, 1870, and *Dirofilaria repens* Railliet and Henry, 1911 in dogs (Alcaíno et al., 1984; López et al., 2012), and *Setaria equina* Abildgaard, 1789 in horses (Alcaíno and Gorman, 1999). Argentina and Chile share several rodent species and some are known to be hosts for this group of nematodes in Argentina.

This study reports the first record of *L. pardinasi* in rodents of Chile. We also report new hosts and the first record of *L. pardinasi* in the exotic rodent *Rattus rattus* Linnaeus, 1758. Finally we discuss the evolution of the host-parasite associations of this nematode.

Materials and methods

We sampled 1 150 native and exotic rodents between the latitudes 31°30'32" S, 35°08'13" S, and between 0 and

1 100 m asl in central Chile, for a total of 19 sampling sites. Localities covered heterogeneous landscapes encompassing several eco-regions, ranging from arid-Mediterranean, to humid-Mediterranean. Rodents were analyzed for the presence of intestinal, peritoneal, and pleural helminths. Trapped native species consisted of 344 *Abrothrix olivaceus* Waterhouse, 1837, 10 *Abrothrix longipilis* Waterhouse, 1837, 58 *O. longicaudatus*, 119 *Phyllotis darwini* Waterhouse, 1837 (Cricetidae), 183 *Octodon degus* Molina, 1782 (Octodontidae), and 7 *Abrocoma bennetti* Waterhouse, 1837 (Abrocomidae), and the exotic rodents were 161 *R. rattus*, 87 *R. norvegicus*, and 181 *Mus musculus* Linnaeus, 1758 (Muridae). Rodents were trapped with Sherman traps baited with rolled oats, and killed using isoflurane anesthesia. Captures followed all ethical guidelines of the Bioethical Committee of the Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile, and were performed with the permission of the Chilean Agriculture and Livestock Bureau (Servicio Agrícola y Ganadero of Chile) and the Chilean National Forest Corporation (Corporación Nacional Forestal).

Worms were recovered directly from the peritoneal and pleural cavities and preserved in 70% ethanol. For examination, samples were cleared with ethanol-glycerin solution and observed under light microscope (Olympus BX51); drawings were made with the aid of a drawing tube. Some worms were studied under Scanning Electron Microscope (SEM Hitachi TM 3000, Pontificia Universidad Católica de Chile, Facultad de Ciencias Biológicas). Measurements are given in micrometers unless otherwise stated; mean values follow the range in parentheses; when two specimens are given, measurements are separated by semicolons.

Prevalence (P) and mean intensity (MI) were calculated following Bush et al. (1997). Fisher's exact test was used to compare prevalences and the negative binomial regression methodology (NBR) was used to analyze intensities. Comparisons with Argentinean P and MI correspond to those reported by Notarnicola and Navone (2011). Statistical analyses were performed in R environment (R Core Team, Vienna, Austria, <http://www.R-project.org/>).

Voucher specimens of worms were deposited at the Museo Nacional de Historia Natural de Chile, Invertebrate Collection (MNHNCL/NEM), and voucher specimen of rodents were deposited at the Museo Nacional de Historia Natural de Chile - Vertebrate Collection (MNHN), and the Patricio Sánchez Reyes Flora and Fauna Collection (UCK) -Pontificia Universidad Católica de Chile.

Results

The filarioids were identified as *L. pardinasi* according to the morphological and morphometric characters stated by Notarnicola and Navone (2011). Specimens show the following characters: the shape of the buccal capsule displays an anterior enlargement in the cuticularized segment (Fig. 1A); the left spicule is constituted with a handle longer than the blade; blade consisting of a filament;




Figure 1. *Litomosoides pardinasi* from *Rattus rattus*. A, buccal capsule lateral view; B, posterior extremity of male showing the spicules and cloacal papillae.

the right spicule is well cuticularized, with a prominent dorsal heel and a terminal cap; the tail is attenuated with one pair of ad-cloacal papillae and three to four pairs of slightly asymmetric postcloacal papillae (Fig. 1B); the arrangement of the head papillae consists of two dorsal cephalic papillae near the amphids and four labial papillae (Fig. 2A); the area rugosa constituted by transverse ridges made of small longitudinal crests (Fig. 2B); and the position of the vulva, posterior to the esophago-intestinal junction. Measurements of the specimens are included in Table 1.

Taxonomic summary

Hosts and localities: *Oligoryzomys longicaudatus* Bennett, 1832 from Los Queñes (35°08'13" S, 70°45'19" W), UCK 423; from Polpaico (33°09' S, 70°53' W), MNHN 1598; from Rinconada de Maipú (33°29' S, 70°49' W), PM84,




Figure 2. Scanning electron microphotographs of *Litomosoides pardinasi*. A, anterior extremity showing the labial papillae (black arrows), cephalic papillae (black star), and amphids (white arrows); B, detail of the area rugosa at mid-length.

Table 1. Measurements of *Litomosoides pardinasi* parasitizing 3 different hosts in Chile

	<i>P. darwini</i>	<i>O. longicaudatus</i>	<i>R. rattus</i>
Females	(n= 2)	(n= 6)	(n= 2)
Body length (mm)	55.4 *	56.1 (36.2-72.2)	53.3 †
Maximum width	-	244 (190-325)	310 †
Width at vulva	186; 207.7	177.1 (160-200)	170.5
Buccal capsule (L x W)	19 x 9; 22 x 9	18 (15-20) x 8 (7-10)	19 x 8; 25 x 12
Esophagus length	415; 533	529 (432-645)	496; 567
Tail length	-	260 (225-312)	855 †
Vulva to apex	772; 961	1 132 (925-1 335)	1 271; 1 559
Males	(n= 3)	(n= 5)	(n= 6)
Body length (mm)	21.5 (20.3-22.7)	15.5 (10.2-18.7)	18.1 (15-22.2)
Maximum width	99 (83-108)	127 (100-140)	111 (105-120)
Buccal capsule (L x W)	19 (16-22) x 8 (6-9)	17 (16-20) x 7 (5-8)	18 (13-20) x 7 (6-9)
Esophagus length	423 (418-427)	405 (350-462)	452 (424-475)
Tail length	139 (99-186)	132 (115-160)	146 (125-170)
Left spicule	199 (180-217)	222 (170-241)	210 (153-232)
Handle	112 (93-130)	132 (115-150)	107 (93-124)
Right spicule	101 (96-108)	105 (87-117)	116 (85-197)
Area rugosa length	1 622 (1 333-2 046)	1 514 (1 350-1 700)	1 467 (1 162-1 725)

*From 1 entire specimen. † One entire female found in this host.

PM183, PM184, PM204, PM207, PM233, PM264, PM266; from Calera de Tango (33°38' S, 70°47' W), PL186. *Phyllotis darwini* Waterhouse, 1837 from Las Chinchillas National Reserve (31°30' S, 71°06' W), MNHN 1599, Dn56, T66, L56, K56. *Rattus rattus* from Fundo El Bosque (33°32' S, 70°48' W), M180; from La Pintana (33°34' S, 70°37' W), M145, Chile.

Site of infection: abdominal and peritoneal cavities.

Specimens deposited: voucher specimens at the Museo Nacional de Historia Natural de Chile, Invertebrates Collection numbers MNHNCL/NEM 11853; 11854; 11855.

Remarks. *Litomosoides pardinasi* was found in the peritoneal and thoracic cavities of 11 *O. longicaudatus* (P= 18.9%; MI= 57.3) and 5 *P. darwini* (P= 4.2%; MI= 10), and in the peritoneal cavity of 2 *R. rattus* (P= 1.24%; MI= 22.5) captured in 7 areas located in Chile. Total prevalence and total mean intensity in native host species were 9% and 42.5, respectively. *Phyllotis darwini* showed significantly lower prevalence than *O. longicaudatus* ($p= 0.003$), and non-significantly higher prevalence than *R. rattus*.

When comparing our data with that from Argentina, total prevalence in Chile, excluding *R. rattus* (9%, n= 177), was significantly lower than in Argentina (25.5%, n= 47) ($p= 0.005$). When comparing prevalence from the same host genus, in Chilean *P. darwini* it was significantly lower (4.2%, n= 119) than in the Argentinean *P. xanthopygus*

(25%, n= 44) ($p < 0.001$), while prevalence in Chilean *O. longicaudatus* (19%, n= 58) was not significantly different from that of the Argentinean *O. longicaudatus* (33.3%, n= 3) ($p= 0.490$). The total mean intensity in Chile excluding *R. rattus* (42.5) was significantly higher than in Argentina (7.08) (Likelihood ratio test (LRt): $p= 0.001$). Analyzing by host species or genus, there were no significant differences between the mean intensities of *O. longicaudatus* from Chile (57.3) and Argentina (6) (LRt: $p= 0.150$); the same was observed between the *Phyllotis* spp. (*P. darwini*: 10 vs. *P. xanthopygus*: 7.2; LRt: $p= 0.560$).

Discussion

Here we report the first record of *L. pardinasi* in Chile and expand the host record to *P. darwini* and *R. rattus*. Identification was based on morphological attributes and supported by morphometric characters. Most of the measurements of Chilean *L. pardinasi* averaged between the minimum and maximum measurements of Argentinean samples (Notarnicola and Navone, 2011). However some differences were observed: female tails of *L. pardinasi* found in *O. longicaudatus* from Chile are shorter than those obtained from Argentina, while the female tail from the specimen found in *R. rattus* is similar to the specimen of *O. longicaudatus* from Argentina (855 versus 800 μm ; Table 1). Male worms from the three host species found in Chile presented shorter tails than those found in Argentina. The

Chilean specimens also had shorter left spicules (153-241 μm versus 210-270 μm). However, the spicular ratio from Chilean specimens fit within the range of the specimens from Argentina (i. e., 1:1.96 in *P. darwini*; 1:2.12 in *O. longicaudatus*, and 1:1.91 in *R. rattus* from Chile versus a range of 1:1.8 to 1:2.4 in *P. xanthopygus* and 1:2.09 in *O. longicaudatus* from Argentina). Morphometric variability was reported within other *Litomosoides* species and these observations were attributed to differences in the host species and/or geographic distribution (Esslinger, 1973; Bain et al., 1989; Guerrero et al., 2002; Notarnicola 2005; Notarnicola et al., 2010).

Prevalences and mean intensities of *L. pardinasi* in *O. longicaudatus* from Chile and Argentina showed non-significant differences, while in *Phyllotis* spp. the prevalence from the Chilean specimens was significantly lower than in Argentina. Thus, it appears that *L. pardinasi* is well established in *O. longicaudatus* based on the similarities in their prevalences and mean intensities in Chile and Argentina. In contrast, this filarioid species seems to be recently acquired by *P. darwini* and the exotic *R. rattus* in Chile, based on the significantly lower prevalence, suggesting that this parasitism is a result of a recent host-switching phenomenon. Moreover, of the 16 *Litomosoides* species registered in Sigmodontinae rodents, 6 are present in Akodontini, 9 in Oryzomyini –including *L. pardinasi*–, and 1 in both Akodontini and Oryzomyini. *Litomosoides pardinasi* is the unique species registered in Phyllotini. This result supports the hypothesis that *L. pardinasi* primarily colonized the Oryzomyini tribe and later, by different phenomena of host-switching, colonized the Phyllotini (*P. xanthopygus* in Argentina and *P. darwini* in Chile) and the exotic host *R. rattus*. The host-switching phenomenon was frequently reported between native rodents living in sympatry (Guerrero et al., 2002; Notarnicola, 2005; Notarnicola and Navone, 2011).

Vogel and Gabaldon (1932) reported the presence of adults and microfilariae of *L. sigmodontis* in *R. norvegicus* from Caracas, Venezuela. This filarioid species was also found to parasitize the cotton rat *Sigmodon hispidus* Say and Ord, 1825 from Mexico and USA (Ochoterena and Caballero, 1932; Forrester and Kinsella, 1973). Female specimens of *L. pardinasi* found in *R. rattus* presented microfilariae in the uterus, indicating that its life cycle can be completed in the host and can be transmitted to other hosts. All these results confirm that filarioid species are able to colonize allochthonous rodents when they share microhabitats with parasitized native rodents.

The mite *Ornithonyssus bacoti* Hirst, 1913 (Acari, Macronyssidae) has been reported as a vector for several species of *Litomosoides* in rodents as well as in bats (Bain et al., 1980, 2002). Another study showed

that *Hoplopleura travassosi* Werneck, 1932 (Anoplura, Hoplopleuridae) presented positive association with *Litomosoides bonaerensis* Notarnicola, Bain and Navone 2000 (Lareschi et al., 2003). In Chile, *O. bacoti* has been recorded in several native and allochthonous rodents (Jofré et al., 2009; Peña Oyarce, 2009); whereas *H. travassosi* and *Hoplopleura aitkeni* Johnson, 1972 have been found in *O. longicaudatus* and *P. darwini*, respectively, in a geographic range that encompasses this work (González-Acuña et al., 2003; 2005). The presence of these arthropods in native and allochthonous rodents provide more information to the hypothesis that filarioids are transmitted by these vectors in rodents living in sympatry.

Acknowledgments

The authors wish to thank Maria Cristina Estivariz from CEPAVE for the drawings and Eileen Smith for editorial support. This work was supported by the Conicyt Ph. D. scholarships awarded to C.L. (number 24110058) and J.P.C. (AT-24100028), and Fondecyt grants awarded to P.E.C (1070960), C.B. (11090086) and F.T.P. (1110664). JN is a member of Conicet.

Literature cited

- Alcaíno, H. and T. R. Gorman. 1999. Parásitos de los animales domésticos en Chile. *Parasitología al Día* 23:33-41.
- Alcaíno, H., T. R. Gorman and M. Puelma. 1984. Filariasis canina en Chile. *Archivos de Medicina Veterinaria* 16:67-73.
- Babero, B. B. and P. E. Cattán. 1975. Helminthofauna de Chile: III. Parásitos del roedor degú, *Octodon degus* Molina, 1782, con la descripción de tres nuevas especies. *Boletín Chileno de Parasitología* 30:68-76.
- Bain, O., S. Babayan, J. Gomes, G. Rojas and R. Guerrero. 2002. First account on the larval biology of a *Litomosoides* filaria from a bat. *Parassitologia* 44:89-92.
- Bain, O., G. Petit and S. Berteaux. 1980. Description de deux nouvelles filaires du genre *Litomosoides* et de leurs stades infestants. *Annales de Parasitologie Humaine et Comparée* 55:225-267.
- Bain, O., G. Petit and M. Diagne. 1989. Étude de quelques *Litomosoides* parasites de rongeurs: Conséquences taxonomiques. *Annales de Parasitologie Humaine et Comparée* 64:268-289.
- Brant, S. V. and S. L. Gardner. 1997. Two new species of *Litomosoides* (Nematoda: Onchocercidae) from *Ctenomys opimus* (Rodentia: Ctenomyidae) on the Altiplano of Bolivia. *Journal of Parasitology* 83:700-705.
- Bush, A. O., K. D. Lafferty, J. M. Lotz and A. W. Shostak. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83:575-583.
- Durette-Desset, M. C., M. A. Denké and R. Murúa. 1976. Présence chez un Rongeur du Chili d'un Nématode Inglamidinae (sub.

- Fam. Nov) appartenant aux Amidostomatidae, famille connue des Mammifères d'Australie. *Annales de Parasitologie Humaine et Comparée* 51:453-460.
- Esslinger, H. 1973. The genus *Litomosoides* Chandler, 1931 (Filarioidea: Onchocercidae) in Colombian bats and rats. *Journal of Parasitology* 59:225-246.
- Forrester, D. F. and J. M. Kinsella. 1973. Comparative morphology and ecology of two species of *Litomosoides* (Nematoda: Filarioidea) of rodents in Florida, with a key to the species of *Litomosoides* Chandler, 1931. *International Journal for Parasitology* 3:255-263.
- González-Acuña, D., D. D. C. Castro and L. Moreno-Salas. 2003. Contribución al conocimiento de los Phthiraptera (Anoplura, Hoplopleura) parásitos de roedores en Chile. *Gayana* 67:117-119.
- Guerrero, R., O. Bain, C. Martin and M. Barbutto. 2011. A new species of *Litomosoides* (Nematoda: Onchocercidae), parasite of *Nectomys palmipes* (Rodentia: Cricetidae: Sigmodontinae) from Venezuela: description, molecular evidence, *Wolbachia pipientis* screening. *Folia Parasitologica* 58:149-156.
- González-Acuña, D., D. D. C. Castro, L. Moreno-Salas, J. Torres-Mura and E. Mey. 2005. New records of sucking lice (Insecta: Phthiraptera: Anoplura) on rodents (Mammalia; Rodentia; Muridae) from Chile. *Mastozoología Neotropical* 12:249-251.
- Guerrero, R., C. Martin and O. Bain. 2003. *Litomosoides yutajensis* n. sp., first record of this filarial genus in a mormoopid bat. *Parasite* 10:219-25.
- Guerrero, R., C. Martin, S. L. Gardner and O. Bain. 2002. New and known species of *Litomosoides* (Nematoda: Filarioidea): Important adult and larval characters and taxonomic changes. *Comparative Parasitology* 69:177-195.
- Jofré, L., I. Noemí, P. Neira, T. Saavedra and C. Diaz. 2009. Animal mites transmissible to humans and associated zoonosis. *Revista Chilena de Infectología* 26:248-257
- Landaeta-Aqueveque, C. A., M. D. R. Robles and P. E. Cattán. 2007a. The community of gastrointestinal helminths in the house mouse, *Mus musculus*, in Santiago, Chile. *Parasitología Latinoamericana* 62:165-169.
- Landaeta-Aqueveque, C. A., M. D. R. Robles and P. E. Cattán. 2007b. Helminthofauna del roedor *Abrothrix olivaceus* (Sigmodontinae) en áreas sub-urbanas de Santiago de Chile. *Parasitología Latinoamericana* 62:134-141.
- Lareschi, M., J. Notarnicola, G. Navone and P. M. Linardi. 2003. Arthropod and filarioid parasites associated with wild rodents in the northeast marshes of Buenos Aires, Argentina. *Memórias do Instituto Oswaldo Cruz* 98:673-677.
- López, J., F. Valiente-Echeverría, M. Carrasco, R. Mercado and K. Abarca. 2012. Identificación morfológica y molecular de filarias caninas en una comuna semi-rural de la Región Metropolitana, Chile. *Revista Chilena Infectología* 29:284-289.
- Moraes-Neto, A. H. A., R. M. Lanfredi and W. De Souza. 1997. *Litomosoides chagasfilhoi* sp. nov. (Nematoda: Filarioidea) parasitizing the abdominal cavity of *Akodon cursor* (Winge, 1887) Rodentia: Muridae from Brazil. *Parasitology Research* 83:137-143.
- Notarnicola, J. 2005. Description of adult and fourth-stage larva of *Litomosoides navonae* n. sp. (Nematoda: Onchocercidae), a parasite of five species of sigmodontine rodents from northeastern Argentina. *Systematic Parasitology* 62:171-183.
- Notarnicola, J., O. Bain and G. T. Navone. 2000. Two new species of *Litomosoides* (Nematoda: Filarioidea) in Sigmodontines (Rodentia: Muridae) from Río de La Plata marshland, Argentina. *Journal of Parasitology* 86:1318-1325.
- Notarnicola, J., O. Bain and G. T. Navone. 2002. *Litomosoides anguyai* n. sp. (Nematoda: Onchocercidae) from *Oxymycterus misionalis* (Rodentia: Muridae) in the rain forest of Misiones, Argentina. *Systematic Parasitology* 52:129-135.
- Notarnicola, J., F. A. Jiménez Ruiz and S. L. Gardner. 2010. *Litomosoides* (Nematoda: Filarioidea) of bats from Bolivia with records for three known species and the description of a new species. *Journal of Parasitology* 96:775-782.
- Notarnicola, J. and G. Navone. 2002. A new species, *Litomosoides odilae* n. sp. (Nematoda: Onchocercidae) from *Oligoryzomys nigripes* (Rodentia: Muridae) in the rainforest of Misiones, Argentina. *Journal of Parasitology* 88:967-971.
- Notarnicola, J. and G. Navone. 2011. *Litomosoides pardinasi* n. sp. (Nematoda: Onchocercidae) from two species of cricetid rodents in Northern Patagonia, Argentina. *Parasitology Research* 108:187-194.
- Ochoterena I. and E. Caballero. 1932. Filaria parasita de las ratas de campo, *Micropleura sigmodoni* spec. nov. *Anales del Instituto de Biología. Universidad Nacional de Mexico* 3:123-125.
- Peña-Oyarce, L. A. 2009. Análisis de la presencia de parasitosis zoonóticas en roedores que cohabitan con la población humana en diferentes sectores de las comunas de Valdivia y San José de la Mariquina. Tesis, Universidad Austral de Chile. Chile, Valdivia. 43 p.
- Travassos, L. 1919. *Filaria carinii* n. sp. *Revista de la Sociedade Brasileira de Ciências* 3:189-190.
- Vogel, H. and A. Gabaldon. 1932. *Vestibulosestaria*, eine neue Filariengattung aus Rattenarten. *Zentralblatt für Bakteriologie, Parasitenkunde und Infektionskrankheiten* 105:119-124.