



ELSEVIER

Available online at www.sciencedirect.com

Physics Letters B 557 (2003) 147–156

PHYSICS LETTERS B

www.elsevier.com/locate/npe

Measurement of W polarisation at LEP

L3 Collaboration

P. Achard^t, O. Adriani^q, M. Aguilar-Benitez^x, J. Alcaraz^{x,r}, G. Alemanni^v, J. Allaby^r, A. Aloisio^{ab}, M.G. Alviggi^{ab}, H. Anderhub^{at}, V.P. Andreev^{f,ag}, F. Anselmo^h, A. Arefiev^{aa}, T. Azemoon^c, T. Aziz^{i,r}, P. Bagnaia^{al}, A. Bajo^x, G. Baksay^y, L. Baksay^y, S.V. Baldew^b, S. Banerjeeⁱ, Sw. Banerjee^d, A. Barczyk^{at,ar}, R. Barillère^r, P. Bartalini^v, M. Basile^h, N. Batalova^{aq}, R. Battiston^{af}, A. Bay^v, F. Becattini^q, U. Becker^m, F. Behner^{at}, L. Bellucci^q, R. Berbeco^c, J. Berdugo^x, P. Berges^m, B. Bertucci^{af}, B.L. Betev^{at}, M. Biasini^{af}, M. Biglietti^{ab}, A. Biland^{at}, J.J. Blaising^d, S.C. Blyth^{ah}, G.J. Bobbink^b, A. Böhm^a, L. Boldizsar^l, B. Borgia^{al}, S. Bottai^q, D. Bourilkov^{at}, M. Bourquin^t, S. Braccini^t, J.G. Branson^{an}, F. Brochu^d, J.D. Burger^m, W.J. Burger^{af}, X.D. Cai^m, M. Capell^m, G. Cara Romeo^h, G. Carlino^{ab}, A. Cartacci^q, J. Casaus^x, F. Cavallari^{al}, N. Cavallo^{ai}, C. Cecchi^{af}, M. Cerrada^x, M. Chamizo^t, Y.H. Chang^{av}, M. Chemarin^w, A. Chen^{av}, G. Chen^g, G.M. Chen^g, H.F. Chen^u, H.S. Chen^g, G. Chiefari^{ab}, L. Cifarelli^{am}, F. Cindolo^h, I. Clare^m, R. Clare^{ak}, G. Coignet^d, N. Colino^x, S. Costantini^{al}, B. de la Cruz^x, S. Cucciarelli^{af}, J.A. van Dalen^{ad}, R. de Asmundis^{ab}, P. Déglon^t, J. Debreczeni^l, A. Degré^d, K. Dehmelt^y, K. Deiters^{ar}, D. della Volpe^{ab}, E. Delmeire^t, P. Denes^{aj}, F. DeNotaristefani^{al}, A. De Salvo^{at}, M. Diemoz^{al}, M. Dierckxsens^b, C. Dionisi^{al}, M. Dittmar^{at,r}, A. Doria^{ab}, M.T. Dova^{j,5}, D. Duchesneau^d, M. Duda^a, B. Echenard^t, A. Eline^r, A. El Hage^a, H. El Mamouni^w, A. Engler^{ah}, F.J. Eppling^m, P. Extermann^t, M.A. Falagan^x, S. Falciano^{al}, A. Favara^{ae}, J. Fay^w, O. Fedin^{ag}, M. Felcini^{at}, T. Ferguson^{ah}, H. Fesefeldt^a, E. Fiandrini^{af}, J.H. Field^t, F. Filthaut^{ad}, P.H. Fisher^m, W. Fisher^{aj}, I. Fisk^{an}, G. Forconi^m, K. Freudenreich^{at}, C. Furetta^z, Yu. Galaktionov^{aa,m}, S.N. Ganguliⁱ, P. Garcia-Abia^{e,r}, M. Gataullin^{ae}, S. Gentile^{al}, S. Giagu^{al}, Z.F. Gong^u, G. Grenier^w, O. Grimm^{at}, M.W. Gruenewald^p, M. Guida^{am}, R. van Gulik^b, V.K. Gupta^{aj}, A. Gurtuⁱ, L.J. Gutay^{aq}, D. Haas^e, R.Sh. Hakobyan^{ad}, D. Hatzifotiadou^h, T. Hebbeker^a, A. Hervé^r, J. Hirschfelder^{ah}, H. Hofer^{at}, M. Hohlmann^y, G. Holzner^{at}, S.R. Hou^{av}, Y. Hu^{ad}, B.N. Jin^g, L.W. Jones^c, P. de Jong^b, I. Josa-Mutuberría^x, D. Käfer^a, M. Kaurⁿ, M.N. Kienzle-Focacci^t, J.K. Kim^{ap}, J. Kirkby^r, W. Kittel^{ad}, A. Klimentov^{m,aa}, A.C. König^{ad}, M. Kopal^{aq}, V. Koutsenko^{m,aa}, M. Kräber^{at}, R.W. Kraemer^{ah},

A. Krüger^{as}, A. Kunin^m, P. Ladron de Guevara^x, I. Laktineh^w, G. Landi^q, M. Lebeau^r,
 A. Lebedev^m, P. Lebrun^w, P. Lecomte^{at}, P. Lecoq^r, P. Le Coultre^{at}, J.M. Le Goff^r,
 R. Leiste^{as}, M. Levtchenko^z, P. Levtchenko^{ag}, C. Li^u, S. Likhoded^{as}, C.H. Lin^{av},
 W.T. Lin^{av}, F.L. Linde^b, L. Lista^{ab}, Z.A. Liu^g, W. Lohmann^{as}, E. Longo^{al}, Y.S. Lu^g,
 C. Luci^{al}, L. Luminari^{al}, W. Luster^{at}, W.G. Ma^u, L. Malgeri^t, A. Malinin^{aa},
 C. Maña^x, D. Mangeol^{ad}, J. Mans^{aj}, J.P. Martin^w, F. Marzano^{al}, K. Mazumdarⁱ,
 R.R. McNeil^f, S. Mele^{r,ab}, L. Merola^{ab}, M. Meschini^q, W.J. Metzger^{ad}, A. Mihul^k,
 H. Milcent^r, G. Mirabelli^{al}, J. Mnich^a, G.B. Mohantyⁱ, G.S. Muanza^w, A.J.M. Muijs^b,
 B. Musicar^{an}, M. Musy^{al}, S. Nagy^o, S. Natale^t, M. Napolitano^{ab}, F. Nessi-Tedaldi^{at},
 H. Newman^{ae}, A. Nisati^{al}, H. Nowak^{as}, R. Ofierzynski^{at}, G. Organtini^{al}, C. Palomares^r,
 P. Paolucci^{ab}, R. Paramatti^{al}, G. Passaleva^q, S. Patricelli^{ab}, T. Paul^j, M. Pauluzzi^{af},
 C. Paus^m, F. Pauss^{at}, M. Pedace^{al}, S. Pensotti^z, D. Perret-Gallix^d, B. Petersen^{ad},
 D. Piccolo^{ab}, F. Pierella^h, M. Pioppi^{af}, P.A. Piroué^{aj}, E. Pistolesi^z, V. Plyaskin^{aa},
 M. Pohl^t, V. Pojidaev^q, J. Pothier^r, D.O. Prokofiev^{aq}, D. Prokofiev^{ag}, J. Quartieri^{am},
 G. Rahal-Callot^{at}, M.A. Rahamanⁱ, P. Raics^o, N. Rajaⁱ, R. Ramelli^{at}, P.G. Rancoita^z,
 R. Ranieri^q, A. Raspereza^{as}, P. Razis^{ac}, D. Ren^{at}, M. Rescigno^{al}, S. Reucroft^j,
 S. Riemann^{as}, K. Riles^c, B.P. Roe^c, L. Romero^x, A. Rosca^{as}, S. Rosier-Lees^d,
 S. Roth^a, C. Rosenbleck^a, B. Roux^{ad}, J.A. Rubio^r, G. Ruggiero^q, H. Rykaczewski^{at},
 A. Sakharov^{at}, S. Saremi^f, S. Sarkar^{al}, J. Salicio^r, E. Sanchez^x, M.P. Sanders^{ad},
 C. Schäfer^r, V. Schegelsky^{ag}, H. Schopper^{au}, D.J. Schotanus^{ad}, C. Sciacca^{ab},
 L. Servoli^{af}, S. Shevchenko^{ae}, N. Shivarov^{ao}, V. Shoutko^m, E. Shumilov^{aa},
 A. Shvorob^{ae}, D. Son^{ap}, C. Souga^w, P. Spillantini^q, M. Steuer^m, D.P. Stickland^{aj},
 B. Stoyanov^{ao}, A. Straessner^r, K. Sudhakarⁱ, G. Sultanov^{ao}, L.Z. Sun^u, S. Sushkov^a,
 H. Suter^{at}, J.D. Swain^j, Z. Szillasi^{y,3}, X.W. Tang^g, P. Tarjan^o, L. Tauscher^e, L. Taylor^j,
 B. Tellili^w, D. Teyssier^w, C. Timmermans^{ad}, Samuel C.C. Ting^m, S.M. Ting^m,
 S.C. Tonwar^{i,r}, J. Tóth^l, C. Tully^{aj}, K.L. Tung^g, J. Ulbricht^{at}, E. Valente^{al},
 R.T. Van de Walle^{ad}, R. Vasquez^{aq}, V. Veszpremi^y, G. Vesztergombi^l, I. Vetlitsky^{aa},
 D. Vicinanza^{am}, G. Viertel^{at}, S. Villa^{ak}, M. Vivargent^d, S. Vlachos^e, I. Vodopianov^y,
 H. Vogel^{ah}, H. Vogt^{as}, I. Vorobiev^{ah,aa}, A.A. Vorobyov^{ag}, M. Wadhwa^e, X.L. Wang^u,
 Z.M. Wang^u, M. Weber^a, P. Wienemann^a, H. Wilkens^{ad}, S. Wynhoff^{aj}, L. Xia^{ae},
 Z.Z. Xu^u, J. Yamamoto^c, B.Z. Yang^u, C.G. Yang^g, H.J. Yang^c, M. Yang^g, S.C. Yeh^{aw},
 An. Zalite^{ag}, Yu. Zalite^{ag}, Z.P. Zhang^u, J. Zhao^u, G.Y. Zhu^g, R.Y. Zhu^{ae},
 H.L. Zhuang^g, A. Zichichi^{h,r,s}, B. Zimmermann^{at}, M. Zöllner^a

^a III. Physikalisches Institut, RWTH, D-52056 Aachen, Germany¹

^b National Institute for High Energy Physics, NIKHEF, and University of Amsterdam, NL-1009 DB Amsterdam, The Netherlands

^c University of Michigan, Ann Arbor, MI 48109, USA

^d Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux cedex, France

^e Institute of Physics, University of Basel, CH-4056 Basel, Switzerland

^f Louisiana State University, Baton Rouge, LA 70803, USA

^g Institute of High Energy Physics, IHEP, 100039 Beijing, China⁶

- ^h University of Bologna and INFN-Sezione di Bologna, I-40126 Bologna, Italy
ⁱ Tata Institute of Fundamental Research, Mumbai (Bombay) 400 005, India
^j Northeastern University, Boston, MA 02115, USA
^k Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania
^l Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary²
^m Massachusetts Institute of Technology, Cambridge, MA 02139, USA
ⁿ Panjab University, Chandigarh 160 014, India
^o KLTE-ATOMKI, H-4010 Debrecen, Hungary³
^p Department of Experimental Physics, University College Dublin, Belfield, Dublin 4, Ireland
^q INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy
^r European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland
^s World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland
^t University of Geneva, CH-1211 Geneva 4, Switzerland
^u Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China⁶
^v University of Lausanne, CH-1015 Lausanne, Switzerland
^w Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne, France
^x Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain⁴
^y Florida Institute of Technology, Melbourne, FL 32901, USA
^z INFN-Sezione di Milano, I-20133 Milan, Italy
^{aa} Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia
^{ab} INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy
^{ac} Department of Physics, University of Cyprus, Nicosia, Cyprus
^{ad} University of Nijmegen and NIKHEF, NL-6525 ED Nijmegen, The Netherlands
^{ae} California Institute of Technology, Pasadena, CA 91125, USA
^{af} INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy
^{ag} Nuclear Physics Institute, St. Petersburg, Russia
^{ah} Carnegie Mellon University, Pittsburgh, PA 15213, USA
^{ai} INFN-Sezione di Napoli and University of Potenza, I-85100 Potenza, Italy
^{aj} Princeton University, Princeton, NJ 08544, USA
^{ak} University of California, Riverside, CA 92521, USA
^{al} INFN-Sezione di Roma and University of Rome “La Sapienza”, I-00185 Rome, Italy
^{am} University and INFN, Salerno, I-84100 Salerno, Italy
^{an} University of California, San Diego, CA 92093, USA
^{ao} Bulgarian Academy of Sciences, Central Lab. of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria
^{ap} The Center for High Energy Physics, Kyungpook National University, 702-701 Taegu, South Korea
^{aq} Purdue University, West Lafayette, IN 47907, USA
^{ar} Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland
^{as} DESY, D-15738 Zeuthen, Germany
^{at} Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland
^{au} University of Hamburg, D-22761 Hamburg, Germany
^{av} National Central University, Chung-Li, Taiwan, ROC
^{aw} Department of Physics, National Tsing Hua University, Taiwan, ROC

Received 16 December 2002; received in revised form 27 January 2003; accepted 31 January 2003

Editor: L. Rolandi

Abstract

The three different helicity states of W bosons produced in the reaction $e^+e^- \rightarrow W^+W^- \rightarrow \ell\nu q\bar{q}'$ at LEP are studied using leptonic and hadronic W decays. Data at centre-of-mass energies $\sqrt{s} = 183\text{--}209$ GeV are used to measure the polarisation of W bosons, and its dependence on the W boson production angle. The fraction of longitudinally polarised W bosons is measured to be $0.218 \pm 0.027 \pm 0.016$ where the first uncertainty is statistical and the second systematic, in agreement with the Standard Model expectation.

© 2003 Published by Elsevier Science B.V. Open access under [CC BY license](https://creativecommons.org/licenses/by/4.0/).

1. Introduction

The existence of all three W boson helicity states, $+1$, -1 and 0 , is a consequence of the non-vanishing mass of the W boson, that, in the Standard Model [1], is generated by the Higgs mechanism of electroweak symmetry breaking. The measurement of the fractions of longitudinally and transversely polarised W bosons constitutes a test of the Standard Model predictions for the triple gauge boson couplings γWW and ZWW .

To determine the W helicity fractions, events of the type $e^+e^- \rightarrow W^+W^- \rightarrow \ell\nu q\bar{q}'$ are used, with ℓ denoting either an electron or a muon. These events are essentially background free and allow a measurement, with good accuracy, of the W momentum vector, the W charge and the polar decay angles. The W helicity states are accessible in a model independent way through the shape of the distributions of the polar decay angle, θ_ℓ^* , between the charged lepton and the W direction in the W rest frame. Transversely polarised W bosons have angular distributions $(1 \mp \cos\theta_\ell^*)^2$ for a W^- with helicity ± 1 , and $(1 \pm \cos\theta_\ell^*)^2$ for a W^+ with helicity ± 1 . For longitudinally polarised W bosons, a $\sin^2\theta_\ell^*$ dependence is expected. For simplicity, we refer in the following only to the fractions f_- , f_+ and f_0 of the helicity states -1 , $+1$ and 0 of the W^- boson, respectively. Assuming CP invariance these equal the fractions of the corresponding helicity states $+1$, -1 and 0 of the W^+ boson.

The differential distribution of leptonic W^- decays at Born level is

$$\frac{1}{N} \frac{dN}{d\cos\theta_\ell^*} = f_- \frac{3}{8} (1 + \cos\theta_\ell^*)^2 + f_+ \frac{3}{8} (1 - \cos\theta_\ell^*)^2 + f_0 \frac{3}{4} \sin^2\theta_\ell^*. \quad (1)$$

¹ Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie.

² Supported by the Hungarian OTKA fund under contract numbers T019181, F023259 and T037350.

³ Also supported by the Hungarian OTKA fund under contract number T026178.

⁴ Supported also by the Comisión Interministerial de Ciencia y Tecnología.

⁵ Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.

⁶ Supported by the National Natural Science Foundation of China.

For hadronic W decays, the quark charge is difficult to reconstruct experimentally and only the absolute value of the cosine of the decay angle, $|\cos\theta_q^*|$, is used

$$\frac{1}{N} \frac{dN}{d|\cos\theta_q^*|} = f_\pm \frac{3}{4} (1 + |\cos\theta_q^*|^2) + f_0 \frac{3}{2} (1 - |\cos\theta_q^*|^2), \quad (2)$$

with $f_\pm = f_+ + f_-$.

After correcting the data for selection efficiencies and background, the different fractions of W helicity states are obtained from a fit to these distributions. The fractions f_- , f_+ and f_0 are also determined as a function of the W^- production angle Θ_{W^-} in the laboratory frame. The helicity composition of the W bosons depends strongly on the centre-of-mass energy, \sqrt{s} .

2. Data and Monte Carlo

The analysis presented in this Letter is based on the whole data set collected with the L3 detector [2] and supersedes our previous results [3] based on about one third of the data. An integrated luminosity of 684.8 pb^{-1} , collected at different centre-of-mass energies between 183 and 209 GeV, as shown in Table 1, is analysed.

The $e^+e^- \rightarrow W^+W^- \rightarrow \ell\nu q\bar{q}'$, $\mu\nu q\bar{q}'$ Monte Carlo events are generated using KORALW [4]. The Standard Model predictions for f_- , f_+ and f_0 are obtained from these samples by fitting the generated decay angular distributions for each value of \sqrt{s} . As an example, the expected fraction of longitudinally polarised W bosons changes from 0.271 at $\sqrt{s} = 183 \text{ GeV}$ to 0.223 at $\sqrt{s} = 206 \text{ GeV}$. The luminosity averaged Standard Model expectations for f_- , f_+ and f_0 are 0.590, 0.169 and 0.241, respectively.

Background processes are generated using KORALW for W pair production decaying to other final states, and PYTHIA [5] and KK2F [6] for $e^+e^- \rightarrow q\bar{q}'(\gamma)$. For studies of systematic effects, signal events are also generated using EEW [7] and EXCALIBUR [8]. The L3 detector response is simulated with the GEANT [9] and GEISHA [10] packages. Detector inefficiencies, as monitored during the data taking period, are included.

Table 1

Average centre-of-mass energies, integrated luminosities and numbers of selected events

$\langle\sqrt{s}\rangle$ [GeV]	182.7	188.6	191.6	195.5	199.5	201.8	205.9
Integrated luminosity [pb^{-1}]	55.5	176.8	29.8	84.1	83.3	37.2	218.1
Selected $e\nu q\bar{q}'$ events	82	293	59	133	110	56	355
Selected $\mu\nu q\bar{q}'$ events	67	255	43	110	99	59	289

A large sample of signal events is generated using the EEWW Monte Carlo program. This program assigns, differently from KORALW, W helicities on an event-by-event basis but uses the zero-width approximation for the W boson and does not include higher order radiative corrections and interference terms. The W^- helicity fractions obtained from a fit to the generated decay angle distributions agree with the input values. A comparison of the fractions obtained from EEWW and YFSWW [11], which includes improved $O(\alpha)$ corrections, with those obtained from KORALW also shows good agreement. Therefore the Born level formulae (1) and (2) are applicable after radiative corrections.

3. Selection of $W^+W^- \rightarrow e\nu q\bar{q}'$, $\mu\nu q\bar{q}'$ events

Only events which contain exactly one electron or one muon candidate are accepted [3]. Electrons are identified as isolated energy depositions in the electromagnetic calorimeter with an electromagnetic shower shape. A match in azimuthal angle with a track reconstructed in the central tracking chamber is required. Muons are identified and measured as tracks reconstructed in the muon chambers which point back to the interaction vertex. All other energy depositions in the calorimeters are assumed to originate from the hadronically decaying W. The neutrino momentum vector is assumed to be the missing momentum vector of the event. The following additional criteria are applied:

- the reconstructed momentum must be greater than 20 GeV for electrons and 15 GeV for muons;
- the neutrino momentum must be greater than 10 GeV and its polar angle, θ_ν , has to satisfy $|\cos\theta_\nu| < 0.95$;
- the invariant mass of the lepton–neutrino system has to be greater than 60 GeV;

- the invariant mass of the hadronic system has to be between 50 and 110 GeV.

Fig. 1 shows some distributions of those variables for data and Monte Carlo.

The number of events selected by these criteria are listed in Table 1. In total, 2010 events are selected with an efficiency of 65.7% and a purity of 96.3%. The contamination from $W^+W^- \rightarrow \tau\nu q\bar{q}'$ and $e^+e^- \rightarrow q\bar{q}'(\gamma)$ is 2.4% and 1.3%, respectively, independent of \sqrt{s} and the W production angle.

4. Analysis of the W helicity states

For the selected events, the rest frames of the W bosons are calculated from the lepton and neutrino momenta, and the decay angles θ_ℓ^* and θ_q^* of the lepton and the quarks are determined. The angle θ_q^* is approximated by the polar angle of the thrust axis with respect to the W direction in the rest frame of the hadronically decaying W.

The fractions of the W helicity states are obtained from the event distributions, $dN/d\cos\theta_\ell^*$ and $dN/d|\cos\theta_q^*|$. For each energy point, the background, as obtained from Monte Carlo simulations, is subtracted from the data, and the resulting distributions are corrected for selection efficiencies as obtained from large samples of KORALW Monte Carlo events. The corrected decay angle distributions at the different centre-of-mass energies are combined into single distributions for leptonic and hadronic decays, which are then fitted to the functions (1) and (2), respectively. A binned fit is performed on the normalised distributions, shown in Fig. 2, using f_- and f_0 as the fit parameters. The fraction f_+ is obtained by constraining the sum of all three parameters to unity.

Detector resolution introduces migration effects that bias the fitted parameters. For example, purely longitudinally polarised leptonically decaying W bo-




Fig. 1. Distributions used for the event selections: (a) momentum of the electrons, (b) momentum of the muons, (c) absolute value of the cosine of the polar angle of the neutrino, (d) mass of the lepton–neutrino system. In each plot, all other selection criteria are applied. The arrows indicate cut positions.

sons at $\sqrt{s} = 206$ GeV would be measured to have a helicity composition: $f_0 = 0.945$, $f_- = 0.043$ and $f_+ = 0.012$. The magnitude of these effects depends on the helicity fractions and on \sqrt{s} . Corrections for this bias as a function of the helicity fractions are determined from EEW Monte Carlo samples. If the ratio of two helicity fractions is constant the bias correction function of the third fraction is linear to a good approximation. For the correction of f_0 in the hadronic W decay, the ratio f_-/f_+ is taken from the measurement in the leptonic W decay, as only the sum

of f_+ and f_- is known from hadronic decays. Bias correction functions are determined for the analysis of the complete data sample, separately for the W^+ and W^- events and in bins of the W^- production angle.

5. Results

The results of the fits to the decay angle distributions for leptonic and hadronic W decays are shown in Fig. 2. The data are well described only if all three




Fig. 2. Corrected decay angle distributions for (a) leptonic W decays and (b) for hadronic W decays at $\sqrt{s} = 183\text{--}209$ GeV. Fit results for the different W helicity hypotheses are also shown.

W helicity states are used. Fits omitting the helicity 0 state fail to describe the data. For leptonic W decays, the χ^2 increases from 12.7 for eight degrees of freedom if all helicity states are included to 56.2 for nine degrees of freedom if only the helicities +1 and -1 are used in the fit. For hadronic W decays, the χ^2 increases from 6.6 for four degrees of freedom if all helicity states are included to 59.1 for four degrees of freedom if only the helicities ± 1 are used.

The measured fractions of the W helicity states in data, at an average centre-of-mass energy $\sqrt{s} = 196.7$ GeV, are presented together with the Standard Model expectation in Tables 2, 3 and 4. The parameters f_- and f_0 derived from the fit are about 90% anti-correlated. These results include a bias correction of 0.005 on f_0 for leptonic decays and 0.044 for hadronic decays. The bias correction adds 0.003 to the statistical uncertainty on f_0 for leptonic decays and 0.007 to the one for hadronic decays. The measured W helicity fractions agree with the Standard Model expectations for the leptonic and hadronic decays, as well as for the combined sample. Longitudinal W polarisation is observed with a significance of seven standard deviations, including systematic uncertainties.

A number of systematic uncertainties are considered. These include selection criteria, binning effects, bias corrections, the contamination due to non-double resonant four fermion processes, background levels, and efficiencies. Selection cuts are varied over a range of one standard deviation of the corresponding reconstruction accuracy. Fits are repeated with one bin more or one bin less in the decay angle distributions. Uncertainties on the bias and efficiency corrections are determined with large Monte Carlo samples, the latter being negligible. The contamination due to non-double resonant four fermion processes is studied by using the EXCALIBUR Monte Carlo. Background levels are varied according to Monte Carlo statistics for both the $e^+e^- \rightarrow W^+W^- \rightarrow \tau\nu q\bar{q}'$ and $e^+e^- \rightarrow q\bar{q}'(\gamma)$ processes. The largest uncertainties arise from selection criteria and binning effects. As an example, Table 5 summarises those effects on f_0 .

Within the Standard Model, CP symmetry is conserved in the reaction $e^+e^- \rightarrow W^+W^-$ and the helicity fractions f_+ , f_- and f_0 for the W^+ are expected to be identical to the fractions f_- , f_+ and f_0 , for the W^- , respectively. CP invariance is tested by measuring the helicity fractions for W^+ and W^- separately. The charge of the W bosons is obtained from the charge of the lepton. We select 1020 $W^+ \rightarrow \ell^+\nu$, and 990 $W^- \rightarrow \ell^-\bar{\nu}$ events. Results of separate fits for the W^- helicity fractions are given in Tables 2, 3 and 4 for leptonic, hadronic and combined fits. Good agreement is found, consistent with CP invariance.

To test the variation of the helicity fractions with the W^- production angle, Θ_{W^-} , the data are grouped in four bins of $\cos \Theta_{W^-}$. The ranges have been cho-

Table 2

W^- helicity fractions for the leptonic decays for the combined data sample. All the helicities are converted to W^- parameters using CP invariance. The first uncertainty is statistical, the second systematic. The corresponding helicity fractions in the Standard Model as implemented in the KORALW Monte Carlo program are also given with their statistical uncertainties

Sample	f_-	f_+	f_0
$W^- \rightarrow \ell^- \nu$ data	$0.559 \pm 0.038 \pm 0.016$	$0.201 \pm 0.026 \pm 0.015$	$0.240 \pm 0.051 \pm 0.017$
$W^+ \rightarrow \ell^+ \nu$ data	$0.625 \pm 0.037 \pm 0.016$	$0.179 \pm 0.023 \pm 0.015$	$0.196 \pm 0.050 \pm 0.017$
$W^\pm \rightarrow \ell^\pm \nu$ data	$0.589 \pm 0.027 \pm 0.016$	$0.189 \pm 0.017 \pm 0.015$	$0.221 \pm 0.036 \pm 0.017$
Monte Carlo	0.592 ± 0.003	0.170 ± 0.002	0.238 ± 0.004

Table 3

W^- helicity fractions for the hadronic decays for the combined data sample. All the helicities are converted to W^- parameters using CP invariance. The first uncertainty is statistical, the second systematic. The corresponding helicity fractions in the Standard Model as implemented in the KORALW Monte Carlo program are also given with their statistical uncertainties

Sample	f_\pm	f_0
$W^- \rightarrow$ hadrons data	$0.750 \pm 0.056 \pm 0.039$	$0.250 \pm 0.056 \pm 0.039$
$W^+ \rightarrow$ hadrons data	$0.833 \pm 0.062 \pm 0.039$	$0.167 \pm 0.062 \pm 0.039$
$W^\pm \rightarrow$ hadrons data	$0.785 \pm 0.042 \pm 0.039$	$0.215 \pm 0.042 \pm 0.039$
Monte Carlo	0.757 ± 0.004	0.243 ± 0.004

Table 4

W^- helicity fractions, measured combining leptonic and hadronic decays. All the helicities are converted to W^- parameters using CP invariance. The first uncertainty is statistical, the second systematic. The corresponding helicity fractions in the Standard Model as implemented in the KORALW Monte Carlo program are also given with their statistical uncertainties

	f_-	f_+	f_0
W^- data	$0.555 \pm 0.037 \pm 0.016$	$0.200 \pm 0.026 \pm 0.015$	$0.245 \pm 0.038 \pm 0.016$
W^+ data	$0.634 \pm 0.038 \pm 0.016$	$0.181 \pm 0.024 \pm 0.015$	$0.185 \pm 0.039 \pm 0.016$
W^\pm data	$0.592 \pm 0.027 \pm 0.016$	$0.190 \pm 0.017 \pm 0.015$	$0.218 \pm 0.027 \pm 0.016$
Monte Carlo	0.590 ± 0.003	0.169 ± 0.002	0.241 ± 0.003

sen such that large and statistically significant variations of the different helicity fractions are expected. Fig. 3 shows the four decay angle distributions for the leptonic W decays. The corrected distributions are fitted for leptonic and hadronic W decays separately in each bin of $\cos \Theta_{W^-}$. The fit results, combining leptonic and hadronic W decays, are shown in Table 6 and Fig. 4, together with the Standard Model expectations from the KORALW Monte Carlo. The results agree with the Standard Model expectation and demonstrate a strong variation of the W helicity fractions with the W^- production angle.

In conclusion, all three helicity states of the W boson are required in order to describe the data. Their fractions and their variations as a function of $\cos \Theta_{W^-}$ are in agreement with the Standard Model expectation.

Table 5

Systematic uncertainties on the measurement of f_0 for leptonic and hadronic W decays

	$W \rightarrow \ell \nu$	$W \rightarrow$ hadrons
Selection	0.013	0.024
Binning effects	0.007	0.029
Bias correction	0.006	0.011
Four fermion contamination	0.005	0.001
Background corrections	0.004	0.001
Total	0.017	0.039

The fraction of longitudinally polarised W bosons at $\sqrt{s} = 183\text{--}209$ GeV is measured as $0.218 \pm 0.027 \pm 0.016$. Separate analyses of the W^+ and W^- events are consistent with CP conservation.

Table 6

The W^- helicity fractions measured as a function of $\cos \Theta_{W^-}$ combining leptonic and hadronic W decays. The first uncertainty is statistical, the second systematic. The KORALW Monte Carlo expectations are also given with their statistical uncertainties

$\cos \Theta_{W^-}$	Fraction	W^\pm data	Monte Carlo
[-1.0, -0.3]	f_-	$0.173 \pm 0.041 \pm 0.033$	0.156 ± 0.006
	f_+	$0.418 \pm 0.060 \pm 0.043$	0.431 ± 0.008
	f_0	$0.409 \pm 0.082 \pm 0.051$	0.413 ± 0.008
[-0.3, 0.3]	f_-	$0.509 \pm 0.055 \pm 0.029$	0.446 ± 0.006
	f_+	$0.303 \pm 0.040 \pm 0.032$	0.282 ± 0.005
	f_0	$0.188 \pm 0.060 \pm 0.043$	0.272 ± 0.006
[0.3, 0.9]	f_-	$0.683 \pm 0.042 \pm 0.026$	0.723 ± 0.004
	f_+	$0.135 \pm 0.027 \pm 0.030$	0.119 ± 0.003
	f_0	$0.182 \pm 0.039 \pm 0.027$	0.158 ± 0.004
[0.9, 1.0]	f_-	$0.708 \pm 0.093 \pm 0.056$	0.647 ± 0.007
	f_+	$-0.010 \pm 0.055 \pm 0.028$	0.029 ± 0.004
	f_0	$0.302 \pm 0.082 \pm 0.059$	0.324 ± 0.007




Fig. 3. Corrected decay angle distributions for leptonic W decays separated into four different W^- production angle ranges, together with the KORALW expectation and fit results. Assuming CP invariance, W^+ decays are included with $\cos \Theta_{W^-} = -\cos \Theta_{W^+}$.




Fig. 4. W helicity fractions f_- , f_+ and f_0 and their statistical uncertainties for four different bins of $\cos \Theta_{W^-}$ in the combined data sample and in the KORALW Monte Carlo for $\sqrt{s} = 183\text{--}209$ GeV.

References

- [1] S.L. Glashow, Nucl. Phys. 22 (1961) 579;
S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264;
A. Salam, in: N. Svartholm (Ed.), Elementary Particle Theory, Almqvist & Wiksell, Stockholm, 1968, p. 367.
- [2] L3 Collaboration, B. Adeva, et al., Nucl. Instrum. Methods A 289 (1990) 35;
L3 Collaboration, O. Adriani, et al., Phys. Rep. 236 (1993) 1;
I.C. Brock, et al., Nucl. Instrum. Methods A 381 (1996) 236;
M. Chemarin, et al., Nucl. Instrum. Methods A 349 (1994) 345;
- [3] L3 Collaboration, M. Acciarri, et al., Phys. Lett. B 474 (2000) 194.
- [4] KORALW version 1.33 is used;
M. Skrzypek, et al., Comput. Phys. Commun. 94 (1996) 216;
M. Skrzypek, et al., Phys. Lett. B 372 (1996) 289.
- [5] PYTHIA version 5.722 is used;
T. Sjöstrand, preprint CERN-TH/7112/93 (1993), revised 1995;
T. Sjöstrand, Comput. Phys. Commun. 82 (1994) 74.

- [6] KK2F version 4.12 is used;
S. Jadach, B.F.L. Ward, Z. Was, *Comput. Phys. Commun.* 130 (2000) 260.
- [7] EEWW version 1.1 is used;
J. Fleischer, et al., *Comput. Phys. Commun.* 85 (1995) 29.
- [8] EXCALIBUR version 1.11 is used;
F.A. Berends, R. Pittau, R. Kleiss, *Comput. Phys. Commun.* 85 (1995) 437.
- [9] GEANT version 3.21 is used;
- R. Brun, et al., preprint CERN DD/EE/84-1 (1984), revised 1987.
- [10] H. Fesefeldt, RWTH Aachen report PITHA 85/02 (1985).
- [11] YFSWW3 version 1.14 is used;
S. Jadach, et al., *Phys. Rev. D* 54 (1996) 5434;
S. Jadach, et al., *Phys. Lett. B* 417 (1998) 326;
S. Jadach, et al., *Phys. Rev. D* 61 (2000) 113010;
S. Jadach, et al., *Phys. Rev. D* 65 (2002) 093010.