



ELSEVIER

6 August 1998

PHYSICS LETTERS B

Physics Letters B 433 (1998) 163–175

# Search for new physics phenomena in fermion-pair production at LEP

## L3 Collaboration

M. Acciarri<sup>ab</sup>, O. Adriani<sup>q</sup>, M. Aguilar-Benitez<sup>aa</sup>, S. Ahlen<sup>l</sup>, J. Alcaraz<sup>aa</sup>,  
G. Alemanni<sup>w</sup>, J. Allaby<sup>r</sup>, A. Aloisio<sup>ad</sup>, M.G. Alviggi<sup>ad</sup>, G. Ambrosi<sup>t</sup>,  
H. Anderhub<sup>aw</sup>, V.P. Andreev<sup>g,al</sup>, T. Angelescu<sup>n</sup>, F. Anselmo<sup>j</sup>, A. Arefiev<sup>ac</sup>,  
T. Azemoon<sup>c</sup>, T. Aziz<sup>k</sup>, P. Bagnaia<sup>ak</sup>, L. Baksay<sup>ar</sup>, S. Banerjee<sup>k</sup>, Sw. Banerjee<sup>k</sup>,  
K. Banicz<sup>at</sup>, A. Barczyk<sup>aw,au</sup>, R. Barillère<sup>r</sup>, L. Barone<sup>ak</sup>, P. Bartalini<sup>w</sup>,  
A. Baschirotto<sup>ab</sup>, M. Basile<sup>j</sup>, R. Battiston<sup>ah</sup>, A. Bay<sup>w</sup>, F. Becattini<sup>q</sup>, U. Becker<sup>p</sup>,  
F. Behner<sup>aw</sup>, J. Berdugo<sup>aa</sup>, P. Berges<sup>p</sup>, B. Bertucci<sup>ah</sup>, B.L. Betev<sup>aw</sup>,  
S. Bhattacharya<sup>k</sup>, M. Biasini<sup>ah</sup>, A. Biland<sup>aw</sup>, G.M. Bilei<sup>ah</sup>, J.J. Blaising<sup>d</sup>,  
S.C. Blyth<sup>ai</sup>, G.J. Bobbink<sup>b</sup>, R. Bock<sup>a</sup>, A. Böhm<sup>a</sup>, L. Boldizsar<sup>o</sup>, B. Borgia<sup>ak</sup>,  
D. Bourilkov<sup>aw</sup>, M. Bourquin<sup>t</sup>, S. Braccini<sup>t</sup>, J.G. Branson<sup>an</sup>, V. Brigljevic<sup>aw</sup>,  
I.C. Brock<sup>ai</sup>, A. Buffini<sup>q</sup>, A. Buijs<sup>as</sup>, J.D. Burger<sup>p</sup>, W.J. Burger<sup>ah</sup>, J. Busenitz<sup>ar</sup>,  
A. Button<sup>c</sup>, X.D. Cai<sup>p</sup>, M. Campanelli<sup>aw</sup>, M. Capell<sup>p</sup>, G. Cara Romeo<sup>j</sup>,  
G. Carlino<sup>ad</sup>, A.M. Cartacci<sup>q</sup>, J. Casaus<sup>aa</sup>, G. Castellini<sup>q</sup>, F. Cavallari<sup>ak</sup>,  
N. Cavallo<sup>ad</sup>, C. Cecchi<sup>t</sup>, M. Cerrada<sup>aa</sup>, F. Cesaroni<sup>x</sup>, M. Chamizo<sup>aa</sup>,  
Y.H. Chang<sup>ay</sup>, U.K. Chaturvedi<sup>s</sup>, S.V. Chekanov<sup>af</sup>, M. Chemarin<sup>z</sup>, A. Chen<sup>ay</sup>,  
G. Chen<sup>h</sup>, G.M. Chen<sup>h</sup>, H.F. Chen<sup>u</sup>, H.S. Chen<sup>h</sup>, X. Chereau<sup>d</sup>, G. Chiefari<sup>ad</sup>,  
C.Y. Chien<sup>e</sup>, L. Cifarelli<sup>am</sup>, F. Cindolo<sup>j</sup>, C. Civinini<sup>q</sup>, I. Clare<sup>p</sup>, R. Clare<sup>p</sup>,  
G. Coignet<sup>d</sup>, A.P. Colijn<sup>b</sup>, N. Colino<sup>aa</sup>, S. Costantini<sup>i</sup>, F. Cotorobai<sup>n</sup>,  
B. de la Cruz<sup>aa</sup>, A. Csilling<sup>o</sup>, T.S. Dai<sup>p</sup>, R. D'Alessandro<sup>q</sup>, R. de Asmundis<sup>ad</sup>,  
A. Degré<sup>d</sup>, K. Deiters<sup>au</sup>, D. della Volpe<sup>ad</sup>, P. Denes<sup>aj</sup>, F. DeNotaristefani<sup>ak</sup>,  
M. Diemoz<sup>ak</sup>, D. van Dierendonck<sup>b</sup>, F. Di Lodovico<sup>aw</sup>, C. Dionisi<sup>ak</sup>,  
M. Dittmar<sup>aw</sup>, A. Dominguez<sup>an</sup>, A. Doria<sup>ad</sup>, M.T. Dova<sup>s,l</sup>, D. Duchesneau<sup>d</sup>,  
P. Duinker<sup>b</sup>, I. Duran<sup>ao</sup>, S. Dutta<sup>k</sup>, S. Easo<sup>ah</sup>, H. El Mamouni<sup>z</sup>, A. Engler<sup>ai</sup>,  
F.J. Eppling<sup>p</sup>, F.C. Erné<sup>b</sup>, J.P. Ernenwein<sup>z</sup>, P. Extermann<sup>t</sup>, M. Fabre<sup>au</sup>,  
R. Faccini<sup>ak</sup>, S. Falciano<sup>ak</sup>, A. Favara<sup>q</sup>, J. Fay<sup>z</sup>, O. Fedin<sup>al</sup>, M. Felcini<sup>aw</sup>,  
T. Ferguson<sup>ai</sup>, F. Ferroni<sup>ak</sup>, H. Fesefeldt<sup>a</sup>, E. Fiandrini<sup>ah</sup>, J.H. Field<sup>t</sup>, F. Filthaut<sup>ai</sup>,  
P.H. Fisher<sup>p</sup>, I. Fisk<sup>an</sup>, G. Forconi<sup>p</sup>, L. Fredj<sup>t</sup>, K. Freudenreich<sup>aw</sup>, C. Furetta<sup>ab</sup>,

Yu. Galaktionov <sup>ac,p</sup>, S.N. Ganguli <sup>k</sup>, P. Garcia-Abia <sup>f</sup>, M. Gataullin <sup>ag</sup>,  
 S.S. Gau <sup>m</sup>, S. Gentile <sup>ak</sup>, N. Gheordanescu <sup>n</sup>, S. Giagu <sup>ak</sup>, S. Goldfarb <sup>w</sup>,  
 J. Goldstein <sup>l</sup>, Z.F. Gong <sup>u</sup>, A. Gougas <sup>e</sup>, G. Gratta <sup>ag</sup>, M.W. Gruenewald <sup>i</sup>,  
 R. van Gulik <sup>b</sup>, V.K. Gupta <sup>aj</sup>, A. Gurtu <sup>k</sup>, L.J. Gutay <sup>at</sup>, D. Haas <sup>f</sup>,  
 B. Hartmann <sup>a</sup>, A. Hasan <sup>ac</sup>, D. Hatzifotiadou <sup>j</sup>, T. Hebbeker <sup>i</sup>, A. Hervé <sup>r</sup>,  
 J. Hirschfelder <sup>ai</sup>, W.C. van Hoek <sup>af</sup>, H. Hofer <sup>aw</sup>, H. Hoorani <sup>ai</sup>, S.R. Hou <sup>ay</sup>, G. Hu <sup>e</sup>,  
 V. Innocente <sup>r</sup>, K. Jenkes <sup>a</sup>, B.N. Jin <sup>h</sup>, L.W. Jones <sup>c</sup>, P. de Jong <sup>r</sup>,  
 I. Josa-Mutuberria <sup>aa</sup>, R.A. Khan <sup>s</sup>, D. Kamrad <sup>av</sup>, J.S. Kapustinsky <sup>y</sup>, Y. Karyotakis <sup>d</sup>,  
 M. Kaur <sup>s,2</sup>, M.N. Kienzle-Focacci <sup>t</sup>, D. Kim <sup>ak</sup>, D.H. Kim <sup>aq</sup>, J.K. Kim <sup>aq</sup>,  
 S.C. Kim <sup>aq</sup>, W.W. Kinnison <sup>y</sup>, A. Kirkby <sup>ag</sup>, D. Kirkby <sup>ag</sup>, J. Kirkby <sup>r</sup>, D. Kiss <sup>o</sup>,  
 W. Kittel <sup>af</sup>, A. Klimentov <sup>p,ac</sup>, A.C. König <sup>af</sup>, A. Kopp <sup>av</sup>, I. Korolko <sup>ac</sup>,  
 V. Koutsenko <sup>p,ac</sup>, R.W. Kraemer <sup>ai</sup>, W. Krenz <sup>a</sup>, A. Kunin <sup>p,ac</sup>, P. Lacentre <sup>av,1</sup>,  
 P. Ladron de Guevara <sup>aa</sup>, I. Laktineh <sup>z</sup>, G. Landi <sup>q</sup>, C. Lapoint <sup>p</sup>, K. Lassila-Perini <sup>aw</sup>,  
 P. Laurikainen <sup>v</sup>, A. Lavorato <sup>am</sup>, M. Lebeau <sup>r</sup>, A. Lebedev <sup>p</sup>, P. Lebrun <sup>z</sup>,  
 P. Lecomte <sup>aw</sup>, P. Lecoq <sup>r</sup>, P. Le Coultre <sup>aw</sup>, H.J. Lee <sup>i</sup>, J.M. Le Goff <sup>r</sup>, R. Leiste <sup>av</sup>,  
 E. Leonardi <sup>ak</sup>, P. Levtchenko <sup>al</sup>, C. Li <sup>u</sup>, C.H. Lin <sup>ay</sup>, W.T. Lin <sup>ay</sup>, F.L. Linde <sup>b,r</sup>,  
 L. Lista <sup>ad</sup>, Z.A. Liu <sup>h</sup>, W. Lohmann <sup>av</sup>, E. Longo <sup>ak</sup>, W. Lu <sup>ag</sup>, Y.S. Lu <sup>h</sup>,  
 K. Lübelmeyer <sup>a</sup>, C. Luci <sup>ak</sup>, D. Luckey <sup>p</sup>, L. Luminari <sup>ak</sup>, W. Lustermann <sup>au</sup>,  
 W.G. Ma <sup>u</sup>, M. Maity <sup>k</sup>, G. Majumder <sup>k</sup>, L. Malgeri <sup>ak</sup>, A. Malinin <sup>ac</sup>, C. Maña <sup>aa</sup>,  
 D. Mangeol <sup>af</sup>, S. Mangla <sup>k</sup>, P. Marchesini <sup>aw</sup>, G. Marian <sup>ar,4</sup>, A. Marin <sup>l</sup>,  
 J.P. Martin <sup>z</sup>, F. Marzano <sup>ak</sup>, G.G.G. Massaro <sup>b</sup>, D. McNally <sup>r</sup>,  
 R.R. McNeil <sup>g</sup>, S. Mele <sup>r</sup>, L. Merola <sup>ad</sup>, M. Meschini <sup>q</sup>, W.J. Metzger <sup>af</sup>,  
 M. von der Mey <sup>a</sup>, D. Migani <sup>j</sup>, A. Mihul <sup>n</sup>, A.J.W. van Mil <sup>af</sup>,  
 H. Milcent <sup>r</sup>, G. Mirabelli <sup>ak</sup>, J. Mnich <sup>r</sup>, P. Molnar <sup>i</sup>, B. Monteleoni <sup>q</sup>,  
 R. Moore <sup>c</sup>, T. Moulík <sup>k</sup>, R. Mount <sup>ag</sup>, F. Muheim <sup>t</sup>, A.J.M. Muijs <sup>b</sup>, S. Nahn <sup>p</sup>,  
 M. Napolitano <sup>ad</sup>, F. Nessi-Tedaldi <sup>aw</sup>, H. Newman <sup>ag</sup>, T. Niessen <sup>a</sup>, A. Nippe <sup>w</sup>,  
 A. Nisati <sup>ak</sup>, H. Nowak <sup>av</sup>, Y.D. Oh <sup>aq</sup>, H. Opitz <sup>a</sup>, G. Organtini <sup>ak</sup>, R. Ostonen <sup>v</sup>,  
 S. Palit <sup>m</sup>, C. Palomares <sup>aa</sup>, D. Pandoulas <sup>a</sup>, S. Paoletti <sup>ak</sup>, P. Paolucci <sup>ad</sup>, H.K. Park <sup>ai</sup>,  
 I.H. Park <sup>aq</sup>, G. Pascale <sup>ak</sup>, G. Passaleva <sup>r</sup>, S. Patricelli <sup>ad</sup>, T. Paul <sup>m</sup>, M. Pauluzzi <sup>ah</sup>,  
 C. Paus <sup>r</sup>, F. Pauss <sup>aw</sup>, D. Peach <sup>r</sup>, Y.J. Pei <sup>a</sup>, S. Pensotti <sup>ab</sup>, D. Perret-Gallix <sup>d</sup>,  
 B. Petersen <sup>af</sup>, S. Petrak <sup>i</sup>, A. Pevsner <sup>e</sup>, D. Piccolo <sup>ad</sup>, M. Pieri <sup>q</sup>, P.A. Piroué <sup>aj</sup>,  
 E. Pistolesi <sup>ab</sup>, V. Plyaskin <sup>ac</sup>, M. Pohl <sup>aw</sup>, V. Pojidaev <sup>ac,q</sup>, H. Postema <sup>p</sup>, J. Pothier <sup>r</sup>,  
 N. Produit <sup>t</sup>, D. Prokofiev <sup>al</sup>, J. Quartieri <sup>am</sup>, G. Rahal-Callot <sup>aw</sup>, N. Raja <sup>k</sup>,  
 P.G. Rancoita <sup>ab</sup>, M. Rattaggi <sup>ab</sup>, G. Raven <sup>an</sup>, P. Razis <sup>ac</sup>, D. Ren <sup>aw</sup>, M. Rescigno <sup>ak</sup>,  
 S. Reucroft <sup>m</sup>, T. van Rhee <sup>as</sup>, S. Riemann <sup>av</sup>, K. Riles <sup>c</sup>, A. Robohm <sup>aw</sup>, J. Rodin <sup>ar</sup>,  
 B.P. Roe <sup>c</sup>, L. Romero <sup>aa</sup>, S. Rosier-Lees <sup>d</sup>, S. Roth <sup>a</sup>, J.A. Rubio <sup>r</sup>, D. Ruschmeier <sup>i</sup>,  
 H. Rykaczewski <sup>aw</sup>, J. Salicio <sup>r</sup>, E. Sanchez <sup>aa</sup>, M.P. Sanders <sup>af</sup>, M.E. Sarakinos <sup>v</sup>,  
 S. Sarkar <sup>k</sup>, C. Schäfer <sup>a</sup>, V. Schegelsky <sup>al</sup>, S. Schmidt-Kaerst <sup>a</sup>, D. Schmitz <sup>a</sup>,

N. Scholz<sup>aw</sup>, H. Schopper<sup>ax</sup>, D.J. Schotanus<sup>af</sup>, J. Schwenke<sup>a</sup>, G. Schwering<sup>a</sup>,  
 C. Sciacca<sup>ad</sup>, D. Sciarrino<sup>t</sup>, L. Servoli<sup>ah</sup>, S. Shevchenko<sup>ag</sup>, N. Shivarov<sup>ap</sup>,  
 V. Shoutko<sup>ac</sup>, J. Shukla<sup>y</sup>, E. Shumilov<sup>ac</sup>, A. Shvorob<sup>ag</sup>, T. Siedenburger<sup>a</sup>, D. Son<sup>aq</sup>,  
 B. Smith<sup>p</sup>, P. Spillantini<sup>q</sup>, M. Steuer<sup>p</sup>, D.P. Stickland<sup>aj</sup>, A. Stone<sup>g</sup>, H. Stone<sup>aj</sup>,  
 B. Stoyanov<sup>ap</sup>, A. Straessner<sup>a</sup>, K. Sudhakar<sup>k</sup>, G. Sultanov<sup>s</sup>, L.Z. Sun<sup>u</sup>,  
 G.F. Susinno<sup>t</sup>, H. Suter<sup>aw</sup>, J.D. Swain<sup>s</sup>, X.W. Tang<sup>h</sup>, L. Tauscher<sup>f</sup>, L. Taylor<sup>m</sup>,  
 Samuel C.C. Ting<sup>p</sup>, S.M. Ting<sup>p</sup>, S.C. Tonwar<sup>k</sup>, J. Tóth<sup>o</sup>, C. Tully<sup>aj</sup>, K.L. Tung<sup>h</sup>,  
 Y. Uchida<sup>p</sup>, J. Ulbricht<sup>aw</sup>, U. Uwer<sup>r</sup>, E. Valente<sup>ak</sup>, G. Vesztegombi<sup>o</sup>,  
 I. Vetlitsky<sup>ac</sup>, G. Viertel<sup>aw</sup>, M. Vivargent<sup>d</sup>, S. Vlachos<sup>f</sup>, R. Völkert<sup>av</sup>, H. Vogel<sup>ai</sup>,  
 H. Vogt<sup>av</sup>, I. Vorobiev<sup>r,ac</sup>, A.A. Vorobyov<sup>al</sup>, A. Vorvolakos<sup>ae</sup>, M. Wadhwa<sup>f</sup>,  
 W. Wallraff<sup>a</sup>, J.C. Wang<sup>p</sup>, X.L. Wang<sup>u</sup>, Z.M. Wang<sup>u</sup>, A. Weber<sup>a</sup>, S.X. Wu<sup>p</sup>,  
 S. Wynhoff<sup>a</sup>, J. Xu<sup>l</sup>, Z.Z. Xu<sup>u</sup>, B.Z. Yang<sup>u</sup>, C.G. Yang<sup>h</sup>, X.Y. Yao<sup>h</sup>, J.B. Ye<sup>u</sup>,  
 S.C. Yeh<sup>az</sup>, J.M. You<sup>ai</sup>, An. Zalite<sup>al</sup>, Yu. Zalite<sup>al</sup>, P. Zemp<sup>aw</sup>, Y. Zeng<sup>a</sup>,  
 Z. Zhang<sup>h</sup>, Z.P. Zhang<sup>u</sup>, B. Zhou<sup>l</sup>, G.Y. Zhu<sup>h</sup>, R.Y. Zhu<sup>ag</sup>, A. Zichichi<sup>j,r,s</sup>,  
 F. Ziegler<sup>av</sup>, G. Zilizi<sup>ar,4</sup>

<sup>a</sup> I. Physikalisches Institut, RWTH, D-52056 Aachen, FRG<sup>5</sup>

III. Physikalisches Institut, RWTH, D-52056 Aachen, FRG<sup>5</sup>

<sup>b</sup> National Institute for High Energy Physics, NIKHEF, and University of Amsterdam, NL-1009 DB Amsterdam, The Netherlands

<sup>c</sup> University of Michigan, Ann Arbor, MI 48109, USA

<sup>d</sup> Laboratoire d'Annecy-le-Vieux de Physique des Particules, LAPP, IN2P3-CNRS, BP 110, F-74941 Annecy-le-Vieux CEDEX, France

<sup>e</sup> Johns Hopkins University, Baltimore, MD 21218, USA

<sup>f</sup> Institute of Physics, University of Basel, CH-4056 Basel, Switzerland

<sup>g</sup> Louisiana State University, Baton Rouge, LA 70803, USA

<sup>h</sup> Institute of High Energy Physics, IHEP, 100039 Beijing, China<sup>6</sup>

<sup>i</sup> Humboldt University, D-10099 Berlin, FRG<sup>5</sup>

<sup>j</sup> University of Bologna and INFN-Sezione di Bologna, I-40126 Bologna, Italy

<sup>k</sup> Tata Institute of Fundamental Research, Bombay 400 005, India

<sup>l</sup> Boston University, Boston, MA 02215, USA

<sup>m</sup> Northeastern University, Boston, MA 02115, USA

<sup>n</sup> Institute of Atomic Physics and University of Bucharest, R-76900 Bucharest, Romania

<sup>o</sup> Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary<sup>7</sup>

<sup>p</sup> Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>q</sup> INFN Sezione di Firenze and University of Florence, I-50125 Florence, Italy

<sup>r</sup> European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland

<sup>s</sup> World Laboratory, FBLJA Project, CH-1211 Geneva 23, Switzerland

<sup>t</sup> University of Geneva, CH-1211 Geneva 4, Switzerland

<sup>u</sup> Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, China<sup>6</sup>

<sup>v</sup> SEFT, Research Institute for High Energy Physics, P.O. Box 9, SF-00014 Helsinki, Finland

<sup>w</sup> University of Lausanne, CH-1015 Lausanne, Switzerland

<sup>x</sup> INFN-Sezione di Lecce and Università Degli Studi di Lecce, I-73100 Lecce, Italy

<sup>y</sup> Los Alamos National Laboratory, Los Alamos, NM 87544, USA

<sup>z</sup> Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, F-69622 Villeurbanne, France

<sup>aa</sup> Centro de Investigaciones Energeticas, Medioambientales y Tecnológicas, CIEMAT, E-28040 Madrid, Spain<sup>8</sup>

<sup>ab</sup> INFN-Sezione di Milano, I-20133 Milan, Italy

<sup>ac</sup> Institute of Theoretical and Experimental Physics, ITEP, Moscow, Russia

<sup>ad</sup> INFN-Sezione di Napoli and University of Naples, I-80125 Naples, Italy

<sup>ae</sup> Department of Natural Sciences, University of Cyprus, Nicosia, Cyprus

<sup>af</sup> University of Nijmegen and NIKHEF, NL-6525 ED Nijmegen, The Netherlands

<sup>ag</sup> California Institute of Technology, Pasadena, CA 91125, USA

<sup>ah</sup> INFN-Sezione di Perugia and Università Degli Studi di Perugia, I-06100 Perugia, Italy

<sup>ai</sup> Carnegie Mellon University, Pittsburgh, PA 15213, USA

<sup>aj</sup> Princeton University, Princeton, NJ 08544, USA

<sup>ak</sup> INFN-Sezione di Roma and University of Rome, "La Sapienza", I-00185 Rome, Italy

<sup>al</sup> Nuclear Physics Institute, St. Petersburg, Russia

<sup>am</sup> University and INFN, Salerno, I-84100 Salerno, Italy

<sup>an</sup> University of California, San Diego, CA 92093, USA

<sup>ao</sup> Dept. de Física de Partículas Elementales, Univ. de Santiago, E-15706 Santiago de Compostela, Spain

<sup>ap</sup> Bulgarian Academy of Sciences, Central Lab. of Mechatronics and Instrumentation, BU-1113 Sofia, Bulgaria

<sup>aq</sup> Center for High Energy Physics, Korea Adv. Inst. of Sciences and Technology, 305-701 Taejeon, South Korea

<sup>ar</sup> University of Alabama, Tuscaloosa, AL 35486, USA

<sup>as</sup> Utrecht University and NIKHEF, NL-3584 CB Utrecht, The Netherlands

<sup>at</sup> Purdue University, West Lafayette, IN 47907, USA

<sup>au</sup> Paul Scherrer Institut, PSI, CH-5232 Villigen, Switzerland

<sup>av</sup> DESY-Institut für Hochenergiephysik, D-15738 Zeuthen, FRG

<sup>aw</sup> Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zürich, Switzerland

<sup>ax</sup> University of Hamburg, D-22761 Hamburg, FRG

<sup>ay</sup> National Central University, Chung-Li, Taiwan, ROC

<sup>az</sup> Department of Physics, National Tsing Hua University, Taiwan, China

Received 27 February 1998

Editor: K. Winter

---

## Abstract

The measurements of hadron and lepton-pair production cross sections and lepton-pair forward–backward asymmetries performed with the L3 detector at centre-of-mass energies between 130 GeV and 172 GeV are used to search for new physics phenomena. New physics effects involving four fermion vertices – contact interactions – are looked for in all channels. For hadron production the exchange of virtual leptoquarks and scalar quarks is studied. No evidence for deviations from the Standard Model expectations is found. Lower limits on the scale  $\Lambda$  of contact interactions in the range 1.2–7.1 TeV are obtained at the 95% confidence level for various models. Upper limits on the coupling strengths of leptoquarks and scalar quarks are derived. © 1998 Published by Elsevier Science B.V. All rights reserved.

---

## 1. Introduction

The successful running of LEP in 1995 and 1996 at centre-of-mass energies well above the Z resonance allows to search for new physics beyond the Standard Model [1]. Any significant deviation from the Standard Model predictions in the electron-positron annihilation into fermion-pairs would herald the presence of new phenomena. Four-fermion contact interactions offer a general framework for describing interactions beyond the Standard Model. They are characterised by a coupling strength,  $g$ , and by an energy scale,  $\Lambda$ , which can be viewed as the typical mass of new heavy particles being exchanged. For instance, if fermions are composite,

---

<sup>1</sup> Also supported by CONICET and Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina.

<sup>2</sup> Also supported by Panjab University, Chandigarh-160014, India.

<sup>3</sup> Supported by Deutscher Akademischer Austauschdienst.

<sup>4</sup> Also supported by the Hungarian OTKA fund under contract numbers T22238 and T026178.

<sup>5</sup> Supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie.

<sup>6</sup> Supported by the National Natural Science Foundation of China.

<sup>7</sup> Supported by the Hungarian OTKA fund under contract numbers T14459, T19181 and T24011.

<sup>8</sup> Supported also by the Comisión Interministerial de Ciencia y Tecnología.

such effects can occur. At energies much lower than  $\Lambda$ , the exchange of virtual new particles is described by an effective Lagrangian [2]:

$$\mathcal{L} = \frac{1}{1 + \delta_{ef}} \sum_{i,j=L,R} \eta_{ij} \frac{g^2}{\Lambda_{ij}^2} (\bar{e}_i \gamma^\mu e_i) (\bar{f}_j \gamma^\mu f_j), \quad (1)$$

where  $e_i$  and  $f_j$  denote the left- and right-handed initial-state electron and final-state fermion fields. The Kronecker symbol,  $\delta_{ef}$ , is zero except for the  $e^+e^-$  final state where it is one. The parameters  $\eta_{ij}$  define the contact interaction model by choosing the helicity amplitudes which contribute to the reaction  $e^+e^- \rightarrow f\bar{f}$ . The value of  $g/\Lambda$  determines the size of the expected effects. In a general search for contact terms the energy scale  $\Lambda$  is chosen by convention such that  $g^2/4\pi = 1$  and  $|\eta_{ij}| = 1$  or  $|\eta_{ij}| = 0$  is satisfied.

For hadronic final states two specific scenarios are investigated where  $t$ - or  $u$ -channel exchange of new particles coupling to quark–lepton pairs contribute. In the first scenario the exchange of leptoquarks [3,4] is studied. In the second scenario the exchange of supersymmetric scalar quarks violating R-parity [5] is investigated.

In this paper, our measurements of hadron and lepton-pair cross sections and lepton-pair forward–backward asymmetries are used to search for the existence of contact interactions. The effects of virtual exchange of leptoquarks and scalar quarks are investigated using our hadron cross section measurements only. In a previous publication we have used our data to search for R-parity breaking scalar neutrino exchange [6]. Limits on contact interactions and on leptoquark and scalar quark couplings have been presented by the OPAL collaboration [7] at LEP and by the CDF and DØ collaborations at the Tevatron [8].

## 2. Measurements of fermion-pair production

Measurements of cross sections and forward–backward asymmetries for the reactions  $e^+e^- \rightarrow f\bar{f}$  have been performed by the L3 experiment at centre-of-mass energies,  $\sqrt{s}$ , of 130.3, 136.3 and 140.2 GeV [9], and at  $\sqrt{s} = 161.3, 170.3$  and 172.3 GeV [10]. The L3 detector and its performance are described in Ref. [11].

For the  $e^+e^-$  final states both leptons have to be in the polar angular range  $44^\circ < \theta < 136^\circ$ , where  $\theta$  is the angle between the incoming electron and the outgoing lepton. Muon- and tau-pair candidates are selected with both leptons in the fiducial volume given by  $|\cos\theta| < 0.9$ . Hadron events are selected in the full solid angle.

In total 4305 hadron events and 1269 lepton-pair events are selected. These correspond to an integrated luminosity of  $26.2 \text{ pb}^{-1}$ . Due to the large  $Z$  exchange cross section the sensitivity to new physics is suppressed for centre-of-mass energies in the vicinity of the  $Z$  resonance. Therefore, a minimum effective centre-of-mass energy,  $\sqrt{s}_{\text{min}}$ , is required to reject radiative returns to the  $Z$ . The remaining samples contain in total 1179 hadron and 869 lepton-pair events. The details of the analyses as selection procedures, efficiencies, background contributions, measured cross sections and forward–backward asymmetries together with the statistical and systematic errors are discussed in Refs. [9,10].

## 3. Virtual effects in $e^+e^- \rightarrow f\bar{f}$

### 3.1. Contact interaction

For a general theory at an universal energy scale  $\Lambda \gg \sqrt{s}$ , new interactions are described by an effective contact interaction as shown in Fig. 1a. The differential cross section including a four-fermion contact




Fig. 1. a) Feynman diagram describing the contact interactions; (b) Feynman diagrams describing the  $t$ -channel exchange, left, and  $u$ -channel exchange, right, of leptoquarks with fermion number  $F = L + 3B$ ; (c) Feynman diagrams describing the  $t$ -channel exchange, left, and  $u$ -channel exchange, right, of scalar quarks.

interaction as a function of the final-state fermion scattering angle  $\theta$  is given by [12]:

$$\frac{1}{N_c} \frac{2s}{\pi\alpha^2} \frac{d\sigma}{d\cos\theta} = \left[ |A_{LR}^{ef}(t)|^2 + |A_{RL}^{ef}(t)|^2 \right] \left( \frac{s}{t} \right)^2 \delta_{ef} + \left[ |A_{LR}^{ef}(s)|^2 + |A_{RL}^{ef}(s)|^2 \right] \left( \frac{t}{s} \right)^2 + \left[ |A_{LL}^{ef}(s)|^2 + |A_{RR}^{ef}(s)|^2 \right] \left( \frac{u}{s} \right)^2, \quad (2)$$

with

$$\delta_{ef} = \begin{cases} 1, & \text{for } f=e \\ 0, & \text{for } f \neq e. \end{cases}$$

The Mandelstam variables are denoted  $s$ ,  $t$  and  $u$ . The electromagnetic fine structure constant and the colour factor are given by  $\alpha$  and  $N_c$ , respectively. The electroweak and the four-fermion contact interaction contribute to the helicity amplitudes:

$$\begin{aligned} A_{ij}^{ef}(y) &= Q_e Q_f + g_i^e g_j^f \chi(y) + \eta_{ij} \frac{y}{\alpha} \frac{1}{\Lambda^2}; & (i \neq j) \\ A_{ij}^{ef}(s) &= Q_e Q_f + g_i^e g_j^f \left[ \chi(s) + \frac{s}{t} \chi(t) \delta_{ef} \right] + \frac{s}{t} \delta_{ef} + (1 + \delta_{ef}) \eta_{ij} \frac{s}{\alpha} \frac{1}{\Lambda^2}; & (i=j), \end{aligned} \quad (3)$$

where  $Q$  is the electric charge,  $\chi$  is the Z propagator and  $y$  denotes either  $s$  or  $t$ . The left- or right-handed coupling,  $g_i^f$ , are

$$g_L^f = \frac{1}{\sin\theta_W \cos\theta_W} (I_3^f - Q_f \sin^2\theta_W), \quad (4)$$

$$g_R^f = \frac{1}{\sin\theta_W \cos\theta_W} (-Q_f \sin^2\theta_W), \quad (5)$$

where  $\theta_W$  is the weak mixing angle and  $I_3^f$  is the third component of the weak isospin. The differential cross section in Eq. (2) is rewritten in a general form as

$$\frac{d\sigma}{d\cos\theta} = \frac{d\sigma^{\text{SM}}}{d\cos\theta} + c_2(s, \cos\theta) \frac{1}{\Lambda^2} + c_4(s, \cos\theta) \frac{1}{\Lambda^4}. \quad (6)$$

The coefficients  $c_i$  represent the deviations from the Standard Model cross section,  $\sigma^{\text{SM}}$ , depending on the contact interaction model. The pure contact interaction amplitude and the interference between the Standard Model and the new physics are given by the terms including  $c_4$  and  $c_2$ , respectively. The helicity combinations of the specific models considered are defined in Table 1.

### 3.2. Leptoquark exchange

Leptoquarks couple to quark–lepton pairs from the same family, and preserve the baryon number  $B$  and the lepton number  $L$ . In  $e^+e^-$  collisions, leptoquarks of the first generation can be exchanged in the  $t$ - or  $u$ -channel leading to hadron final states as depicted in Fig. 1b. For leptoquarks the notation used in Ref. [13] is adopted where scalar leptoquarks,  $S$ , and vector leptoquarks,  $V$ , are classified as follows:

- Based on spin and weak isospin,  $I$ , the leptoquarks are divided into  $S_I$  and  $V_I$ , where an additional tilde indicates isomultiplets with different hypercharges.
- Leptoquark couplings are denoted  $g_L, g_R$ , with L,R referring to the chirality of the lepton. In the  $t$ - and  $u$ -channel leptoquarks can only be exchanged for special helicity combinations  $e^+e_i^- \rightarrow q_j\bar{q}$  with  $i, j = \text{L,R}$ .
- Leptoquarks carry fermion numbers,  $F = L + 3B$ . In the  $u$ -channel exchange  $F = 2$  while in the  $t$ -channel exchange  $F = 0$ .

The total cross section of quark-pair production in Born approximation, including the exchange of one leptoquark with either left or right coupling, is described by [4]

$$\sigma(e^+e^- \rightarrow q\bar{q}) = \sigma_{q\bar{q}}^{\text{SM}} + \frac{N_c}{128\pi s} \sum_{i=1}^4 k_i(g_{L,R}^2) C_i \left( \frac{m_{\text{LQ}}^2}{s} \right). \quad (7)$$

The interference between  $\gamma/Z$  and the leptoquark is described by the coupling coefficients  $k_{1,2}$  and the functions  $C_{1,2}$  and the squared leptoquark amplitude is given by the coupling coefficients  $k_{3,4}$  and the functions  $C_{3,4}$ .

Table 1

Models of contact interaction considered. The parameters  $\eta_{ij}$  ( $i, j = \text{L,R}$ ) define to which helicity amplitudes,  $A_{ij}$ , the contact interaction contributes. The models cover the interference of contact terms with single as well as with a combination of helicity amplitudes

Model	LL	RR	LR	RL	VV	AA	V0	A0	LL–RR
$\eta_{\text{LL}}$	$\pm 1$	0	0	0	$\pm 1$	$\pm 1$	$\pm 1$	0	$\pm 1$
$\eta_{\text{RR}}$	0	$\pm 1$	0	0	$\pm 1$	$\pm 1$	$\pm 1$	0	$\mp 1$
$\eta_{\text{LR}}$	0	0	$\pm 1$	0	$\pm 1$	$\mp 1$	0	$\pm 1$	0
$\eta_{\text{RL}}$	0	0	0	$\pm 1$	$\pm 1$	$\mp 1$	0	$\pm 1$	0

In the limit of  $m_{LQ} \gg \sqrt{s}$  the particle propagator approach of Eq. (7) reduces to the contact interaction approach of Eq. (6). The individual masses and couplings of leptoquarks are then related to the contact interaction scale by

$$\frac{g_{L,R}^2}{m_{LQ}^2} \sim \frac{1}{A_{ij}^2}. \quad (8)$$

### 3.3. Exchange of $R$ -parity breaking scalar quarks

Even in a minimal supersymmetric model the most general superpotential contains interactions violating  $R$ -parity in the trilinear couplings of superfields [14]. The only renormalisable gauge invariant operator that couples leptons, fermions and their scalar partners is given by [15]:

$$W_{\mathbb{R}} = \lambda_{ijk} L_L^i L_L^j \bar{E}_R^k + \lambda'_{ijk} L_L^i Q_L^j \bar{D}_R^k, \quad (9)$$

where  $L_L$  stands for the left-handed doublets of leptons,  $Q_L$  for the left-handed quark doublets and  $E_R$  and  $D_R$  for the right-handed singlets of charged leptons and down-type quarks. The family indices are  $i$ ,  $j$  and  $k$ . Limits on the couplings of scalar tau neutrinos to electrons,  $\lambda_{131}$ , and to muons,  $\lambda_{232}$ , are derived in Ref. [6] from our measurements of lepton-pair cross sections and forward–backward asymmetries. Similar studies for hadronic final states give limits on the couplings of scalar quarks to electrons and quarks,  $\lambda'_{1jk}$ , ( $j, k = 1, 2, 3$ ). The two possible scenarios are shown in Fig. 1c.

In the  $t$ -channel the relevant coupling is  $\lambda'_{1jk}$ , where  $j = 1, 2, 3$  is the family index of the exchanged scalar left-handed up-type quark, and  $k = 1, 2, 3$  is the family index of the final state down-type quark. In this case the scalar quarks couple to fermions like the  $\tilde{S}_{1/2}$  leptoquark. In the  $u$ -channel the relevant coupling is  $\lambda'_{1jk}$ , where  $j = 1, 2$  is the family index of the final state up-type quark and  $k = 1, 2, 3$  is the family index of the exchanged scalar right-handed down-type quark. In this case the scalar quarks couple to fermions in the same way as  $S_0$  leptoquarks with left-handed coupling  $g_L$ . At LEP centre-of-mass energies the coupling  $\lambda'_{13k}$  contributes only to the  $t$ -channel exchange since it is not possible to produce  $t\bar{t}$  final states. The scalar quark-exchange contributions are calculated using Eq. (7).

## 4. Analysis and results

The contributions of contact interaction and leptoquarks as given in Eqs. (2) and (7) are included into the improved Born cross section calculated with the program ZFITTER [16] and are convoluted to account for QED radiative corrections:

$$\sigma_{t,fb} = \int_{s'_{\min}}^s \sigma_{t,fb}^0(s') R_{t,fb}(s, s') ds', \quad (10)$$

$$A_{fb} = \frac{\sigma_{fb}}{\sigma_t}. \quad (11)$$

The total Born cross section,  $\sigma_t^0$ , and the difference between forward and backward Born cross sections,  $\sigma_{fb}^0$ , include electroweak and QCD corrections for the Standard Model part. Initial and final state QED corrections are taken into account in Eq. (10) by the radiator functions  $R_t$  and  $R_{fb}$ .

In order to determine the contributions of contact interactions or leptoquarks, the Standard Model parameters are fixed to  $\alpha_s(m_Z) = 0.118$  [17],  $\alpha(m_Z) = 1/128.894$  [18],  $m_Z = 91.195$  GeV [19],  $m_t = 175.6$  GeV [20] and  $m_H = 300$  GeV. The results of our analysis are not sensitive to the uncertainties on these parameters. The contributions of new physics are determined performing a  $\chi^2$ -fit to cross section and asymmetry measurements. Statistical and systematic errors of the measurements are incorporated as determined in Refs. [9,10]. No



Table 2

The one-sided 95% confidence level lower limits on the parameter  $\Lambda$  of contact interaction derived from fits. The limits  $\Lambda_+$  ( $\Lambda_-$ ) given in TeV correspond to the upper (lower) sign of the parameters  $\eta_{ij}$  in Table 1, respectively

	$f\bar{f}$		$q\bar{q}$		$u\bar{u}$		$d\bar{d}$	
	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$
LL	3.2	4.0	2.1	3.0	2.8	3.3	3.0	2.7
RR	3.2	3.9	2.7	2.3	2.4	1.2	1.4	2.0
LR	3.0	3.3	2.6	2.4	1.9	1.5	1.6	1.8
RL	3.5	3.8	3.2	2.0	1.7	1.7	2.0	1.4
VV	5.8	7.1	3.9	3.2	3.8	1.5	1.8	3.1
AA	4.2	5.1	2.9	4.3	3.5	1.7	1.7	3.5
V0	4.4	5.4	2.8	3.2	3.5	4.3	3.5	3.1
A0	4.5	5.2	3.7	2.4	2.2	1.8	2.2	1.8
LL-RR	2.5	3.6	2.5	3.6	2.3	1.7	1.6	2.5

deviations from the Standard Model expectations are seen. In their absence one sided lower limits at the 95% confidence level are derived for  $\Lambda$  and for  $g_{L,R}/m_X$  ( $X = LQ, \tilde{q}$ ).

In some cases the  $\chi^2$ -curve has more than one minimum. This is due to the quadratic form of the differential cross section in  $1/\Lambda^2$  as given in Eq. (6). In these cases the minimum with the smallest  $\Lambda$  that satisfies  $\chi^2(1/\Lambda) \geq \chi^2(1/\Lambda_{\min}) + 3.84$  is taken as a conservative lower limit.

#### 4.1. Limits on the scale $\Lambda$ of four-fermion contact interactions

The different models considered are summarised in Table 1. Atomic physics parity violation experiments probe with high precision the couplings of electrons to quarks of the first family, and place severe constraints on




Fig. 2. One-sided 95% confidence level lower limits on the scales  $\Lambda_+$  and  $\Lambda_-$  for contact interactions in hadronic channels and in all channels combined. The limits correspond to the values given in Table 2.

Table 3

The one-sided 95% confidence level lower limits on the parameter  $\Lambda$  of contact interaction derived from fits. The limits  $\Lambda_+$  ( $\Lambda_-$ ) given in TeV correspond to the upper (lower) sign of the parameters  $\eta_{ij}$  in Table 1, respectively

	$\ll$		$e^+e^-$		$\mu^+\mu^-$		$\tau^+\tau^-$	
	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$	$\Lambda_-$	$\Lambda_+$
LL	3.1	4.0	2.4	2.7	2.4	3.6	2.8	2.4
RR	3.0	3.9	2.4	2.7	2.2	3.5	2.6	2.3
LR	3.0	3.4	2.8	3.3	1.5	2.4	1.4	2.1
RL	3.0	3.4	2.8	3.3	1.5	2.4	1.4	2.1
VV	5.8	7.1	4.8	6.4	4.4	5.3	4.5	3.7
AA	4.1	4.9	1.9	3.3	3.3	4.9	3.9	2.9
V0	4.4	5.5	3.2	4.1	3.5	4.9	4.0	3.1
A0	4.3	4.8	3.9	4.8	1.6	3.1	1.5	2.8
LL-RR	2.1	2.3	1.9	1.9	2.1	2.4	1.7	1.8

the scale  $\Lambda$  of the order of 15 TeV [21]. The VV, AA, V0 and A0 models are parity conserving and hence are not constrained by these measurements.

The four-fermion contact interactions for the different types of final-state fermions are tested separately as well as for all flavours combined and lower limits on the scale  $\Lambda$  are derived. For hadron final states the following cases are analysed: the contact interaction affects only one flavour of up-type or down-type quarks, or all flavours at the same time. The results are given in Table 2 and depicted in Fig. 2. The lepton-pair final states are analysed separately for the three lepton channels and for all leptons combined. The lower limits obtained for  $\Lambda$  are summarised in Table 3 and Fig. 3. The limits  $\Lambda_+$  ( $\Lambda_-$ ) correspond to the upper (lower) sign of the parameters  $\eta_{ij}$  in Table 1, respectively.




Fig. 3. One-sided 95% confidence level lower limits on the scale  $\Lambda_+$  and  $\Lambda_-$  for contact interactions in leptonic channels. The limits correspond to the values given in Table 3.

#### 4.2. Limits on leptoquark couplings

Assuming leptoquarks with a mass of a few hundred GeV, upper limits on their couplings to quark-lepton pairs,  $g_{L,R}$ , are derived. The exchange of different types of leptoquarks is studied separately. The states  $S_0$ ,  $S_{1/2}$  and  $V_0$ ,  $V_{1/2}$  couple to both left- and right-handed quarks. Here, only one coupling,  $g_L$  or  $g_R$ , is assumed to be non-zero since low energy processes and rare decays of  $\pi$  and K largely constrain the product  $g_L g_R$  [22,23].

The allowed values for  $g_{L,R}$  are symmetrically distributed around zero. Our limits are presented in Fig. 4 for scalar leptoquarks and in Fig. 5 for vector leptoquarks. For a leptoquark with  $m_{LQ} = 200$  GeV upper limits on the absolute value of the couplings,  $g_L$  and  $g_R$  in the range 0.2 – 1.0 at the 95% confidence level are obtained.

The exchange of leptoquarks with higher masses is described by the interference and squared contact terms of Eq. (6). In this case the left- and right-handed couplings satisfy Eq. (8) and our limits approach those on contact interactions derived in the previous section.

#### 4.3. Determination of limits on R-parity violating scalar quarks

Upper limits on  $|\lambda'_{1jk}|$  ( $j, k = 1, 2, 3$ ) are derived assuming one single Yukawa coupling to be much larger than the others which are neglected. A strong constraint on the coupling  $\lambda'_{111}$  is derived from the neutrinoless double beta decay [24].

Here, two cases are analysed:

$$m_{\tilde{u}} \gg m_{\tilde{d}} \text{ with } \tilde{u} = \tilde{u}, \tilde{c}, \tilde{t} \text{ and } \tilde{d} = \tilde{d}, \tilde{s}, \tilde{b}$$

$$m_{\tilde{u}} \ll m_{\tilde{d}}$$

Only the exchange of the much lighter scalar quark type is important. Due to quark-universality all limits derived for the case  $m_{\tilde{u}} \gg m_{\tilde{d}}$  are the same. Analogously, all limits coming from the hypothesis  $m_{\tilde{u}} \ll m_{\tilde{d}}$  agree with each other.




Fig. 4. The 95% confidence level upper limits on  $|g_{L,R}|$  as a function of  $m_{LQ}$  for various scalar leptoquarks derived from hadronic final state cross sections. Limits are shown for fermion number  $F = 2$  (a) and for  $F = 0$  (b). Bounds on  $|\lambda'_{1jk}|$  for the exchange of scalar down-type quarks in the  $u$ -channel and scalar up-type quarks in the  $t$ -channel agree with limits on  $|g_L|$  for the  $S_0$  and  $\tilde{S}_{1/2}$  leptoquark exchange, respectively.




Fig. 5. The 95% confidence level upper limits on  $|g_{L,R}|$  as a function of  $m_{LQ}$  for various vector leptoquarks derived from hadronic cross sections. Limits are shown for fermion number  $F = 2$  (a) and for  $F = 0$  (b).

The amplitudes of right-handed scalar down-type quarks interfere with the helicity amplitude,  $A_{LL}$ , of the Standard Model, while amplitudes of left-handed scalar up-type quarks interfere with  $A_{LR}$  which is suppressed at centre-of-mass energies well above the Z resonance. Therefore the R-parity breaking Yukawa couplings are mainly restricted by virtual exchange of scalar down-type quarks in the  $u$ -channel. These limits are identical to those on  $|g_L|$  for  $S_0$  leptoquark exchange shown in Fig. 4a. In case of  $t$ -channel exchange of scalar quarks the limits on  $|\lambda'|$  agree with the result of the  $\tilde{S}_{1/2}$  leptoquark exchange shown in Fig. 4b. Assuming scalar up-type and down-type quark masses to be equal and both contributing to the hadronic cross section yields basically the same limits as for the case  $m_{\tilde{u}} \gg m_{\tilde{d}}$ . For scalar quark masses with  $m_{\tilde{q}} = 200$  GeV limits at the 95% confidence level of 0.4 and 0.8 for the two cases are obtained.

## 5. Conclusions

Our fermion-pair cross section and forward–backward asymmetry measurements are used to search for effects of new physics in terms of four-fermion contact interactions. No hint of deviations from the Standard Model is found. Limits on the energy scale  $\Lambda$  in the range 1.2–7.1 TeV are obtained.

The effects of the exchange of leptoquarks or R-parity violating scalar quarks in the  $u$ - or  $t$ -channel of hadron production are studied. In both cases, upper limits on the coupling constants,  $|g_L|$  and  $|g_R|$ , or  $|\lambda'|$  between 0.2 and 1.0 for masses  $m_{\tilde{X}} = 200$  GeV at the 95% confidence level are determined.

## Acknowledgements

We express our gratitude to the CERN accelerator divisions for the excellent performance of the LEP machine. We acknowledge the effort of all engineers and technicians who have participated in the construction

and maintenance of this experiment. We are grateful to J. Kalinowski, R. Rückl, H. Spiesberger, P. Zerwas and F. Teubert for stimulating discussions.

## References

- [1] S.L. Glashow, Nucl. Phys. 22 (1961) 579; S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264; A. Salam, Elementary Particle Theory, N. Svartholm (Ed.), Stockholm, Almquist & Wiksell, 1968, p. 367.
- [2] E. Eichten, K. Lane, M. Peskin, Phys. Rev. Lett. 50 (1983) 811.
- [3] W. Buchmüller, R. Rückl, D. Wyler, Phys. Lett. B 191 (1987) 442.
- [4] J. Kalinowski, R. Rückl, H. Spiesberger, P. Zerwas, Z. Phys. C 74 (1997) 595.
- [5] P. Fayet, Phys. Lett. B 69 (1977) 489; G. Farrar, P. Fayet, Phys. Lett. B 76 (1978) 575; N. Sakai, T. Yanagida, Nucl. Phys. B 179 (1982) 533.
- [6] L3 Collaboration, M. Acciarri et al., Phys. Lett. B 414 (1997) 373.
- [7] OPAL Collaboration, K. Akerstaff et al., Tests of the Standard Model and Constraints on New Physics from Measurements of Fermion-pair Production at 130–172 GeV at LEP, Preprint CERN-PPE-97-101, CERN, 1997, to be published in Z. Phys. C.
- [8] CDF Collaboration, F. Abe et al., Phys. Rev. Lett. 79 (1997) 2198; DØ Collab., B. Abbott et al., Phys. Rev. Lett. 80 (1998) 666.
- [9] L3 Collaboration, M. Acciarri et al., Phys. Lett. B 370 (1996) 195.
- [10] L3 Collaboration, M. Acciarri et al., Phys. Lett. B 407 (1997) 361.
- [11] L3 Collaboration, B. Adeva et al., Nucl. Inst. Meth. A 289 (1990) 35; M. Acciarri et al., Nucl. Inst. Meth. A 351 (1994) 300; M. Chemarin et al., Nucl. Inst. Meth. A 349 (1994) 345; I.C. Brock et al., Nucl. Inst. Meth. A 381 (1996) 236; A. Adam et al., Nucl. Inst. Meth. A 383 (1996) 342.
- [12] H. Kroha, Phys. Rev. D 46 (1992) 58.
- [13] A. Djouadi, T. Köhler, M. Spira, J. Tutas, Z. Phys. C 46 (1990) 679; B. Schrempp, in: W. Buchmüller, G. Ingelman (Eds.), Proceedings, Physics at HERA, Hamburg, 1991.
- [14] J. Wess, B. Zumino, Nucl. Phys. B 70 (1974) 39.
- [15] S. Dimopoulos, L. Hall, Phys. Lett. B 207 (1987) 210. Phys. Rev. D 40 (1989) 2987.
- [16] D. Bardin et al., FORTRAN package ZFITTER, and preprint CERN-TH. 6443/92; Z. Phys. C 44 (1989) 493; Nucl. Phys. B 351 (1991) 1; Phys. Lett. B 255 (1991) 290.
- [17] Particle Data Group, R.M. Barnett et al., Phys. Rev. D 54 (1996) 1.
- [18] S. Eidelmann, F. Jegerlehner, Z. Phys. C 67 (1995) 585.
- [19] L3 Collaboration, M. Acciarri et al., Z. Phys. C 62 (1994) 223.
- [20] CDF Collaboration, J. Lys, Top Mass Measurements at CDF, in: Proc. ICHEP96, Warsaw, 25–31 July 1996, 1196; DØ Collab., S. Abachi et al., Phys. Rev. Lett. 79 (1997) 1197; R. Raja, Top Quark Mass Measurements from the Tevatron Proc. XXXIInd Rencontres de Moriond, Les Arcs, 16–22 March 1997, p. 77.
- [21] C.S. Wood et al., Science 275 (1997) 1759; V. Barger, K. Cheung, K. Hagiwara, D. Zeppenfeld, Phys. Lett. B 404 (1997) 147; N. Di Bartolomeo, M. Fabbrichinesi, Phys. Lett. B 406 (1997) 237.
- [22] M. Leurer, Phys. Rev. D 49 (1994) 333; D 50 (1994) 536.
- [23] S. Davidson, D. Bailey, D. Campbell, Z. Phys. C 61 (1994) 613; M. Hirsch, H.V. Klapdor-Kleingrothaus, S.G. Kovalenko, Phys. Rev. D 54 (1996) R4207.
- [24] M. Hirsch, H.V. Klapdor-Kleingrothaus, S.G. Kovalenko, Phys. Rev. D 53 (1996) 1329.