Short communication Received: August 17, 2010 Revised version: September 10, 2010 Accepted: September 15, 2010

## Antioxidant, Allelopathic and Toxic Activity of Ochna serrulata

Guilherme COLLA, Mariana A. da SILVA, Gustavo S. QUEIROZ, Moacir G. PIZZOLATTI & Inês M.C. BRIGHENTE\*

Laboratório de Química de Produtos Naturais, Departamento de Química, Universidade Federal de Santa Catarina, Campus Universitário, 88040-900, Florianópolis, SC, Brasil

SUMMARY. Ochna serrulata (Ochnaceae) is an ornamental plant introduced into Brazil from Asia and Africa. Species of the Ochna genus are rich in phenolic compounds, mainly flavonoids. The biological screening of extracts and fractions showed that this plant exhibited a significant antioxidant activity, when evaluated by the DPPH and reducing potential assays. Ochna serrulata also demonstrated slight toxic activity against Artemia salina and a potential inhibitory allelopathic activity, when evaluated using the Lactuca sativa seed germination test. The ethyl acetate fraction, the most active one, was partitioned on a silica gel column to obtain epicatechin, which showed potential antioxidant activity.

KEY WORDS: Allelopathy, Antioxidants, Artemia salina, Epicatechin, Ochna serrulata.

\* Author to whom correspondence should be addressed. E-mail: ines@qmc.ufsc.br

ISSN 0326-2383