

## References

Amaral JF, Teixeira CG, Pinheiro ED, 1956. O bactério causador da mancha aureolada do cafeeiro. *Arquivos do Instituto Biológico* 23, 151–5.

Braun-Kiewnick A, Sands DC, 2001. *Pseudomonas*. In: Schaad NW, Jones JB, Chun W, eds. *Laboratory Guide for Identification of Plant Pathogenic Bacteria*. St. Paul, MN, USA: APS Press, 84–120.

Lelliott RA, Billing E, Hayward AC, 1966. A determinative scheme for the fluorescent plant pathogenic pseudomonads. *Journal of Applied Bacteriology* 29, 470–89.

Rodrigues-Neto J, Figueiredo P, Mariotto PR, Robbs CF, 1981. *Pseudomonas andropogonis* (Smith, 1911) Stapp, 1928, agente causal da “mancha escura bacteriana” em folhas de cafeeiro (*Coffea arabica* L.). *Arquivos do Instituto Biológico* 48, 31–6.

\*E-mail: julio@biologico.sp.gov.br. Accepted 21 July 2010 at <http://www.bspp.org.uk/ndr> where figures relating to this paper can be viewed.

*Plant Pathology* (2010) 59, 1163

Doi: 10.1111/j.1365-3059.2010.02291.x

## First report of *Pseudomonas mediterranea* causing tomato pith necrosis in Argentina

A. M. Alippi\* and A. C. López

CIDEFI - Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, calles 60 y 119, c.c. 31, 1900 La Plata, Argentina

During the summers of 2007 and 2008 fruiting tomato plants (*Solanum lycopersicum* cv. Orco) from commercial glasshouses near La Plata, Argentina (35°S 57°W) showed abundant adventitious root production, apical chlorosis of leaves and a brown discoloration of the stem pith. These symptoms were similar to those reported by López *et al.* (1994) and Catara *et al.* (2002) on tomatoes affected by *Pseudomonas corrugata* or *P. mediterranea*. Bacteria consistently isolated from stem lesions formed cream-coloured, glistening, convex colonies on sucrose peptone agar and were non-fluorescent on King's medium B. Four strains were selected for further study. All were aerobic, Gram-negative rods with PHB inclusions. In LOPAT tests, all induced a hypersensitive response in tobacco plants, were oxidase positive, did not cause soft rot of potato tubers, and were negative for levan and arginine dihydrolase. Colonies developed at 28°C and 37°C but not at 41°C. Additional characterization was achieved by API 20 NE tests strips (Biomérieux®). Reference strains 5367 (Spain), 5924 (Spain) and CFBP 10906 (France) of *P. mediterranea* and strain NCPPB 2445 of *P. corrugata* were included in all tests for comparison.

Further identity was confirmed by PCR, using species-specific primers PCS/1-PCS/2 for *P. mediterranea* and primers PC1/1-PC1/2 for *P. corrugata* (Catara *et al.*, 2002). All the strains were identified by the amplification of a 600 bp DNA fragment characteristic of *P. mediterranea* (Catara *et al.*, 2002). The strains of *P. mediterranea* were also differentiated from those of *P. corrugata* by PCR/RFLP analysis of 16S rDNA gene by using endonuclease AluI.

Pathogenicity was verified on four-week-old tomato plants (cv. Presto) by injecting bacterial suspensions at 10<sup>7</sup> cfu mL<sup>-1</sup> or sterile distilled water for controls, after which the plants were kept for 72 h in a humid chamber before incubation at 25°C. After 45 days inoculated plants

showed necrotic pith symptoms similar to those observed on field-grown plants, whereas no lesions were observed on controls.

Pith necrosis caused by *P. corrugata* and *P. viridiflava* has been previously reported in Argentina (Alippi *et al.*, 1993, 2003). This is the first report of a disease caused by *P. mediterranea* on glasshouse-grown tomatoes in Argentina and South America.

### Acknowledgements

A.M. Alippi and A.C. López are Career Researchers of CIC and CONICET (Argentina), respectively.

### References

Alippi AM, Ronco L, Alippi HE, 1993. Tomato pith necrosis caused by *Pseudomonas corrugata* in Argentina. *Plant Disease* 77, 428.

Alippi AM, Dal Bo E, Ronco LB, López MV, López AC, Aguilar OM, 2003. *Pseudomonas* populations causing pith necrosis of tomato and pepper in Argentina are highly diverse. *Plant Pathology* 52, 287–302.

Catara V, Sutra L, Morineau A, Achouak W, Christen R, Gardan L, 2002. Phenotypic and genomic evidence for the revision of *Pseudomonas corrugata* and proposal of *Pseudomonas mediterranea* species sp. nov. *International Journal of Systematic and Evolutionary Microbiology* 52, 1749–58.

López MM, Siverio F, Albiach MR, García F, Rodríguez R, 1994. Characterization of Spanish isolates of *Pseudomonas corrugata* from tomato and pepper. *Plant Pathology* 43, 80–90.

\*E-mail: alippi@biol.unlp.edu.ar. Accepted 13 January 2010 at <http://www.bspp.org.uk/ndr> where figures relating to this paper can be viewed.