

## ORIGINAL ARTICLE

**Genotypic and phenotypic characterization of *Bordetella pertussis* strains used in different vaccine formulations in Latin America**

D. Bottero, M.E. Gaillard, L.A. Basile, M. Fritz and D.F. Hozbor

Laboratorio VacSal, Instituto de Biotecnología y Biología Molecular, CONICET – Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, La Plata, Argentina

**Keywords**

disease, pertussis, proteomic, re-emergence, vaccine strains.

**Correspondence**Daniela Hozbor, Laboratorio VacSal, Instituto de Biotecnología y Biología Molecular, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, CCT La Plata CONICET, Calles 47 y 115, 1900 La Plata, Argentina.  
E-mail: hozbor@biol.unlp.edu.ar

2011/2142: received 19 December 2011, revised 19 March 2012 and accepted 26 March 2012

doi:10.1111/j.1365-2672.2012.05299.x

**Abstract****Aim:** To characterize *Bordetella pertussis* vaccine strains in comparison with current circulating bacteria.**Methods and Results:** Genomic and proteomic analyses of Bp137 were performed in comparison with other vaccine strains used in Latin America (Bp509 and Bp10536) and with the clinical Argentinean isolate Bp106. Tohama I strain was used as reference strain. Pulse-field gel electrophoresis (PFGE) and pertussis toxin promoter (*ptxP*) sequence analysis revealed that Bp137 groups with Bp509 in PFGE group III and contains *ptxP2* sequence. Tohama I (group II) and Bp10536 (group I) contain *ptxP1* sequence, while Bp106 belongs to a different PFGE cluster and contains *ptxP3*. Surface protein profiles diverged in at least 24 peptide subunits among the studied strains. From these 24 differential proteins, Bp10536 shared the expression of ten proteins with Tohama I and Bp509, but only three with Bp137. In contrast, seven proteins were detected exclusively in Bp137 and Bp106.**Conclusions:** Bp137 showed more features in common with the clinical isolate Bp106 than the other vaccine strains here included.**Significance and Impact of the Study:** The results presented show that the old strains included in vaccines are not all equal among them. These findings together with the data of circulating bacteria should be taken into account to select the best vaccine to be included in a national immunization programme.**Introduction**

Pertussis or whooping cough is an immune-preventable respiratory disease that is still endemic worldwide among infants. This age group is most at risk of morbidity, hospitalization and mortality. Estimates from WHO suggest that in 2008, about 16 million cases of pertussis occurred world-wide, 95% of which were in developing countries, and that about 195 000 children died from the disease (World Health Organization 2010). The best way to prevent this highly contagious disease is to get vaccinated. Two types of pertussis vaccines are available: whole-cell (wP) vaccines based on killed aetiological pathogen (*Bordetella pertussis*) and acellular (aP) vaccines based on

highly purified, selected bacterial components. Although for paediatric population, wP or aP vaccines could be used, for adolescent and adults, only aP vaccine with lower dose of immunogens is recommended to reduce the reactogenicity associated with the other vaccine formulations (World Health Organization 2010).

The optimal pertussis immunization schedule and the appropriate time for booster dose in a country are normally assessed based on its current epidemiological situation. Because of that, epidemiological surveillance of pertussis is encouraged worldwide. Moreover, the reported shift in the antigenic characteristics of *Bord. pertussis* circulating strains (Mooi *et al.* 1998; Hozbor *et al.* 2009) makes such surveillance crucial to evaluate the

potential impact of bacterial shift on the overall immunity of a population. To control the increasing number of pertussis cases, many countries that do not produce vaccines must import the vaccine doses required to handle the demands of its population. In countries where wP vaccines are still being used, the selection of the vaccine to be imported is a challenge in itself because not all vaccines are formulated with the same strain or the same combination of strains. Latin American countries are using wP vaccines that contain among others the *Bord. pertussis* strains *Bp10536*, *Bp509* and *Bp137*. In our previous work, we have characterized the first two vaccine strains (*Bp10536* and *Bp509*) and have observed not only differences between them but also a representative isolate of the currently circulating bacterial population. *Bp137* strain has been included in a Brazilian vaccine successfully used in their national vaccination programme for more than 17 years (Pereira *et al.* 2005). However, the properties of this strain are scarcely studied. In this work, we present the results obtained from proteomic and genomic studies on this strain and their comparison with those from other vaccine strains. Results from the current clinical isolate *Bp106* were also included.

## Materials and methods

### Bacterial strains and growth conditions

The strains of *Bord. pertussis* used in this study were Tohama I (Kasuga *et al.* 1954a,b,c) obtained from the collection of the Pasteur Institute, France, *Bp509* (van Hemert 1969) obtained from the Netherlands Vaccine Institute, and *Bp10536* (Stainer and Scholte 1970) and *Bp137* (Pereira *et al.* 2005) obtained from the National Administration of Laboratories and Institutes of Health. The last three strains are widely used in wP vaccines in Latin America (Table 1). The Argentinean clinical isolate,

*Bp106*, which was collected in 2001 from an infant patient residing in Buenos Aires, was also included (Bottero *et al.* 2007). The strains and isolates were cultured on Bordet–Gengou agar (BGA, Difco) supplemented with 1% glycerol, Bacto-peptone (Difco) 10 g l<sup>-1</sup> and 10% (v/v) defibrinated sheep blood and incubated at 36°C for 3 days. Then, the bacteria were replated in the same medium for 24 h. Bacterial suspensions prepared from these plates were used for genomic analysis [PCR, sequencing and pulse-field gel electrophoresis (PFGE)].

For proteomic experiments, subcultures were grown in Stainer–Scholte liquid medium (Stainer and Scholte 1970) for 20 h at 36°C until the optical density at 650 nm reached 1.0.

### PCR, sequencing and PFGE

PCR, sequencing and PFGE were performed as previously described (Mooi *et al.* 2000, 2009; Hardwick *et al.* 2002b; van Loo *et al.* 2002; Fiett *et al.* 2003; Advani *et al.* 2004; Schouls *et al.* 2004; Borisova *et al.* 2007; Bottero *et al.* 2007). The sequences of the primers used to amplify and sequence the promoter region of pertussis toxin (*ptxP*), subunit A of pertussis toxin (*ptxA*), pertactin (*prn*), and type 2 (*fim2*) and type 3 (*fim3*) fimbriae are given in Table 2.

The obtained *Xba*I PFGE profiles were analysed using BioNUMERICS (Applied Maths, Sint-Martens-Latem, Belgium) software version 3.5. The unweighted pair group method with arithmetic mean (UPGMA) algorithm was used as the clustering method, with a 1% band tolerance and 1% optimization settings with the Dice's coefficient. The band pattern of each strain was verified by visual comparison. PFGE profiles were classified into groups based on a criterion of similarity higher than 82%.

**Table 1** Vaccine strains used in this study

Vaccine strain	Origin of the strain	Year of isolation	wP vaccine—manufacturing countries in Latin America	
			Before 1996	At present
Tohama I	Japan	1954	Chile	None
<i>Bp509</i>	the Netherlands	1950	Cuba	Cuba
			Mexico	Mexico
<i>Bp10536</i>	USA	Before 1940	Venezuela	Venezuela
			Colombia	None
<i>Bp137</i>	USA	No data available	Ecuador	Brazil
			Uruguay	Ecuador
			Brazil	

**Table 2** Primers used in this study

Gene	Primer sequence	References
<i>ptxP</i>	F: 5'-AATCGTCTGCTCAACCGCC-3'	Schouls <i>et al.</i> (2004), Mooi <i>et al.</i> (2009)
	R: 5'-GGTATACGGTGCGGGAGGA-3'	
<i>ptxA</i>	F: 5'-CCCCTGCCATGGTGTGATC-3'	Fiett <i>et al.</i> (2003)
	R: 5'-TCAATTACCGGAGTTGGGCG-3'	
<i>prn</i>	F: 5'-CAATGTCACGGTCCAA-3'	Mooi <i>et al.</i> (2000)
	R: 5'-GCAAGGTGATCGACAGGG-3'	
<i>fim2</i>	F: 5'-GCGCCGGCCCTGCATGCAC-3'	Van Loo and Mooi (2002), Borisova <i>et al.</i> (2007)
	R: 5'-GGGGGGTTGGCGATTCCAGTTCTC-3'	
<i>fim3</i>	F: 5'-GACCTGATATTCTGATCCG-3'	Borisova <i>et al.</i> (2007)
	R: 5'-AAGGCTTGCCGTTTTTTTGG-3'	

### Serotyping

Serotype analysis was performed using an agglutination assay with monoclonal antibodies against type 2 fimbriae (Fim2; NIBSC, 04/154) and type 3 fimbriae (Fim3, NIBSC, 04/156) according to EU pertstrain group recommendations (<http://www.eupertstrain.org>). Briefly, 15  $\mu$ l of bacterial suspension in PBS was mixed on slide with an equal volume of 1/10 dilution of monoclonal antibodies against Fim2 and 1/100 dilution of monoclonal antibodies against Fim3. If the agglutination reaction was obtained with either Fim2, Fim3, or both antibodies, the serotype was defined as Fim2, Fim3 or Fim2,3, respectively. If no reaction was detected, the serotype was defined as untypeable. Autoagglutination was examined with phosphate-buffered saline in parallel with monoclonal antibodies.

### Membrane protein enrichment for two-dimensional polyacrylamide gel electrophoresis (2-DE)

Membrane fractions were prepared as described previously (Bottero *et al.* 2007). Briefly, *Bord. pertussis* cells were harvested by centrifugation (10 000 g; 30 min; 4°C) and washed twice with low-salt washing buffer containing 3 mmol l<sup>-1</sup> KCl, 68 mmol l<sup>-1</sup> NaCl, 1.5 mmol l<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub> and 9 mmol l<sup>-1</sup> NaH<sub>2</sub>PO<sub>4</sub>. The cells were suspended in 10 mmol l<sup>-1</sup> Tris-HCl (pH 8.5) supplemented with phenylmethylsulphonyl fluoride and protease inhibitor cocktail tablets (Roche Applied Science, Buenos Aires, Argentina) and then disrupted with an ultrasonicator (Sonics & Materials, Inc., Danbury, CT, USA). DNase and RNase (20  $\mu$ g ml<sup>-1</sup> each) were added to the cell suspension, and the mixture was incubated at 37°C for 1 h. The unbroken cells were removed by centrifugation (12 000 g; 30 min; 4°C), and the supernatant was retained. Total membrane proteins were then collected by centrifugation (30 000 g, 1 h; 4°C) and resuspended in 7 mol l<sup>-1</sup> urea, 2 mol l<sup>-1</sup> thiourea, 10% isopropanol and 2% Triton X-100. Membrane proteins were divided into aliquots and stored at -20°C.

Sample preparation, 2-DE and protein identification were repeated at least four times for each strain.

### Protein quantification

Protein concentrations were determined by the Bradford's method (Bradford 1976) with bovine serum albumin (Sigma) as a standard.

### 2-DE

The method previously described by Bottero *et al.* (2007) was followed. Seven-centimetre Immobiline DryStrip

(IPG, pH 4-7; Amersham Biosciences) dissolving 200  $\mu$ g of the membrane proteins in a volume of 125  $\mu$ l of rehydration buffer (7 mol l<sup>-1</sup> urea, 2 mol l<sup>-1</sup> thiourea, 10% isopropanol and 2% Triton X-100) plus 1.25  $\mu$ l 28% dithiothreitol (DTT), 0.62  $\mu$ l 0.5% ampholyte (pH 4.0-7.0 [Amersham]) and 0.01% bromophenol blue was rehydrated overnight at room temperature. Three preset programmes were executed with slight modifications so that the focusing conditions consisted of the conditioning step, voltage ramping and final focusing. After IEF, the strips were equilibrated in 50 mmol l<sup>-1</sup> Tris buffer (pH 8.8) containing 6 mol l<sup>-1</sup> urea, 2% sodium dodecyl sulphate, 30% glycerol and 1% DTT, followed by another 1-h equilibration step with the same buffer supplemented with 4.5% iodoacetamide. SDS-PAGE was performed according to (Laemmli 1970) with a 12.5% resolving polyacrylamide gel without a stacking gel. Separation in the second dimension was carried out at 40 V at 4°C until the running dye reached the bottom of the gel.

Proteins were visualized using a colloidal Coomassie staining method (<http://prospector.ucsf.edu>) with the modifications described previously (Bottero *et al.* 2007). A gel image was captured in a UVP Bioimaging system Epi Chemi3 Darkroom with a Hamamatsu Photonic systems camera, model 1394 C8484-51-03G, controlled by Labworks image acquisition and analysis software version 4.6.00.0. The 8-bit grey-scale tif files obtained were later processed with the IMAGE MASTER 2D PLATINUM software ver. 6.0 (GE Healthcare Argentina S.A., CABA, Argentina).

### MALDI-TOF-MS analysis and database search

Coomassie-stained spots were excised from 2-DE gels for tryptic in-gel digestion and MALDI-TOF-MS with an Ultraflex (Bruker) (Bottero *et al.* 2007). Peptide mass fingerprint (PMF) data were searched against the NCBI database in MASCOT server (<http://www.matrix-science.com>) for sequence match. The MASCOT search parameters were as follows: (i) species, bacteria (eubacteria); (ii) allowed number of missed cleavages (only for trypsin digestion), 1; (iii) variable post-translational modification, methionine oxidation; (iv) fixed modification, carbamidomethylation; (v) peptide tolerance,  $\pm$ 50 ppm; (vi) peptide charge, +; and (vii) mono-isotopic peptide masses that were used to search the database, allowing a molecular mass range for 2-DE analyses of  $\pm$ 15%. Only significant hits as defined by MASCOT probability analysis were considered. Prediction of protein localization was carried out using a PSORTb.2, PSORTb.3 algorithm available at <http://psort.nibb.ac.jp> and Proteome Analyst (PA) (Lu *et al.* 2004).

## Results

### Genotypic analysis

Chromosomal DNA samples from *Bp137* and two other vaccine strains (*Bp10536* and *Bp509*) used in some Latin America countries were digested with *Xba*I and examined by PFGE. The profiles obtained were compared with that from the reference strain Tohama I (Fig. 1a). The profiles were distributed in three groups classified according to a criterion of similarity higher than 80%. The vaccine strain *Bp137* grouped with *Bp509* in PFGE group III. The similarity between these strains was 83%. Group I included *Bp10536*, and group II was composed of the Japanese vaccine strain Tohama I.

The representative isolate *Bp106* collected after the introduction of a massive vaccination programme in Argentina is clearly separated from vaccine strains as we previously reported (Bottero *et al.* 2007).

Regarding the genotypification of well-known polymorphic sequences described for virulence factors of *Bord. pertussis*, vaccine strains *Bp137* and *Bp509* present pertussis toxin promoter *ptxP2* and the allele *fim2-2*. These genotypes are different from those of the other vaccine strains (Fig. 1b).

In contrast, the representative clinical isolate *Bp106* contained *ptxP3*, *ptxA1*, *prn2*, *fim2-1* and *fim3-B* alleles. In fact, we observed this genotype in the majority of the current members of our collection of circulating clinical isolates (data not shown). Regarding the *fim2* and *fim3* alleles, 97% of the collection, including the *Bp106* representative strain, is *fim2-1* and 76% has the variant B for the *fim3* allele. In relation to the *fim3* allele, the vaccine strains included in our study have the variant A.


The Fim serotypes are Fim2 for Tohama I strain and Fim2,3 for *Bp10536*, *Bp137* and *Bp509*. In our study, the Fim serotype for *Bp106* and for 97% of clinical isolates of our collection was Fim3.

### Proteomic analysis


We characterized *Bp137* strain by proteomic analysis and compared its surface proteome with the proteomes of the other strains previously reported but repeated here (Bottero *et al.* 2007; Supporting Information, Fig. S1). In the analysis, we also included the data of human and murine immunoproteomes already performed (Altindis *et al.* 2009; Zhu *et al.* 2010; Tefon *et al.* 2011).

The 2-DE profile of *Bp137* revealed more than 90 protein spots from which 50 proteins were successfully identified (Fig. 2). For this work, we have repeated the 2-DE of surface proteins of the other four strains (*Bp10536*, *Bp509*, Tohama I and *Bp106*). In all instances, we have confirmed previously published data, but in addition, we have identified more spots (64 spots in total). Of the total identified peptide subunits, 12 were predicted to be associated with the external membrane/extracellular localization, ten had periplasmic localization, nine had cytoplasmic membrane localization, eight had an unknown or undefined origin and 25 had a cytoplasmic localization (Table 3). As observed for the other vaccine strains, some of the proteins separated by 2-DE were present as multiple spots exhibiting variability in pI values (horizontal spot patterns, Fig. 2). Charge variants included the following proteins: EF-Tu, 60-kDa chaperonin, outer membrane porin protein precursor, serum resistance protein and serine protease. These may represent natural isoforms or an artefact caused by sample preparation or 2-DE.

From the proteins identified by MALDI-TOF-MS, 14 are involved in small-molecule metabolism (BP2360, BP0277, BP2439, BP2386, BP3288, BP3125, BP0995, BP0379, BP3215, BP1126, BP0844, BP1499, BP0843 and BP0047), seven are associated with macromolecule biosynthesis and degradation (BP2434, BP0007, BP3642, BP2361, BP1420, BP1455 and BP2470), 15 are classified in the category cell structure (BP1146, BP1296, BP3405,



**Figure 1** (a) Genomic analysis of *Bordetella pertussis* strains used for vaccine production. The chromosomal DNA profiles obtained after digestion with *Xba*I are shown on the left side and the identifier of strains on the right side. (b) Characteristics of vaccine strains used in this work.



**Figure 2** 2-D proteome of *Bordetella pertussis* vaccine strain Bp137. Preparations of membrane-enriched protein samples were separated by IEF at pH 4–7 in the first dimension and then by 12.5% SDS-PAGE in the second dimension. Protein spots were visualized by colloidal Coomassie staining. The spot numbers refer to the identified peptide subunits by MALDI-TOF.

BP0840, BP1440, BP3862, BP0943, BP2513, BP2755, BP3150, BP1630, BP2750, BP3559, BP3077 and BP1485), 14 are associated with cellular processes (BP3757, BP1487, BP3322, BP0965, BP3495, BP1285, BP2761, BP3794, BP2747, BP3552, BP1774, BP2235, BP2499 and BP2744), two have general regulatory roles (BP2483 and BP2435), three are associated with phages, transposons and pathogenicity islands (BP2667, BP3494 and BP1054) and, finally, six have unknown function (BP3441, BP2196, BP3128, BP3515, BP2964 and BP1203) according to Riley categories (Riley 1993).

Twenty of the 64 identified proteins were not detected in at least one of the strains studied, and four proteins were detected only in the local isolate *Bp106* (Table 3). Tohama I and *Bp509* have very similar protein profiles with only one differential subunit peptide (spot 10). However, these two strains share the expression of only three of the 24 differential proteins with *Bp137*. Interestingly, we note that seven of 24 peptide subunits were expressed exclusively by the vaccine strain *Bp137* and the clinical isolate *Bp106*. Peptide subunit Bp2235 (spot 53), a potential protein of type III secretion system (TTSS), belongs to this group of seven subunits. Two other proteins identified only in *Bp106* and *Bp137*, but not detected in the rest of the vaccine strains, are BP3150 and BP1630, assigned to polysaccharide biosynthesis and capsule biosynthesis (spot 41 and spot 42, respectively, in *Bp137*).

Human immunoproteomic data recently published (Zhu *et al.* 2010) include 16 of the 64 polypeptides here

identified, indicating that they are immunogenic (Table 3). Other ten were detected to be reactive against murine immune serum. Five of them were reactive against both sera. Three of the five are present in all the strains here included and correspond to well-known antigens of *Bord. pertussis*: 60-kDa chaperonin (spot 14), pertactin (spot 19) and serum resistance protein (spot 32) (Altindis *et al.* 2009; Zhu *et al.* 2010; Tefon *et al.* 2011) (Table 3). Other proteins such as BP1285, BP3642 and BP0844 are among the differential proteins here detected.

## Discussion

Here, we showed that the PFGE of the *Bord. pertussis* strain *Bp137* and two other strains included in wP vaccines in Latin America were distributed in three groups classified according to a criterion of similarity higher than 80%. Although this observation of the vaccine strain PFGE profiles is similar to that previously reported in other countries (Caro *et al.* 2005), it is still important for our region. The current PFGE classifies strains that were not studied before and that are currently included in the national immunization schedules of Latin American countries (e.g. the Brazilian vaccine strain *Bp137* and strain *Bp10536*, which is included in vaccines used in Argentina). The representative isolate *Bp106*, collected after the introduction of generalized vaccination in Argentina, is clearly separated from vaccine strains as we previously reported (Bottero *et al.* 2007).

**Table 3** Surface proteome of *Bordetella pertussis* vaccine strains and an Argentinean clinical isolate Bp106. Numbers in parentheses indicate corresponding spot number of Fig. 2

GI:	Gene locust	Localization†	Protein name/function	MW (kDa)	pI	Spot detection in strain (spot number in Bottero et al. 2007 or this work)				Murine serum reactive	Human serum reactive	Spot detection in strain (spot number in Bottero et al. 2007)*
						Bp137	Bp509	Bp10536	Tohama I			
33592278	<i>bp1146</i>	Outer membrane	Competence lipoprotein precursor	298	5.0	Yes(1)	Yes	Yes	Yes			Yes
33594323	<i>bp3441</i>	Cytoplasmic membrane	Conserved hypothetical protein	198	5.1	Yes(2)	Yes	Yes	Yes			Yes
33593636	<i>bp2667</i>	Outer membrane	Adhesin	263.6	9.7	Yes(3)	Yes	Yes	Yes			Yes
33592419	<i>bp1296</i>	Unknown	Putative lipoprotein	30.6	7.4	Yes(4)	Yes	Yes	Yes			Yes
33593352	<i>bp2360</i>	Not defined	Succinate dehydrogenase catalytic subunit	27.2	6.2	Yes(5)	Yes	Yes	Yes			Yes
33594289	<i>bp3405</i>	Outer membrane	Outer membrane protein OMPQ	39.1	5.7	Yes(6)	Yes	Yes	Yes			Yes
33592006	<i>bp0840</i>	Outer membrane	Outer membrane porin protein precursor	41.0	5.4	Yes(7)	Yes	Yes	Yes			Yes
33594616	<i>bp3757</i>	Cytoplasmic membrane	Putative ABC transport ATP binding protein	29.6	5.1	Yes(8)	Yes	Yes	Yes			Yes
33592580	<i>bp1487</i>	Periplasmic	Putative periplasmic solute binding protein	40.0	7.8	Yes(9)	Yes	Yes	Yes			Yes
33594215	<i>bp3322</i>	Periplasmic	Putative binding protein-dependent transport protein	40.9	6.9	No	No	Yes(10)	Yes(10)			Yes(10)
33592538	<i>bp1440</i>	Cytoplasmic membrane	Putative membrane protein	33.4	5.3	Yes(11)	Yes	Yes	Yes			Yes
33592121	<i>bp0965</i>	Cytoplasmic	Antioxidant protein	23.7	5.7	Yes(12)	Yes	Yes	Yes			Yes
33593418	<i>bp2434</i>	Periplasmic	Serine protease	52.1	8.8	Yes(13)	Yes	Yes	Yes	Yes		Yes
33594370	<i>bp3495</i>	Cytoplasmic	Chaperonin 60 kDa	57.4	4.9	Yes(14)	Yes	Yes	Yes	Yes		Yes
33594369	<i>bp3494</i>	Outer membrane	Serum resistance protein	103.3	7.1	Yes(15)	Yes	Yes	Yes	Yes		Yes
33591281	<i>bp0007</i>	Cytoplasmic	Elongation factor Tu	42.9	5.1	Yes(16)	Yes	Yes	Yes	Yes		Yes
33592409	<i>bp1285</i>	Periplasmic	Leu/ile/Val protein precursor	39.6	6.8	No	Yes(17)	No	Yes(17)	Yes		No
33594507	<i>bp3642</i>	Cytoplasmic	DNA direct RNA $\alpha$ subunit polymerase	36.1	5.7	Yes(18)	Yes	No	Yes	Yes		Yes
33592195	<i>bp1054</i>	Outer membrane	Pertactin	93.4	10.0	Yes(19)	Yes	Yes	Yes	Yes		Yes
3593200	<i>bp2196</i>	Outer membrane	Putative quino protein	40.0	8.7	Yes(20)	Yes	Yes	Yes	Yes		Yes
33594713	<i>bp3862</i>	Cytoplasmic membrane	Putative extracellular solute binding protein	57.3	9.7	No	Yes(21)	Yes(21)	Yes(21)	Yes(21)		No
33599458	<i>bb0468</i>	Periplasmic	Putative molybdopterin oxidoreductase	121.6	7.3	No	No	Yes(22)	No	Yes(22)		No

Table 3 (Continued)

GI:	Gene locus†	Localization‡	Protein name/function	MW (kDa)	pI	Spot detection in strain (spot number in Bottero et al. 2007 or this work)				Murine serum reactive	Human serum reactive	Spot detection in strain (spot number in Bottero et al. 2007)*
						Bp137	Bp509	Bp10536	Tohama I			
33592100	<i>bp0943</i>	Outer membrane	Outer membrane protein A precursor	20.9	9.2	Yes(23)	Yes	Yes	Yes			Yes
33593721	<i>bp2761</i>	Periplasmic	Superoxide dismutase	21.2	6.5	Yes(24)	Yes	Yes		Yes		Yes
33593496	<i>bp2513</i>	Periplasmic	Putative exported protein	34.9	10.2	Yes(25)	Yes	Yes				Yes
33591513	<i>bp0277</i>	Cytoplasmic membrane	Ubiquinol cytochrome C reductase iron sulfur subunit	22.8	5.2	Yes(26)	Yes	Yes				Yes
33593715	<i>bp2755</i>	Cytoplasmic/membrane	Putative exported protein	189.0	6.2	Yes(27)	Yes	Yes				No
33593423	<i>bp2439</i>	Cytoplasmic	3-oxoacyl-(acyl carrier protein) synthase	43.6	5.7	Yes(28)	No	No				Yes
33594649	<i>bp3794</i>	Unknown/multiple	Putative bacterial secretion system protein	29.4	6.8	Yes(29)	Yes	Yes		Yes		Yes
33593375	<i>bp2386</i>	Cytoplasmic	Enolase	45.9	4.5	Yes(30)	Yes	Yes				Yes
33593466	<i>bp2483</i>	Cytoplasmic membrane	Two-component sensor protein	97.4	8.7	Yes(31)	Yes	No				No
33594369	<i>bp3494</i>	Outer membrane	Serum resistance protein	103.3	7.1	Yes(32)	Yes	Yes		Yes		Yes
33593353	<i>bp2361</i>	Cytoplasmic membrane	Succinate dehydrogenase flavo subunit	64.8	6.5	Yes(33)	Yes	Yes		Yes		Yes
33563780	<i>bp2747</i>	Periplasmic	Putative ABC transport solute binding protein	40.6	6.5	Yes(34)	Yes	Yes		Yes		Yes
33594186	<i>bp3288</i>	Cytoplasmic	ATP synthase subunit B	50.5	4.7	Yes(35)	Yes	Yes		Yes		Yes
33592518	<i>bp1420</i>	Cytoplasmic	Elongation factor Ts	30.9	5.1	Yes(36)	Yes	Yes		Yes		Yes
33594046	<i>bp3125</i>	Cytoplasmic	Ribose phosphate pyrophosphokinase	34.1	5.1	Yes(37)	No	No				Yes
33592145	<i>bp0995</i>	Cytoplasmic	Dihydroliipoamide dehydrogenase	62.3	5.8	No	No	No				Yes(38)
33594049	<i>bp3128</i>	Unknown/multiple	Hypothetical protein	68.5	6.1	No	Yes(39)	Yes(39)				Yes(39)
33594387	<i>bp3515</i>	Cytoplasmic	Hypothetical protein	35.9	6.6	Yes(40)	Yes	Yes				Yes
33594071	<i>bp3150</i>	Cytoplasmic	Polysaccharide biosynthesis protein	46.7	5.6	Yes(41)	No	No				Yes
33592714	<i>bp1630</i>	Cytoplasmic	Capsular polysaccharide biosynthesis protein	37.3	5.5	Yes(42)	No	No				Yes
33594122	<i>bp0379</i>	Cytoplasmic	Putative L lactactodehydrogenase	37.2	5.8	No	No	No		Yes		Yes(43)

Table 3 (Continued)

GI:	Gene locus†	Localization‡	Protein name/function	MW (kDa)	pI	Spot detection in strain (spot number in Bottero et al. 2007 or this work)				Murine serum reactive	Human serum reactive	Spot detection in strain (spot number in Bottero et al. 2007)*
						Bp137	Bp509	Bp10536	Tohama I			
33593899	<i>bp2964</i>	Cytoplasmic	Hypothetical protein	48.5	6.2	No	Yes(44)	No	Yes(44)			Yes(44)
33592332	<i>bp1203</i>	Unknown	Hypothetical protein	42.7	6.0	No	Yes(45)	Yes(45)	Yes(45)			Yes(45)
33593419	<i>bp2435</i>	Periplasmic	Putative sigma factor regulatory protein	39.2	9.6	No	No	No	No			Yes(46)
33594122	<i>bp3215</i>	Cytoplasmic membrane	Enoyl-acyl carrier protein	27.6	5.8	Yes(47)	Yes	Yes	Yes			Yes
33593710	<i>bp2750</i>	Unknown	Lipoprotein	23.1	7.7	No	Yes(48)	Yes(48)	Yes(48)			Yes(48)
33594422	<i>bp3552</i>	Cytoplasmic	Alkyl hydroperoxide reductase	20.1	4.9	Yes(49)	Yes	Yes	Yes			Yes
33591361	<i>bp0102</i>	Periplasmic	Putative penicillin binding protein precursor	44.9	7.8	No	No	No	No		Yes	Yes(50)
33571906	<i>bp1126</i>	Cytoplasmic	2-oxoglutarate dehydrogenase complex, E3 component	50.3	6.3	Yes(51)	No	No	No			Yes
33592841	<i>bp1774</i>	Cytoplasmic	Trigger factor	47.5	4.9	Yes(52)	Yes	Yes	Yes			Yes
3593235	<i>bp2235</i>	Outer membrane	Putative type III secretion system	63.3	5.9	Yes(53)	No	No	No			Yes
33564503	<i>bp3559</i>	Not defined	Hypothetical protein	37.9	4.7	Yes(54)	Yes	Yes	Yes			Yes
33592552	<i>bp1455</i>	Cytoplasmic	Probable phosphoglycerate mutase 2	23.8	5.9	Yes(55)	Yes	Yes	Yes			Yes
33594004	<i>bp3077</i>	Outer membrane	Putative outer membrane protein	77.7	6.1	Yes(56)	Yes	Yes	Yes			Yes
39931027	<i>bp2499</i>	Cytoplasmic	Molecular chaperone DnaK	69.7	4.9	Yes(57)	Yes	Yes	Yes	Yes		Yes
33592578	<i>bp1485</i>	Extracellular	Putative membrane protein	51.6	6.8	Yes(58)	Yes	Yes	Yes			Yes
33592010	<i>bp0844</i>	Cytoplasmic	NADH dehydrogenase delta subunit	47.7	5.8	Yes(59)	Yes	No	Yes	Yes		Yes
33593704	<i>bp2744</i>	Not defined	Putative ABC transport protein, ATP binding component	29.1	6.3	Yes(60)	Yes	Yes	Yes			Yes
33592591	<i>bp1499</i>	Cytoplasmic	Glutathione synthetase	34.7	5.4	Yes(61)	No	No	No			Yes
33593453	<i>bp2470</i>	Cytoplasmic	Seryl-tRNA synthetase	50.0	5.4	Yes(62)	Yes	Yes	Yes			Yes
33592009	<i>bp0843</i>	Cytoplasmic	NADH dehydrogenase subunit C	24.1	5.1	No	Yes(63)	Yes(63)	Yes(63)			Yes(63)
33591314	<i>bp0047</i>	Cytoplasmic	Homoserine O-acetyltransferase	44.9	5.7	No	Yes(64)	No	Yes(64)			Yes(64)

\*Numbers in parentheses correspond to spot number in this work or from Bottero et al. (2007).

†Gene loci are named according to NCBI (<http://www.ncbi.nlm.nih.gov/>).‡Protein localization is as predicted by PSORT (<http://psort.nibb.ac.jp/>).



As expected, the above-mentioned *Bord. pertussis* wP vaccine strains contain the characteristic *ptxA*, *prn* and *fim3* gene alleles of the old *Bord. pertussis* strains (Fig. 1b) (Cassiday *et al.* 2000; Gzyl *et al.* 2001; Hardwick *et al.* 2002a; Fielt *et al.* 2003). The vaccine strains *Bp137* and *Bp509*, however, present different characteristics from those of the other vaccine strains: pertussis toxin promoter *ptxP2* and the allele *fim2-2* instead *ptxP1* and *fim2-1*. The *ptxP2* allele was found in the Netherlands at a frequency of 43% and in the United States at 29% during the prevaccination period. In the Netherlands, this allele was also detected during the 1999–2000 period, but at a very low frequency (0.003%). Bart *et al.* (2010) showed that strains that harbour this *ptxP2* allele represented a distinct lineage that diverged from other strains relatively early in the evolutive history of *Bord. pertussis*. The *ptxP2* and also *ptxP1* strains are nearly completely replaced in the late 1990s by the *ptxP3* strains. In the Netherlands, the increase in the frequency of *ptxP3* strains was associated with the resurgence of pertussis. The *ptxP3* strains produced more Ptx than the *ptxP1* strain, and epidemiological data suggest that *ptxP3* strains are more virulent. The *ptxP3* strains have spread worldwide, being the predominant allele in our country (Mooi *et al.* 2009; Bart *et al.* 2010).

Regarding circulating bacteria, we observed that *Bp106*, as well as the majority of the current members of our collection, contains *ptxP3*, *ptxA1*, *prn2*, *fim2-1* and *fim3-B* alleles (data not shown). The replacement of *ptxP1*, *ptxA2* or *ptxA4*, *prn1* or *prn7* strains by *ptxP3*, *ptxA1* and *prn2* strains in recent times is a global phenomenon that has been observed in other countries (van Gent *et al.* 2009; Kallonen and He 2009; Mooi 2010; Advani *et al.* 2011).

Regarding the *fim2* allele, 97% of the collection, including the *Bp106* representative strain, is *fim2-1*. This finding agrees with observations made in the UK, where *fim2-1* has been the prevalent allele since 1920, and in the Netherlands, where it has been the prevalent allele since 1965 (Van Loo and Mooi 2002; Packard *et al.* 2004). In relation to the *fim3* allele, vaccine strains included in our study have the variant A, which was found in 24% of the isolates of our collection. The representative local strain, *Bp106*, has the allele B, similar to 76% of the circulating bacteria. This finding agrees with results from Finland prior to 1999, Canada prior to 1990 and Russia prior to 1969, as all isolates in those countries at those times contained the variant A. Isolates obtained from those countries after those years contained the predominant allele B (Tsang *et al.* 2004; Kallonen and He 2009).

The Fim serotypes are Fim2 for Tohama I strain and Fim2,3 for *Bp10536*, *Bp137* and *Bp509*. In our study, the Fim serotype for *Bp106* and for 97% of clinical isolates

was Fim3. The serotype for these circulating bacteria correlated with observations in other populations where Fim3 is the most frequent (Tsang *et al.* 2004; Heikkinen *et al.* 2008; Kallonen and He 2009; Kurova *et al.* 2010; Zhang *et al.* 2010; Advani *et al.* 2011).

Regarding the proteomic analysis here performed, the 2-DE profile of *Bp137* revealed more than 90 protein spots from which 50 proteins were successfully identified (Fig. 2). Sixteen polypeptides from the total identified seem to be immunogenic in humans as it was recently published (Zhu *et al.* 2010).

Comparative analysis of the proteomes showed that *Bp137* and *Bp106* present seven proteins that are not detected in the other strains. One of these seven proteins is BP2235, which is a potential protein of TTSS (spot 52). This result is striking because previously we found that the TTSS is expressed in bacteria that have recently been in contact with the host, whereas in laboratory-adapted vaccine strains, this expression would not occur (Gaillard *et al.* 2011). In contrast to those findings, here we observed the expression of TTSS components in the vaccine strain *Bp137*, even when this strain is adapted to growth in laboratory conditions. This result suggests that the expression of TTSS in this strain is governed by a different regulatory mechanism than in other vaccine strains. Whatever the molecular mechanism, whose identification is not within the scope of this work, the expression of TTSS components in *Bp137* is a desirable feature in a vaccine strain, not only because the TTSS is immunogenic but also because it shares a property with circulating clinical isolates (Fennelly *et al.* 2008; Medhekar *et al.* 2008; Zongfu Wu *et al.* 2008).

Two other proteins identified only in *Bp106* and *Bp137* but not detected in the rest of the vaccine strains are BP3150, which is assigned to polysaccharide biosynthesis, and BP1630, which is assigned to capsule biosynthesis (spot 41 and spot 42, respectively, in *Bp137*). Bacterial capsules allow pathogens to evade host defences. The expression of capsule proteins in these strains, therefore, could indicate the need to overcome the host immune response induced by vaccination.

Our results show that, among the vaccine strains studied here, the strain *Bp137* is the one that shares the highest number of proteins detected in the surface proteome with the representative circulating bacteria *Bp106*. Interestingly, some of these common proteins have immunogenic properties. Based on these results and taking into account the previous reports showing that phenotypic and genotypic divergence between strains could have an impact in protection (King *et al.* 2001; Bottero *et al.* 2007), we suggest that vaccines containing *Bp137* could be appropriate to improve the control of pertussis in our region.

## Acknowledgements

This work was supported by Agencia Nacional de Promoción Científica y Tecnológica – ANCPyT and Comisión de Investigaciones Científicas de Buenos Aires – CICBA (Argentina) grants to DFH. DFH is a member of the Scientific Career of CICBA. DB and MEG have fellowships from Consejo Nacional de Investigaciones Científicas y Tecnológicas – CONICET. LB and MF have fellowships from ANCPyT.

## References

- Advani, A., Donnelly, D. and Hallander, H. (2004) Reference - system for characterization of *Bordetella pertussis* pulsed-field gel electrophoresis profiles. *J Clin Microbiol* **42**, 2890–2897.
- Advani, A., Gustafsson, L., Ahren, C., Mooi, F.R. and Hallander, H.O. (2011) Appearance of Fim3 and ptxP3-*Bordetella pertussis* strains, in two regions of Sweden with different vaccination programs. *Vaccine* **29**, 3438–3442.
- Altindis, E., Tefon, B.E., Yildirim, V., Ozcengiz, E., Becher, D., Hecker, M. and Ozcengiz, G. (2009) Immunoproteomic analysis of *Bordetella pertussis* and identification of new immunogenic proteins. *Vaccine* **27**, 542–548.
- Bart, M.J., van Gent, M., van der Heide, H.G., Boekhorst, J., Hermans, P., Parkhill, J. and Mooi, F.R. (2010) Comparative genomics of prevaccination and modern *Bordetella pertussis* strains. *BMC Genomics* **11**, 627.
- Borisova, O., Kombarova, S.Y., Zakharova, N.S., van Gent, M., Aleshkin, V.A., Mazurova, I. and Mooi, F.R. (2007) Antigenic divergence between *Bordetella pertussis* clinical isolates from Moscow, Russia, and vaccine strains. *Clin Vaccine Immunol* **14**, 234–238.
- Bottero, D., Gaillard, M.E., Fingerhann, M., Weltman, G., Fernandez, J., Sisti, F., Graieb, A., Roberts, R. *et al.* (2007) Pulsed-field gel electrophoresis, pertactin, pertussis toxin S1 subunit polymorphisms, and surfaceome analysis of vaccine and clinical *Bordetella pertussis* strains. *Clin Vaccine Immunol* **14**, 1490–1498.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248–254.
- Caro, V., Njamkepo, E., Van Amersfoort, S.C., Mooi, F.R., Advani, A., Hallander, H.O., He, Q., Mertsola, J. *et al.* (2005) Pulsed-field gel electrophoresis analysis of *Bordetella pertussis* populations in various European countries with different vaccine policies. *Microbes Infect* **7**, 976–982.
- Cassiday, P., Sanden, G., Heuvelman, K., Mooi, F., Bisgard, K.M. and Popovic, T. (2000) Polymorphism in *Bordetella pertussis* pertactin and pertussis toxin virulence factors in the United States, 1935–1999. *J Infect Dis* **182**, 1402–1408.
- Fennelly, N.K., Sisti, F., Higgins, S.C., Ross, P.J., van der Heide, H., Mooi, F.R., Boyd, A. and Mills, K.H. (2008) *Bordetella pertussis* expresses a functional type III secretion system that subverts protective innate and adaptive immune responses. *Infect Immun* **76**, 1257–1266.
- Fiett, J., Letowska, I., Gniadkowski, M. and Hryniewicz, W. (2003) The new strategy for allele identification of the genes coding for pertussis toxin subunit S1 (ptx S1) and pertactin (prn) in *Bordetella pertussis*. *J Microbiol Methods* **55**, 651–666.
- Gaillard, M.E., Bottero, D., Castuma, C.E., Basile, L.A. and Hozbor, D. (2011) Laboratory adaptation of *Bordetella pertussis* is associated with the loss of type three secretion system functionality. *Infect Immun* **79**, 3677–3682.
- van Gent, M., de Greeff, S.C., van der Heide, H.G. and Mooi, F.R. (2009) An investigation into the cause of the 1983 whooping cough epidemic in the Netherlands. *Vaccine* **27**, 1898–1903.
- Gzyl, A., Augustynowicz, E., van Loo, I. and Slusarczyk, J. (2001) Temporal nucleotide changes in pertactin and pertussis toxin genes in *Bordetella pertussis* strains isolated from clinical cases in Poland. *Vaccine* **20**, 299–303.
- Hardwick, T.H., Cassidy, P., Weyant, R.S., Bisgard, K.M. and Sanden, G.N. (2002a) Changes in predominance and diversity of genomic subtypes of *Bordetella pertussis* isolated in the United States, 1935 to 1999. *Emerg Infect Dis* **8**, 44–49.
- Hardwick, T.H., Plikaytis, B., Cassidy, P.K., Cage, G., Peppler, M.S., Shea, D., Boxrud, D. and Sanden, G.N. (2002b) Reproducibility of *Bordetella pertussis* genomic DNA fragments generated by XbaI restriction and resolved by pulsed-field gel electrophoresis. *J Clin Microbiol* **40**, 811–816.
- Heikkinen, E., Xing, D.K., Olander, R.M., Hytonen, J., Viljanen, M.K., Mertsola, J. and He, Q. (2008) *Bordetella pertussis* isolates in Finland: serotype and fimbrial expression. *BMC Microbiol* **8**, 162.
- van Hemert, P.A. (1969) Specific properties of acid precipitated pertussis vaccine. *Prog Immunobiol Stand* **3**, 297–301.
- Hozbor, D., Mooi, F., Flores, D., Weltman, G., Bottero, D., Fossati, S., Lara, C., Gaillard, M.E. *et al.* (2009) Pertussis epidemiology in Argentina: trends over 2004–2007. *J Infect* **59**, 225–231.
- Kallonen, T. and He, Q. (2009) *Bordetella pertussis* strain variation and evolution postvaccination. *Expert Rev Vaccines* **8**, 863–875.
- Kasuga, T., Nakase, Y., Ukishima, K. and Takatsu, K. (1954a) Studies on *Haemophilus pertussis*. III. Some properties of each phase of *H. pertussis*. *Kitasato Arch Exp Med* **27**, 37–47.
- Kasuga, T., Nakase, Y., Ukishima, K. and Takatsu, K. (1954b) Studies on *Haemophilus pertussis*. IV. Preventive potency of each phase organisms of *H. pertussis* in mice. *Kitasato Arch Exp Med* **27**, 49–55.
- Kasuga, T., Nakase, Y., Ukishima, K. and Takatsu, K. (1954c) Studies on *Haemophilus pertussis*. V. Relation between the phase of bacilli and the progress of the whooping-cough. *Kitasato Arch Exp Med* **27**, 57–62.

- King, A.J., Berbers, G., van Oirschot, H.F., Hoogerhout, P., Knipping, K. and Mooi, F.R. (2001) Role of the polymorphic region 1 of the *Bordetella pertussis* protein pertactin in immunity. *Microbiology* **147**, 2885–2895.
- Kurova, N., Njamkepo, E., Brun, D., Tseneva, G. and Guiso, N. (2010) Monitoring of *Bordetella* isolates circulating in Saint Petersburg, Russia between 2001 and 2009. *Res Microbiol* **161**, 810–815.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- van Loo, I.H., Heuvelman, K.J., King, A.J. and Mooi, F.R. (2002) Multilocus sequence typing of *Bordetella pertussis* based on surface protein genes. *J Clin Microbiol* **40**, 1994–2001.
- Lu, Z., Szafron, D., Greiner, R., Lu, P., Wishart, D.S., Poulin, B., Anvik, J., Macdonell, C. *et al.* (2004) Predicting subcellular localization of proteins using machine-learned classifiers. *Bioinformatics* **20**, 547–556.
- Medhekar, B., Shrivastava, R., Mattoo, S., Gingery, M. and Miller, J.F. (2008) *Bordetella* Bsp22 forms a filamentous type III secretion system tip complex and is immunoprotective *in vitro* and *in vivo*. *Mol Microbiol* **71**, 492–504.
- Mooi, F.R. (2010) *Bordetella pertussis* and vaccination: the persistence of a genetically monomorphic pathogen. *Infect Genet Evol* **10**, 36–49.
- Mooi, F.R., van Oirschot, H., Heuvelman, K., van der Heide, H.G., Gaastra, W. and Willems, R.J. (1998) Polymorphism in the *Bordetella pertussis* virulence factors P.69/pertactin and pertussis toxin in The Netherlands: temporal trends and evidence for vaccine-driven evolution. *Infect Immun* **66**, 670–675.
- Mooi, F.R., Hallander, H., Wirsing von Konig, C.H., Hoet, B. and Guiso, N. (2000) Epidemiological typing of *Bordetella pertussis* isolates: recommendations for a standard methodology. *Eur J Clin Microbiol Infect Dis* **19**, 174–181.
- Mooi, F.R., van Loo, I.H., van Gent, M., He, Q., Bart, M.J., Heuvelman, K.J., de Greeff, S.C., Diavatopoulos, D. *et al.* (2009) *Bordetella pertussis* strains with increased toxin production associated with pertussis resurgence. *Emerg Infect Dis* **15**, 1206–1213.
- Packard, E.R., Parton, R., Coote, J.G. and Fry, N.K. (2004) Sequence variation and conservation in virulence-related genes of *Bordetella pertussis* isolates from the UK. *J Med Microbiol* **53**, 355–365.
- Pereira, A., Pereira, A.S., Moreira-Filho, C.A., Bando, S.Y. and Tambourgi, D.V. (2005) Comparative analysis of a *Bordetella pertussis* patient isolated strain and classical strains used in the pertussis vaccine. *Vaccine* **23**, 4353–4358.
- Riley, M. (1993) Functions of the gene products of *Escherichia coli*. *Microbiol Rev* **57**, 862–952.
- Schouls, L.M., van der Heide, H.G., Vauterin, L., Vauterin, P. and Mooi, F.R. (2004) Multiple-locus variable-number tandem repeat analysis of Dutch *Bordetella pertussis* strains reveals rapid genetic changes with clonal expansion during the late 1990s. *J Bacteriol* **186**, 5496–5505.
- Stainer, D.W. and Scholte, M.J. (1970) A simple chemically defined medium for the production of phase I *Bordetella pertussis*. *J Gen Microbiol* **63**, 211–220.
- Tefon, B.E., Maass, S., Ozcengiz, E., Becher, D., Hecker, M. and Ozcengiz, G. (2011) A comprehensive analysis of *Bordetella pertussis* surface proteome and identification of new immunogenic proteins. *Vaccine* **29**, 3583–3589.
- Tsang, R.S., Lau, A.K., Sill, M.L., Halperin, S.A., Van Caesele, P., Jamieson, F. and Martin, I.E. (2004) Polymorphisms of the fimbria fim3 gene of *Bordetella pertussis* strains isolated in Canada. *J Clin Microbiol* **42**, 5364–5367.
- Van Loo, I.H. and Mooi, F.R. (2002) Changes in the Dutch *Bordetella pertussis* population in the first 20 years after the introduction of whole-cell vaccines. *Microbiology* **148**, 2011–2018.
- World Health Organization, W. (2010) Pertussis vaccines: WHO position paper. *Wkly Epidemiol Rec* **85**, 385–400.
- Wu, Z., Zhang, W. and Lu, C. (2008) Immunoproteomic assay of surface proteins of *Streptococcus suis* serotype 9. *FEMS Immunol Med Microbiol* **53**, 52–59.
- Zhang, L., Xu, Y., Zhao, J., Kallonen, T., Cui, S., Hou, Q., Li, F., Wang, J. *et al.* (2010) Effect of vaccination on *Bordetella pertussis* strains, China. *Emerg Infect Dis* **16**, 1695–1701.
- Zhu, Y.Z., Cai, C.S., Zhang, W., Guo, H.X., Zhang, J.P., Ji, Y.Y., Ma, G.Y., Wu, J.L. *et al.* (2010) Immunoproteomic analysis of human serological antibody responses to vaccination with whole-cell pertussis vaccine (WCV). *PLoS ONE* **5**, e13915.

## Supporting information

Additional Supporting Information may be found in the online version of this article:

**Figure S1** Proteome reference map of *Bordetella pertussis* vaccine strains Bp509, Tohama I, Bp10536 and clinical isolate Bp106.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.