

# Vibrational Properties of Bismutine, BiH<sub>3</sub>

Enrique J. Baran

Centro de Química Inorgánica (CEQUINOR/CONICET, UNLP), Facultad de Ciencias Exactas,  
Universidad Nacional de La Plata, C. Correo 962, 1900-La Plata, Argentina

Reprint requests to Prof. Dr. E. J. B.; Fax: (54) 0221 4259485; E-mail: baran@quimica.unlp.edu.ar

Z. Naturforsch. **58a**, 126–128 (2003); received January 7, 2003

Structural and vibrational-spectroscopic data of BiH<sub>3</sub> have recently been reported for the first time. On the basis of these data the force constants, mean amplitudes of vibration and thermodynamic functions have been calculated. Some comparisons with the structurally related SbH<sub>3</sub> molecule are made.

**Key words:** BiH<sub>3</sub>; SbH<sub>3</sub>; Force Constants; Mean Amplitudes of Vibration; Bond Properties; Thermodynamic Functions.

Bismutine, BiH<sub>3</sub>, is the least stable of the group-15 trihydrides. F. Paneth first detected it in minute traces in 1918 using a radiochemical technique involving <sup>212</sup>Bi<sub>2</sub>Mg<sub>3</sub> [1]. The best preparation method seems to be the one proposed by Amberger in 1961, based on the disproportionation of CH<sub>3</sub>BiH<sub>2</sub> at –45 °C [2]. This relatively complex synthesis has recently been successfully reproduced and allowed to obtain the vibrational and rotational spectra of BiH<sub>3</sub> [3]. Using the spectroscopic and structural data reported in this study, we have now performed the calculation of the most important vibrational properties and thermodynamic functions of this molecule. For comparative purposes, similar calculations have also been made for SbH<sub>3</sub>, the immediately adjacent lower-mass trihydride in group-15 of the Periodic System.

It is expected that on going from NH<sub>3</sub> to BiH<sub>3</sub> an increase of the p-orbital character in the three bonding orbitals occurs, generating a stepwise diminution of the respective bond angle [4]. This expectation is clearly fulfilled, as BiH<sub>3</sub> presents the lowest bond angle (90.48°) [3], even lower as that of SbH<sub>3</sub> (91.55°) [5]. This means that in the vibrational ground state BiH<sub>3</sub> is an oblate symmetric top molecule ( $I_a = I_b < I_c$ ) close to a spherical top ( $I_a = I_b = I_c$ ), even more than SbH<sub>3</sub>.

A parameter that can be used to quantify the spherical character of an oblate top molecule is the relation between the rotational constants  $C$  and  $B$ , i.e.,  $\gamma = (C - B_c)/1/2(C + B_c)$  [5]. Bismutine with  $\gamma = -0.015$  represents the most quasi-spherical oblate top so far investigated, whereas for stibine  $\gamma = -0.052$ .

Table 1. Force constants (mdyn/Å) of BiH<sub>3</sub> and SbH<sub>3</sub>.

	$f_r$	$f_{rr}$	$f_\alpha$	$f_{\alpha\alpha}$
BiH <sub>3</sub>	1.79	0.01	0.16	–0.003
SbH <sub>3</sub>	2.11	–0.002	0.19	–0.005

For the calculation of force constants the geometrical and vibrational data reported by Jerzembeck et al. [3] were used. **G**- and **F**-matrixes were constructed as detailed in [6] and a modified valence force field (MVFF) ( $F_{12} = F_{34} = 0$ ) was employed. Results of this calculation are shown in Table 1, in which  $f_r$  refers to the Bi-H bonds and  $f_\alpha$  to the H-Bi-H angle, whereas  $f_{rr}$  and  $f_{\alpha\alpha}$  refer to the bond/bond and angle/angle interactions, respectively.


A similar calculation was performed for SbH<sub>3</sub>. In this case the structural parameters and vibrational frequencies were taken from the paper of Fusina et al. [5] (cf. also [7]). The results are included in Table 1.

As can be seen, Sb-H bonds are somewhat stronger than Bi-H bonds, whereas the deformational constants are comparable. In both cases interaction constants are negligible. The potential energy distribution of both molecules showed that all the vibrations are highly characteristic, as expected for species with heavy central atoms.

With the now obtained results it is interesting to compare the bond characteristics of the full series of the trihydrides of group-15 elements. A plot of the force constants of the X-H bonds as a function of the atomic number is presented in Fig. 1 (the force constants of the three lighter trihydrides were taken from the book of Siebert [8]). From this figure it becomes

Table 2. Mean vibration amplitudes (in Å) of BiH<sub>3</sub> and SbH<sub>3</sub>.

Temperature (K)	BiH <sub>3</sub>		SbH <sub>3</sub>	
	$u_{\text{Bi-H}}$	$u_{\text{H...H}}$	$u_{\text{Sb-H}}$	$u_{\text{H...H}}$
0	0.0984	0.179	0.0944	0.171
100	0.0984	0.179	0.0944	0.171
200	0.0984	0.180	0.0944	0.171
298.16	0.0985	0.183	0.0944	0.173
300	0.0985	0.183	0.0944	0.173
400	0.0986	0.188	0.0945	0.177
500	0.0991	0.195	0.0948	0.183
600	0.1000	0.204	0.0954	0.190
700	0.1013	0.212	0.0963	0.197
800	0.1030	0.221	0.0976	0.204
900	0.1048	0.230	0.0991	0.212
1000	0.1069	0.239	0.1008	0.220

Fig. 1. X-H force constant  $f_r$  (in mdyn/Å) of the X-H bond as function of the atomic number ( $Z$ ) of the group-15 trihydrides.

evident that with increasing p-orbital character of the bonds, which occurs with increasing  $Z$ -values, a continuous diminution of the X-H bond strength takes place. The comparatively high value of ammonia is clearly supported by the involvement of the  $sp^3$  hybrid orbital from the N-atom in bonding, and this is the only trihydride of this group in which this hybrid orbital is used (NH<sub>3</sub> bond angle = 107° [4, 9]). Already in the next compound of the series, PH<sub>3</sub>, the s-contribution can be considered as practically negligible (PH<sub>3</sub> bond angle = 94° [4, 9]). In going from the second row elements to higher rows, the bonding overlap is mainly concentrated on the p orbitals, where the s electron density is concentrated into the lone pair regions [9].

Table 3. Statistical thermodynamic functions of BiH<sub>3</sub> and SbH<sub>3</sub> (values in cal/deg.mole).

$T(K)$	$C_p$	$(H^0 - H_0^0)/T$	$-(G^0 - H_0^0)/T$	$S^{00}$
<b>BiH<sub>3</sub>:</b>				
100	7.96	7.95	40.22	48.17
200	8.77	8.10	45.76	53.86
298.16	10.30	8.57	49.08	57.65
300	10.32	8.58	49.13	57.71
400	11.80	9.21	51.68	60.89
500	13.11	9.86	53.80	63.66
600	14.22	10.49	55.66	66.15
700	15.15	11.10	57.32	68.42
800	15.91	11.65	58.84	70.49
900	16.62	12.16	60.24	72.40
1000	17.02	12.62	61.55	74.17
1100	17.42	13.04	62.77	75.81
1200	17.74	13.42	63.92	77.34
<b>SbH<sub>3</sub>:</b>				
100	7.96	7.95	38.38	46.33
200	8.54	8.05	43.91	51.96
298.16	9.89	8.43	47.19	55.61
300	9.91	8.43	47.24	55.67
400	11.30	8.98	49.74	58.72
500	12.56	9.57	51.81	61.38
600	13.67	10.17	53.60	63.77
700	14.61	10.74	55.21	65.95
800	15.41	11.27	56.68	67.96
900	16.06	11.77	58.04	69.81
1000	16.60	12.23	59.30	71.53
1100	17.04	12.64	60.49	73.13
1200	17.41	13.03	61.61	74.63

For a wider characterization of the metal-hydrogen bonds the mean amplitudes of vibration of BiH<sub>3</sub> and SbH<sub>3</sub> were also calculated. These calculations were performed with the method of characteristic vibrations of Müller et al. [10] (cf. also [11, 12]). The obtained results, in the temperature range between 0 and 1000 K, are shown in Table 2.

The mean amplitudes for the metal-hydrogen bonds show the expected trend; i.e., they are lower in the case of SbH<sub>3</sub>, as this molecule presents the higher bond force constant. Interestingly, both hydrides present relatively low temperature dependencies, and this dependency is also lower in the case of stibine. On the other hand, and as expected, the calculated mean amplitudes are somewhat higher than those of AsH<sub>3</sub> [11], PH<sub>3</sub> [11] and NH<sub>3</sub> [13], which present stronger bond force constants (cf. Fig. 1). Regarding the amplitudes for the non bonded H...H pairs, they lie also slightly higher in bismutene.

Finally, we have calculated the thermodynamic functions, in a wide temperature range, for the ideal gaseous state (unit fugacity) using the rigid rotator,

harmonic oscillator approximation [14, 15]. The moments of inertia were calculated with the MOLDRAW program [16]. The symmetry number is 3.

The results, specific heat ( $C_p$ ), reduced enthalpy [ $(H^0 - H_0^0)/T$ ], reduced free enthalpy [ $(G^0 - H_0^0)/T$ ] and entropy ( $S^0$ ), in the temperature range between 100 and 1200 K, are presented in Table 3. A comparison of the tabulated data shows comparable values for the thermodynamic functions of both species which are, as expected, slightly higher in the case of BiH<sub>3</sub> in the whole temperature range.

To conclude, these calculations provide a first insight into the most important vibrational properties of BiH<sub>3</sub>, the heaviest group-15 trihydride, and show that this molecule follows very well the expected general trends, when compared with its lighter congeners [9].

#### Acknowledgements

This work was supported by the “Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina”. The author is a member of the Research Career of this organism.

- [1] N. N. Greenwood and A. Earnshaw, *Chemistry of the Elements*, 2<sup>nd</sup>. Ed. Butterworth-Heinemann, Oxford 1997.
- [2] E. Amberger, *Chem. Ber.* **94**, 1447 (1961).
- [3] W. Jerzembeck, H. Bürger, L. Constantin, L. Margulès, J. Demaison, J. Breidung, and W. Thiel, *Angew. Chem. Int. Ed.* **41**, 2550 (2002).
- [4] H. B. Gray, *Electrons and Chemical Bonding*, W. A. Benjamin, Inc., New York 1965.
- [5] L. Fusina, G. Di Lonardo, and P. De Natale, *J. Chem. Phys.* **109**, 997 (1998).
- [6] J. R. Ferraro and J. S. Ziomek, *Introductory Group Theory and its Application to Molecular Structure*, Plenum Press, New York 1969.
- [7] J. Breidung and W. Thiel, *J. Mol. Spectrosc.* **169**, 166 (1995).
- [8] H. Siebert, *Anwendungen der Schwingungsspektroskopie in der Anorganischen Chemie*, Springer, Berlin 1966.
- [9] D. M. P. Mingos, *Essential Trends in Inorganic Chemistry*, Oxford University Press, Oxford 1998.
- [10] A. Müller, C. J. Peacock, H. Schulze, and U. Heidborn, *J. Mol. Struct.* **3**, 252 (1969).
- [11] A. Müller, E. J. Baran, and K. H. Schmidt, *Characteristic Mean Amplitudes of Vibration*; in S. J. Cyvin (Ed.), *Molecular Structures and Vibrations*, Elsevier, Amsterdam 1972.
- [12] E. J. Baran, *An. Asoc. Quím. Argent.* **61**, 141 (1973).
- [13] S. J. Cyvin, *Molecular Vibrations and Mean Square Amplitudes*, Elsevier, Amsterdam 1968.
- [14] K. S. Pitzer, “*Quantum Chemistry*”, Prentice Hall, New Jersey 1953.
- [15] G. Allen and H. O. Pritchard, “*Statistical Mechanics and Spectroscopy*”, Butterworths, London 1974.
- [16] P. Ugliengo, D. Viterbo, and G. Chiari, *Z. Kristallogr.* **207**, 9 (1993).