

Review

# Stretch-elicited $\text{Na}^+/\text{H}^+$ exchanger activation: the autocrine/paracrine loop and its mechanical counterpart

Horacio E. Cingolani<sup>a,\*</sup>, Néstor G. Pérez<sup>a,1</sup>, Burket Pieske<sup>b</sup>, Dirk von Lewinski<sup>b</sup>,  
María C. Camilión de Hurtado<sup>a,1</sup>

<sup>a</sup>Centro de Investigaciones Cardiovasculares, Facultad de Ciencias Médicas, Universidad Nacional de La Plata, 1900 La Plata, Argentina

<sup>b</sup>Abteilung Kardiologie und Pneumologie, Universität Göttingen, Göttingen, Germany

Received 9 September 2002; accepted 4 November 2002

## Abstract

The stretch of the cardiac muscle is immediately followed by an increase in the contraction strength after which occurs a slow force increase (SFR) that takes several minutes to fully develop. The SFR was detected in a wide variety of experimental preparations including isolated myocytes, papillary muscles and/or trabeculae, left ventricle strips of failing human myocardium, in vitro isovolumic and in vivo volume-loaded hearts. It was established that the initial increase in force is due to an increase in myofilament  $\text{Ca}^{2+}$  responsiveness, whereas the SFR results from an increase in the  $\text{Ca}^{2+}$  transient. However, the mechanism(s) for this increase in the  $\text{Ca}^{2+}$  transient has remained undefined until the proposal of  $\text{Na}^+/\text{H}^+$  exchanger (NHE) activation by stretch. Studies in multicellular cardiac muscle preparations from cat, rabbit, rat and failing human heart have shown evidence that the stretch induces a rise in intracellular  $\text{Na}^+$  ( $[\text{Na}^+]_i$ ) through NHE activation, which subsequently leads to an increase in  $\text{Ca}^{2+}$  transient via reverse-mode  $\text{Na}^+/\text{Ca}^{2+}$  (NCX) exchange. These experimental data agree with a theoretical ionic model of cardiomyocytes that predicted an increased  $\text{Na}^+$  influx and a concurrent increase in  $\text{Ca}^{2+}$  entry through NCX as the cause of the SFR to muscle stretch. However, there are aspects that await definitive demonstration, and perhaps subjected to species-related differences like the possibility of an autocrine/paracrine loop involving angiotensin II and endothelin as the underlying mechanism for stretch-induced NHE activation leading to the rise in  $[\text{Na}^+]_i$  and reverse-mode NCX.

© 2003 European Society of Cardiology. Published by Elsevier Science B.V. All rights reserved.

**Keywords:** Angiotensin; Endothelins; Na/Ca-exchanger; Na/H-exchanger; Stretch/m-e coupling

## 1. Introduction

There are intrinsic mechanisms in the heart by which it can adjust cardiac output to changes in hemodynamic conditions. The increase in ventricular end-diastolic volume (EDV) caused either by an increase in venous return or a rise in aortic resistance is immediately followed by an increase in the strength of the heartbeat. This rapid adaptation, known as the Frank–Starling mechanism, allows cardiac output to match venous return or remain constant even when the heart faces higher afterloads. Next,

there is a further increase in force that takes several minutes to fully develop and allows the return of EDV towards baseline. These load- and time-dependent changes in heart contractility were reproduced in isolated ventricular strips by Parmley and Chuck in 1973 [1]. These authors showed that the sudden increase of cardiac muscle length leads to a rapid initial increase in twitch force followed by a second slower force increase over several minutes as exemplified in Fig. 1.

After the description by Parmley and Chuck [1] the SFR was detected in a wide variety of experimental preparations including isolated rat and guinea pig cardiomyocytes, [2,3] papillary muscles from cat, ferret and rabbit and rat trabeculae [4–9], in vitro isovolumic dog hearts [10,11], and in vivo volume-loaded canine hearts [12]. Although

\*Corresponding author. Tel.: +54-221-483-4833; fax: +54-221-425-5861.

E-mail address: [cicmes@infovia.com.ar](mailto:cicmes@infovia.com.ar) (H.E. Cingolani).

<sup>1</sup>Established investigators of Consejo Nacional de Investigaciones Científicas y Técnicas (Conicet) Argentina.

**Time for primary review 21 days.**




Fig. 1. Characteristic myocardial contractile response to a sudden increase in muscle length. After stretching a papillary muscle, there is an immediate increase in force (from a to b, panel A) due to an increase in myofilament  $Ca^{2+}$  responsiveness. After that, a progressive increase in force develops during the next 10–15 min, the SFR, that is due to an increase in the  $Ca^{2+}$  transient (panel B). This increase in  $Ca^{2+}$  transient is due to the increase in  $[Na^+]_i$  (panel D, solid circles) caused by  $Na^+/H^+$  exchange (NHE) activation since both the SFR and the increase in  $[Na^+]_i$  are abolished by blocking NHE activity (panels C and D, open squares). \* indicates  $P < 0.05$  vs. initial rapid phase. § indicates  $P < 0.05$  vs. pre-stretched control, # indicates  $P < 0.05$  between curves. (Adapted from Cingolani et al. [48] and from Alvarez et al. [6]).

one study in humans failed to detect the SFR after the elevation of EDV by intracardiac catheterization [13], the SFR was found to be present in failing human myocardium. Fig. 2 shows the characteristic biphasic force response to stretch in a left ventricular muscle strip isolated from an end-stage failing human heart. This

observation in human myocardium is very much alike the functional response to stretch of a cat papillary muscle shown in Fig. 1, suggesting that the SFR to stretch is an ubiquitous intrinsic response of mammalian myocardium to mechanical load, even present in failing human myocardium.




Fig. 2. The SFR is also present in failing human myocardium. A. Typical force record of the effect of stretch in a left ventricular muscle strip from an end-stage failing human heart. Stretch resulted in an immediate, followed by a slowly developing second phase in force increase. B. Overall results from eight independent experiments showing the relative contribution of the immediate force response vs. the SFR to the total force increase. (Pieske et al., unpublished results). \* indicates  $P < 0.05$  vs. 88%; # indicates  $P < 0.05$  vs 1st phase.

## 2. Mechanisms of the SFR to stretch

It has been established that the mechanisms that participate in each phase of the increase in force after stretch are quite different. There is general agreement that the initial rapid phase is due to an increase in myofilament  $\text{Ca}^{2+}$  responsiveness [14], with no change in the amount of  $\text{Ca}^{2+}$  delivered to the contractile elements. Instead, the SFR is the result of an increase in the  $\text{Ca}^{2+}$  transient (Fig. 1), a finding first reported by Allen and Kurihara in 1982 [4], and subsequently by several (including our) laboratories [2,5,6,10]. Even though the phenomenon received considerable attention, the mechanism responsible for the increase in the  $\text{Ca}^{2+}$  transient during the SFR was not completely defined and several alternatives were raised.

Increases in the  $\text{Ca}^{2+}$  transient might be accounted by increases in  $\text{Ca}^{2+}$  entry, decreases in  $\text{Ca}^{2+}$  efflux and/or changes in the sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$  handling. A main  $\text{Ca}^{2+}$  entry pathway in cardiac cells is the L-type  $\text{Ca}^{2+}$  current, which could be sensitive to stretch or affected by the prolongation of action potential duration induced by stretch [14,15]. So far, the experimental evidence is against a possible increase in  $\text{Ca}^{2+}$  entry through L-type channels as the cause of the augmented  $\text{Ca}^{2+}$  transients during the SFR. No change in L-type  $\text{Ca}^{2+}$  current after stretch was detected by Hongo et al. in isolated cardiomyocytes [2], nor was the SFR abolished by  $\text{Ca}^{2+}$  channel antagonists in the experiments by Chuck and Parmley performed in cat papillary muscles [16]. Besides the L-type  $\text{Ca}^{2+}$  current, stretch-activated ion channels might represent another potential  $\text{Ca}^{2+}$  entry pathway contributing to the enhancement of  $\text{Ca}^{2+}$  transient [17]. However, recent reports showed that gadolinium failed to modify the SFR in rabbit [8] and rat [18] papillary muscle.

Allen et al. [19] proposed the alternative that the increase of the  $\text{Ca}^{2+}$  transient might have resulted from an increase in  $\text{Ca}^{2+}$  influx that, increasing resting  $\text{Ca}^{2+}$  would load the SR and increase  $\text{Ca}^{2+}$  transients in subsequent beats.

The increase in  $\text{Ca}^{2+}$  binding to TnC immediately after the stretch [14] causes a decrease in  $\text{Ca}^{2+}$  transient. Trafford et al. [20] proposed that a decrease in  $\text{Ca}^{2+}$  transient causes a decrease in  $\text{Ca}^{2+}$  efflux that may load the SR. The greater loading of the SR will subsequently lead to the increase in the amplitude of the  $\text{Ca}^{2+}$  transient and force. Although a gain in the SR  $\text{Ca}^{2+}$  content was reported in rabbit myocardium [8,21] the studies of Kentish and Wrzosek [5], Bluhm and Levy [21], and Kentish et al. [22] would indicate that a functional SR is not a requirement for the development of the SFR since it can be elicited even in the presence of specific SR inhibitors. The appearance of an SFR in failing myocardium, a condition of severely depressed SR  $\text{Ca}^{2+}$ -ATPase activity [23,24], would also argue against a predominant role of the SR. In contrast, a report by Chuck and Parmley [16] described the reversal of the SFR in cat papillary

muscles by caffeine; but changes in myofilament  $\text{Ca}^{2+}$  responsiveness promoted by caffeine could have interfered with their results. Despite the above mentioned studies showing that a functional SR is not mandatory for the development of the SFR, it was shown that the time-course of the SFR was delayed after SR inhibition [4,13].

Other mechanisms proposed to probably mediate the increase in  $\text{Ca}^{2+}$  transient were stretch-induced changes in intracellular second messengers. Increases in cAMP after stretch were reported in isolated ferret papillary muscles [9], and intact canine hearts [10], whereas increased production of  $\text{InsP}_3$  was found in rat cardiomyocytes [25]. Also in isolated rat cardiomyocytes, Vila Petroff et al. [26] proposed that endogenous NO, released from stretched myocytes themselves, could act on ryanodine receptors, and thus enhance the SR  $\text{Ca}^{2+}$  releasing capacity. Though interesting, these results are difficult to reconcile with the aforementioned findings showing that the SFR is elicited even with non-functional SR.

## 3. Role of NHE and reverse-mode NCX in the development of the SFR

An alternative hypothesis for the increase in  $\text{Ca}^{2+}$  transient amplitude during the SFR emerged from the results in cat papillary muscles by Cingolani et al. [27] showing stretch-induced activation of the sarcolemmal  $\text{Na}^+/\text{H}^+$  exchanger (NHE). Although this early study was not focused on the contractile response to stretch, subsequent work by the same group demonstrated the implication of NHE activation by stretch in SFR development. Their results showed a marked increase in intracellular  $\text{Na}^+$  ( $[\text{Na}^+]_i$ ) with a time-course similar to, or slightly preceding, the SFR and both the rise in  $[\text{Na}^+]_i$  and SFR were abolished by NHE inhibitors (Fig. 1) [6,7]. Since the rise in  $[\text{Na}^+]_i$  changes the thermodynamic balance of the  $\text{Na}^+/\text{Ca}^{2+}$  exchange (NCX), the possibility of an increase in  $\text{Ca}^{2+}$  influx through reverse-mode ( $\text{Ca}^{2+}_{in} - \text{Na}^+_{out}$ ) NCX was proposed. The participation of this mechanism in the increase in  $\text{Ca}^{2+}$  transient and the generation of the SFR was further supported by the experiments of Pérez et al. [7] in cat papillary muscles. In these experiments, the authors demonstrated the suppression of the SFR after inhibition of the NCX either by extracellular  $\text{Na}^+$  deprivation or with the NCX blocker KB-R7943 (Fig. 3). The involvement of the NHE and reverse-mode NCX in the SFR elicited by stretch in rabbit myocardium is also reported in this Spotlight issue by von Lewinski et al. [8]. In accordance with these results, preliminary experiments revealed a clear dependency of SFR on NHE and reverse-mode NCX activation in failing human cardiac tissue (Pieske et al. unpublished observation). Therefore, the increase in  $[\text{Na}^+]_i$  induced by NHE activation is the underlying mechanism for the increase in  $\text{Ca}^{2+}$  transient and force. It may be




Fig. 3. Role of reverse-mode of NCX in the development of the SFR. The NCX operating in reverse-mode is a requirement for the development of the SFR since any maneuver inhibiting the reverse-mode NCX like extracellular  $\text{Na}^+$  removal (panels A and B, solid circles) or KB-R7943 (panel C, solid circles) cancelled the SFR. (Adapted from Pérez et al. [7]).

argued that NHE activation should cause intracellular alkalinization. However, it is not the case when bicarbonate-dependent mechanisms are operative [6] because the simultaneous activation of the  $\text{Na}^+$ -independent  $\text{Cl}^-/\text{HCO}_3^-$  exchanger minimizes the change in  $\text{pH}_i$  [28]. In the absence of bicarbonate NHE activation raises  $\text{pH}_i$ , and under this condition the SFR results from a combination of two mechanisms: the increase in  $[\text{Na}^+]_i$  leading to the increase in  $\text{Ca}^{2+}$  influx through reverse-mode NCX and the increase in myofilament  $\text{Ca}^{2+}$  responsiveness due to intracellular alkalosis. As a consequence, the magnitude of the SFR is almost doubled in the absence of bicarbonate and KB-R7943 reduces it by half [29].

A potential point of controversy could be how important

is the rise in  $[\text{Na}^+]_i$  for the development of the SFR. In light of the results in multicellular muscle preparations by Alvarez et al. [6] and Perez et al. [7] the increase in  $[\text{Na}^+]_i$  appears to be mandatory (Fig. 1). Instead Hongo et al. [2] showed that the stretch did not promote an increase in  $[\text{Na}^+]_i$  in isolated myocytes. It may be speculated that there is at least two mechanisms for the SFR: a  $[\text{Na}^+]_i$ -independent mechanism seen in isolated cardiomyocytes, and another one dependent on nonmyocyte cells of multicellular muscle preparations acting to increase  $[\text{Na}^+]_i$ . However, it should be noted that NHE and NCX have been found expressed colocalized along the transverse tubular system in cardiomyocytes [30,31]. Their close proximity probably enhances the effect of changes in  $[\text{Na}^+]_i$  able to drive the reverse-mode NCX. Interestingly, an elegant theoretical ionic model was recently used by Bluhm et al. [32] to analyze the changes in the parameters of sarcolemmal ion fluxes that reproduced the effect of step changes in cardiac muscle length. The results suggested that the slow change in force that follows a sudden increase in muscle length may be caused by length-induced step changes in sarcolemmal  $\text{Na}^+$  influx, leading to an increase in  $[\text{Na}^+]_i$  and a concurrent increase in systolic  $\text{Ca}^{2+}$  entry through the NCX. Therefore, the theoretical model and the experimental results are in good agreement about a role of NHE and reverse-mode NCX in the development of the SFR.

#### 4. Role of angiotensin II (Ang II) and endothelin in the SFR

The study by Cingolani et al. [27] in cat papillary muscles provided experimental evidence to support the participation of an autocrine/paracrine mechanism involving endogenous Ang II and endothelin in the activation of NHE and the SFR. Blockade of Ang II- $\text{AT}_1$  or endothelin  $\text{ET}_A$  receptors with selective inhibitors abolished stretch-induced NHE activation as well as the increase in the  $\text{Ca}^{2+}$  transient and the SFR (Fig. 4). Coincidentally, Calaghan and White recently reported that the blockade of endothelin  $\text{ET}_A$  receptors reduced the SFR by half in ferret papillary muscle [33], but they were unable to demonstrate a contribution of Ang II. In contrast, blockade of  $\text{AT}_1$  and  $\text{ET}_A$  receptors failed to affect the SFR in rabbit papillary muscle [8]. Therefore, although NHE activation seems to be responsible for the development of the SFR, the mechanism through which NHE is activated remains controversial.

A relevant aspect is to define, in relation to the proposed activation of the renin–angiotensin and endothelin systems, if it occurs in sequential steps of a single process or they are independently activated by stretch. There is an increasing body of evidence showing that many effects thought to be due to Ang II are actually the result of endogenous endothelin. The mediation of endothelin in




Fig. 4. The SFR is the mechanical counterpart of an autocrine/paracrine mechanism triggered by the stretch that involves AngII-endothelin release. Both the increase in Ca<sup>2+</sup> transient and the SFR to stretch are cancelled by the blockade of Ang II-AT<sub>1</sub> receptors with losartan (left panels) or endothelin ET<sub>A</sub> receptors with BQ 123 (right panels). (Adapted from Alvarez et al. [6]).

great variety of Ang II effects such as its hypertensive action [34], induction of hypertrophy in both cardiac myocytes exposed to mechanical stress [35–37] and transgenic hypertensive rats [38], and activation of membrane transport mechanisms [27,39], has been reported. In connection with this, a recent study by Aiello et al. [40] reported that the increase in outward NCX current (*I*<sub>NCX</sub>) produced by exogenous Ang II in isolated cat cardiomyocytes was cancelled by blocking either Ang II-AT<sub>1</sub> or ET-1 receptors (Fig. 5).

Another important aspect to be considered is the origin




Fig. 5. Ang II increases outward *I*<sub>NCX</sub> through AT<sub>1</sub> receptors in isolated myocytes. The effect of exogenous Ang II increasing outward *I*<sub>NCX</sub> is cancelled after the blockade of endothelin receptors with TAK044 indicating that Ang II-effect is mediated by endogenous endothelin. The results also indicate that cardiomyocytes themselves are at the same time, source and target of endothelin. (Adapted from Aiello et al. [39]).

of Ang II and endothelin in stretched myocardial preparations. Even though it is known that myocardial cells have local renin–angiotensin [41] and endothelin systems [42], and that the stretch stimulates the secretion of Ang II [36,43,44] and endothelin [44], most of the knowledge about the events that couple mechanical stress with intracardiac peptide secretion and/or production has come from studies performed in isolated neonatal myocytes. Papillary muscles are multicellular with cardiomyocytes being surrounded by various cells, mainly fibroblasts and endothelial cells, which also release endothelin [45]. Whether cardiomyocytes themselves or any other type of intramyocardial cells are the source of endothelin is unknown at present. In cat papillary muscle, Pérez et al. [7] demonstrated that the SFR was preserved after rendering nonfunctional vascular and endocardial endothelial cells, whereas Calaghan and White [33] showed that the removal of endocardial endothelial cells reversed the slow response to stretch. We do not have a clear explanation for the discrepancy besides the simplistic one about species-related differences. Pioneering studies by Sadoshima et al. [43] established that the mechanical stretch caused the release of preformed Ang II to the surrounding medium of cultured neonatal rat cardiomyocytes but not in nonmyocyte cultures. These authors also showed that Ang II acted as the initial stimulus of the stretch-induced hypertrophic response. Further elucidation of the process came from the experiments mentioned above by Ito et al. [35] who showed in cultured isolated myocytes that Ang II causes a PKC-dependent increase in prepro endothelin-1 mRNA levels and release of endothelin-1. The experiments by Aiello et al. previously cited [40] in isolated cat cardiomyocytes also provided evidence favoring the pro-

posal that myocytes themselves are both the source and the target of endothelin.

As a consequence of the peptide release, multiple downstream signal transduction pathways are activated, including phospholipases C, D and A<sub>2</sub>, as well as many types of protein kinases, such as PKC and MAP kinase cascades [43,44,46]. The study by Yamazaki et al. [44] also showed that whereas ion channels inhibitors like gadolinium, streptomycin, glibenclamide or CsCl did not have any inhibitory effect, NHE blockade could partially attenuate MAP kinase activation and protein synthesis induced by stretch. However, these investigators have called recently the attention to the fact that stretch can trigger myocyte growth through Ang II-independent pathways in addition to the autocrine/paracrine mechanism [46,47]. In this context, Cingolani and collaborators provided a link between the molecular events and contractile effects triggered by myocardial stretch as it is schematized in Fig. 6. The phenomenon of myocardial response to mechanical stress can, therefore, be viewed as a complex

mechanism, which through a combination of responses (increase in force and hypertrophy) meets the common end-point of how to face an increased hemodynamic load. However, we should consider that although the activation of NHE first proposed by Cingolani and collaborators [6,27] to be the mechanism underlying the SFR has been confirmed by other laboratories [8,44], the pathway for this activation remains controversial and perhaps it is subjected to species-dependent differences.

## 5. Conclusions

The SFR seems to be a general phenomenon intrinsic to the mammalian myocardium. This functional response appears to be mediated by stretch-dependent activation of NHE and a consecutive,  $[\text{Na}^+]_i$ -dependent  $\text{Ca}^{2+}$  transient increase mediated through reverse-mode NCX. This slowly developing inotropic response may serve as an additional physiological mechanism to adapt stroke volume to increases in hemodynamic load even in the failing human heart. However, a number of questions remain unanswered: is it the only pathway elicited by stretch with implication in cardiac hypertrophy; or, as proposed by Yamazaki et al., could there be parallel Ang II-dependent and independent mechanisms? If endothelin is involved, which subtype of receptors ( $\text{ET}_A$  or  $\text{ET}_B$ ) are involved? Is it necessary to stimulate both of them for the NHE activation? How does stretch cause the release of peptides, and which is the mechanosensor? How important are these mechanisms for determining cardiac hypertrophy? An attractive hypothesis would be that the chain of events leading to the increase in  $[\text{Ca}^{2+}]_i$  could elicit cardiac hypertrophy through  $\text{Ca}^{2+}$ -calmodulin–calcineurin-dependent pathways. The fact that the interruption of this chain of events by inhibition of NHE activity decreases cardiac hypertrophy [49,50] seems to support the hypothesis.




Fig. 6. Schematic representation of the autocrine/paracrine proposed loop. Upon myocardial stretch, preformed-stored Ang II is released from cardiomyocytes; this peptide either acting on the cardiomyocyte (autocrine mode, panel A) or other intramyocardial type of cells, namely fibroblasts and/or endothelial cells in a paracrine mode (panel B), will induce endothelin (ET) release. In turn, ET, through  $\text{ET}_A$  receptors will induce NHE activation, leading to an increase in  $[\text{Na}^+]_i$  and  $\text{Ca}^{2+}$  transient through reverse-mode NCX. (Adapted from Pérez et al. [7]).

## References

- [1] Parmley WW, Chuck L. Length-dependent changes in myocardial contractile state. *Am J Physiol* 1973;224:1195–1199.
- [2] Hongo K, White E, Le Guennec J, Orchard CH. Changes in  $[\text{Ca}^{2+}]_i$ ,  $[\text{Na}^+]_i$  and  $\text{Ca}^{2+}$  current in isolated rat ventricular myocytes following an increase in cell length. *J Physiol (Lond)* 1996;491:609–619.
- [3] White E, Boyett MR, Orchard CH. The effects of mechanical loading and changes of length on single guinea-pig ventricular myocytes. *J Physiol* 1995;482:93–107.
- [4] Allen DG, Kurihara S. The effects of muscle length on intracellular calcium transients in mammalian cardiac muscle. *J Physiol (Lond)* 1982;327:79–94.
- [5] Kentish JC, Wrzosek A. Changes in force and cytosolic  $\text{Ca}^{2+}$  concentration after length changes in isolated rat ventricular trabeculae. *J Physiol* 1998;506(2):431–444.
- [6] Alvarez BV, Pérez NG, Ennis IL, Camilión de Hurtado MC, Cingolani HE. Mechanisms underlying the increase in force and

- calcium transient that follows stretch of cardiac muscle: a possible explanation of the Anrep effect. *Circ Res* 1999;85:716–722.
- [7] Pérez NG, Camilión de Hurtado MC, Cingolani HE. Reverse mode of the  $\text{Na}^+/\text{Ca}^{2+}$  exchange following myocardial stretch. Underlying mechanism of the slow force response. *Circ Res* 2001;88:376–382.
  - [8] Von Lewinsky D, Stumme B, Maier LS, Luers C, Bers DM, Pieske B. Stretch-dependent slow force response in isolated rabbit myocardium is  $\text{Na}^+$  dependent. *Cardiovasc Res* 2003;57:1052–1061.
  - [9] Calaghan SC, Colyer J, White E. Cyclic AMP but not phosphorylation of phospholamban contributes to the slow inotropic response to stretch in ferret papillary muscle. *Pflügers Arch* 1999;437(5):780–782.
  - [10] Todaka K, Ogino K, Gu AG, Burkhoff D. Effect of ventricular stretch on contractile strength, calcium transient, and cAMP in intact canine hearts. *Am J Physiol Heart Circ Physiol* 1998;43:H990–H1000.
  - [11] Tucci PJF, Bregagnollo EA, Spadaro J, Cicogna AC, Ribiero MCL. Length dependence of activation studied in the isovolumic blood-perfused dog heart. *Circ Res* 1984;55:59–66.
  - [12] Lew WY. Mechanisms of volume-induced increase in left ventricular contractility. *Am J Physiol Heart Circ Physiol* 1993;265:H1778–H1786.
  - [13] Noble MIM, Drake-Holland AJ. Lack of importance of the slow component of the response of force to an increase in cardiac muscle length. *Clin Sci* 1994;87:547–551.
  - [14] Allen DG, Kentish JC. The cellular basis of the length-tension relation in cardiac muscle. *J Mol Cell Cardiol* 1985;17:821–840.
  - [15] Allen DG. On the relationship between action potential duration and tension in cat papillary muscle. *Cardiovasc Res* 1977;11:210–218.
  - [16] Chuck LH, Parmley WW. Caffeine reversal of length-dependent changes in myocardial contractile state in the cat. *Circ Res* 1980;47:592–598.
  - [17] Zeng T, Bett GC, Sachs F. Stretch-activated whole cell currents in adult rat cardiac myocytes. *Am J Physiol Heart Circ Physiol* 2000;278:H548–H557.
  - [18] Lamberts RR, van Rijen MHP, Sipkema P, Franssen P, Sys SU, Westerhof N. Coronary perfusion and muscle lengthening increase cardiac contraction: different stretch-triggered mechanisms. *Am J Physiol Heart Circ Physiol* 2002;283:H1515–H1522.
  - [19] Allen DG, Nichols CG, Smith GL. The effects of changes in muscle length during diastole on the calcium transient in ferret ventricular muscle. *J Physiol* 1988;406:359–370.
  - [20] Trafford AW, Diaz ME, Eisner DA. Coordinated control of cell  $\text{Ca}^{2+}$  loading and triggered release from the sarcoplasmic reticulum underlies the rapid inotropic response to increased L-type  $\text{Ca}^{2+}$  current. *Circ Res* 2001;88:195–201.
  - [21] Bluhm WF, Lew WYW. Sarcoplasmic reticulum in cardiac length-dependent activation in rabbits. *Am J Physiol Heart Circ Physiol* 1995;269:H965–H972.
  - [22] Kentish JC, Davey R, Largen P. Isoprenaline reverses the slow force responses to a length change in isolated rabbit papillary muscle. *Pflügers Arch* 1992;421:519–521.
  - [23] Pieske B, Kretschmann B, Meyer M, Holubarsch Ch, Posival H, Just H, Hasenfuss G. Alterations in intracellular calcium handling associated with the inverse force–frequency relation in human dilated cardiomyopathy. *Circulation* 1995;92:1169–1178.
  - [24] Pieske B, Maier LS, Bers DM, Hasenfuss G.  $\text{Ca}^{2+}$  handling and SR  $\text{Ca}^{2+}$  content in isolated failing and nonfailing human myocardium. *Circ Res* 1999;85:38–46.
  - [25] Dassouli A, Sulpice J, Roux S, Crozatier B. Stretch-induced inositol triphosphate and tetrakisphosphate production in rat cardiomyocytes. *J Mol Cell Cardiol* 1993;25:973–982.
  - [26] Vila Petroff MG, Kim SH, Pepe S, Dessy C, Marban E, Balligand JL, Sollott SJ. Endogenous nitric oxide mechanisms mediate the stretch dependence of  $\text{Ca}^{2+}$  release in cardiomyocytes. *Nat Cell Biol* 2001;3:867–873.
  - [27] Cingolani HE, Alvarez BV, Ennis IL, Camilión de Hurtado MC. Stretch-induced alkalinization of feline papillary muscle. An autocrine-paracrine system. *Circ Res* 1998;83:775–780.
  - [28] Camilión de Hurtado MC, Alvarez BV, Pérez NG, Ennis IL, Cingolani HE. Angiotensin II activates  $\text{Na}^+$ -independent  $\text{Cl}^-$ – $\text{HCO}_3^-$  exchanger. *Circ Res* 1998;82:473–481.
  - [29] Cingolani HE, Pérez NG, Ennis IL, Camilión de Hurtado MC.  $\text{Na}^+/\text{H}^+$  activation by myocardial stretch: an autocrine/paracrine loop. In: Karmazyn M, Avkiran M, Fliebel L, editors. The sodium–hydrogen exchanger. From molecule to its role in disease, Kluwer Academic Publishers, 2002 (in press).
  - [30] Yan Z, Pascarel C, Steele DS, Komukai K, Brette F, Orchard CH.  $\text{Na}^+/\text{Ca}^{2+}$  exchange activity is localized in the T-tubules of rat ventricular myocytes. *Circ Res* 2002;91:315–322.
  - [31] Petreca K, Atansiu R, Grinstein S, Orłowski J, Shrier A. Subcellular localization of the  $\text{Na}^+/\text{H}^+$  exchanger NHE1 in rat myocardium. *Am J Physiol Heart Circ Physiol* 1999;276:H709–H717.
  - [32] Bluhm WF, Lwe WY, Grafinkel A, McCulloch AD. Mechanism of length history-dependent tension in an ionic model of the cardiac myocyte. *Am J Physiol Heart Circ Physiol* 1998;274:H1032–H1040.
  - [33] Calaghan SC, White E. Contribution of angiotensin II, endothelin I and the endothelium to the slow inotropic response to stretch in ferret papillary muscle. *Pflügers Arch* 2001;441:514–520.
  - [34] Rajagopalan S, Laursen JB, Borthayre A, Kurs S, Keiser J, Haleen S, Giaid A, Harrison DG. Role for endothelin-1 in angiotensin II-mediated hypertension. *Hypertension* 1997;30:29–34.
  - [35] Ito H, Hirata Y, Adachi S, Tanaka M, Tsujino M, Koike A, Nogami A, Marumo F, Hiroe M. Endothelin-1 is an autocrine/paracrine factor in the mechanism of angiotensin II-induced hypertrophy in cultured rat cardiomyocytes. *J Clin Invest* 1993;92:398–403.
  - [36] Yamazaki T, Komuro I, Kudoh S, Zou Y, Shiojima I, Hiroi Y, Mizuno T, Maemura K, Kurihara H, Aikawa R, Takano H, Yazaki Y. Endothelin-1 is involved in mechanical stress-induced cardiomyocyte hypertrophy. *J Biol Chem* 1996;271:3221–3228.
  - [37] Gray MO, Long CS, Kalinyak JE, Li HT, Karliner JS. Angiotensin II stimulates cardiac myocyte hypertrophy via paracrine release of  $\text{TGF-}\beta_1$  and endothelin-1 from fibroblasts. *Cardiovasc Res* 1998;40:352–363.
  - [38] Zol O, Quattek J, Seeland U, El-Armouche A, Eschenhagen T, Böhm M. Activation of the cardiac endothelin system in left ventricular hypertrophy before onset of heart failure in TG(mREN2)27 rats. *Cardiovasc Res* 2002;53:363–371.
  - [39] Camilión de Hurtado MC, Alvarez BV, Ennis IL, Cingolani HE. Stimulation of myocardial  $\text{Na}^+$ -independent  $\text{Cl}^-$ – $\text{HCO}_3^-$  exchanger by angiotensin II is mediated by endogenous endothelin. *Circ Res* 2000;86:622–627.
  - [40] Aiello EA, Villa-Abrille MC, Cingolani HE. Autocrine stimulation of cardiac  $\text{Na}^+$ – $\text{Ca}^{2+}$  exchanger currents by endogenous endothelin released by angiotensin II. *Circ Res* 2002;90:374–376.
  - [41] Baker KM, Booz GW, Dostal DE. Cardiac actions of angiotensin II: role of an intracardiac renin–angiotensin system. *Ann Rev Physiol* 1992;54:227–241.
  - [42] Battistini B, Chailier P, Dorleans-Juste P, Briere N, Sirois P. Growth regulatory properties of endothelins. *Peptides* 1993;14:385–399.
  - [43] Sadoshima J, Xu Y, Slayter HS, Izumo S. Autocrine release of angiotensin II mediates stretch-induced hypertrophy of cardiac myocytes in vitro. *Cell* 1993;75:977–984.
  - [44] Yamazaki T, Komuro I, Kudoh S, Zou Y, Nagai R, Aikawa R, Uozumi H, Yazaki Y. Role of ion channels and exchangers in mechanical stress-induced cardiac hypertrophy. *Circ Res* 1998;82:430–437.
  - [45] Harada M, Itoh H, Nakagawa O, Ogawa Y, Miyamoto Y, Kuwahara K, Ogawa E, Igaki T, Yamashita J, Masuda I, Yoshimasa T, Tanaka I, Saito Y, Nakao K. Significance of ventricular myocytes and nonmyocytes interaction during cardiocyte hypertrophy: evidence for endothelin-1 as a paracrine hypertrophic factor from cardiac nonmyocytes. *Circulation* 1997;96:3737–3744.

- [46] Yamazaki T, Komuro I, Kudoh S, Zou Y, Shiojima I, Mizuno T, Takano H, Hiroi Y, Ueki K, Tobe K, Kadowaki T, Nagai R, Yazaki Y. Angiotensin II partly mediates mechanical stress-induced cardiomyocyte hypertrophy. *Circ Res* 1995;77:258–265.
- [47] Yamazaki T, Komuro I, Yazaki Y. Role of the renin–angiotensin system in cardiac hypertrophy. *Am J Cardiol* 1999;83:53H–57H.
- [48] Cingolani HE, Pérez NG, Camilión de Hurtado MC. An autocrine/paracrine mechanism triggered by myocardial stretch induces changes in contractility. *NIPS* 2001;16:88–91.
- [49] Camilión de Hurtado MC, Portiansky EL, Pérez NG, Rebolledo OR, Cingolani HE. Regression of cardiomyocyte hypertrophy in SHR following chronic inhibition of the Na<sup>+</sup>/H<sup>+</sup> exchanger. *Cardiovasc Res* 2002;53:862–868.
- [50] Kusumoto K, Haist JV, Karmazyn M. Na<sup>+</sup>/H<sup>+</sup> exchange inhibition reduces hypertrophy and heart failure after myocardial infarction in rats. *Am J Physiol Heart Circ Physiol* 2001;280:H738–745.