

NGC 4337: an overlooked old cluster in the inner disc of the Milky Way[★]

Giovanni Carraro,¹ † Edgard E. Giorgi,² Edgardo Costa³ and Ruben A. Vázquez²

¹ESO, Alonso de Cordova 3107, Casilla 19001, Santiago de Chile, Chile

²Facultad de Ciencias Astronómicas y Geofísicas (UNLP), Instituto de Astrofísica de La Plata (CONICET, UNLP), Paseo del Bosque s/n, B1900 La Plata, Argentina

³Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago de Chile, Chile

Accepted 2014 March 10. Received 2014 March 10; in original form 2014 February 27

ABSTRACT

Galactic open clusters do not survive long in the high-density regions of the inner Galactic disc. Inside the solar ring only 11 open clusters are known with ages older than 1 Gyr. We show here, basing on deep, high-quality photometry, that NGC 4337, contrary to earlier findings, is indeed an old open cluster. The cluster is located very close to the conspicuous star cluster Trumpler 20, as well misclassified in the past, and that has received so much attention in recent years. NGC 4337 shows a significant clump of He-burning stars which was not detected previously. Its beautiful colour–magnitude diagram is strikingly similar to the one of the classical old open clusters IC 4651, NGC 752, and NGC 3680, and this suggests similar age and composition. A spectroscopic study is much needed to confirm our findings. This, in turn, would also allow us to better define the inner disc radial abundance gradient and its temporal evolution. To this aim, a list of clump star candidates is provided.

Key words: open clusters and associations: general – open clusters and associations: individual: NGC 4337.

1 INTRODUCTION

Recent studies have consolidated a picture of a dual Milky Way disc. The corotation radius roughly divides the disc into two regions, the anticentre and the inner regions, that appear to have experienced very different chemical enrichment histories. Among the others, Magrini et al. (2010) and Frinchaboy et al. (2013) provided plenty of new data that lend strong support to the earlier findings by Twarog, Ashman & Anthony-Twarog (1997) that the radial abundance gradient is not monotonic over the whole disc extent. In agreement with observations in external spiral galaxies (Bresolin, Kennicutt & Ryan-Weber 2012), at about corotation, the radial abundance gradient flattens out in the anticentre region, while being quite steep in the inner regions. Most of these pieces of evidence come from studies of Galactic open clusters, especially the oldest ones (Friel 1995). These clusters are of invaluable importance since they allow us to trace the chemical properties of the disc since its very early assembly. Ideally, by using these clusters, one would be able to understand, e.g., how the abundance gradient built up and changed with time.

However, as widely known, they are very few in number, and mostly located in the disc periphery, where star clusters can survive

longer. In the inner disc, only 11 old open clusters are known, whose ages are larger than 1.0 Gyr (see Table 1). We caution the reader that this number is much smaller than what one can blindly extract from public catalogues, like for instance the WEBDA data base.¹ We restricted ourselves to clusters with a solid determination of the age, located between $l = 0^\circ$ and 85° in the first quadrant, and from $l = 275^\circ$ to 360° in the fourth quadrant, and within a Galactocentric distance smaller than the Sun value (8.5 kpc). For most clusters listed on WEBDA as older than 1 Gyr, either the available data or the analysis is poor (see the discussion, e.g., in Paunzen et al. 2010), and their age cannot be considered reliable enough.

This table contains also NGC 6791, one of the oldest and more metal rich old open clusters in the Milky Way. Its membership to the disc, however, has been questioned, since overall its parameters bear more similarity to the stellar populations in the Galactic bulge (Carraro 2013).

The most recent member of this restricted family is Trumpler 20. Originally misclassified as a young cluster, Trumpler 20 was then correctly found to be a rare example of a rich, old open cluster in the inner parts of the Milky Way disc (Platais et al. 2008; Seleznev et al. 2010). This triggered many follow-up studies (Donati et al. 2014, Carraro et al. 2014b, and references therein).

We report here on another very similar case: the old open cluster NGC 4337.

[★]Based on observations obtained at the Cerro Tololo Inter-American Observatory, Chile.

† On leave of absence from Padova University.

‡ E-mail: gcarraro@eso.org

¹ <http://www.univie.ac.at/webda>

Table 1. Star clusters in the inner disc older than 1.0 Gyr.

Cluster	l ($^{\circ}$)	b ($^{\circ}$)	Age (Gyr)	[Fe/H]	d_{GC} (kpc)	Reference
NGC 6583	9.28	-2.53	1.0	+0.37	6.5	Carraro, Méndez & Costa (2005)
Berkeley 44	53.21	+3.35	1.3		7.6	Carraro, Subramaniam & Janes (2006a)
Berkeley 52	67.89	-3.13	2.0		8.1	Carraro et al. (2006a)
NGC 6791	69.96	+10.90	8.0	+0.40	8.0	Carraro et al. (2006b)
NGC 6819	73.98	+8.48	3.0	+0.09	8.2	Carraro et al. (2014a)
NGC 3680	286.77	+16.92	1.85	-0.03	8.3	Anthony-Twarog & Twarog (2004)
Trumpler 20	301.47	+2.22	1.5	+0.09	7.3	Carraro et al. (2014b)
Collinder 261	301.68	-5.53	7.0	+0.13	7.5	Gozzoli et al. (1996)
NGC 6005	325.78	-2.98	1.2		6.5	Piatti et al. (1998)
NGC 6253	335.46	-6.25	3.0	+0.40	7.0	Piatti et al. (1998)
IC 4651	340.01	-7.91	1.7	+0.11	8.0	Meibom (2000)

This cluster has 2000.0 equinox equatorial coordinates $RA = 12^{\text{h}}24^{\text{m}}04^{\text{s}}$, $Dec. = -58^{\circ}07'24''$, while its Galactic coordinates are $l = 299^{\circ}.313$, $b = 4^{\circ}.556$. It is therefore located very close to Trumpler 20 in longitude but significantly higher on to the Galactic plane.

As a part of their southern sky search for star clusters, Moffat & Vogt (1973) describe NGC 4337 with the following words: In spatial order, the stars 9, 6, 10, 7, 11, 13, 14, 2, 4, 5, and 8 appear to form an open arc, revealing some similarity to the ‘stellar rings’ of Issersted (1968). However, the photometry shows no physical connection among these stars; they most likely represent a random arrangement of field stars. The actual cluster may be a small group (diameter ~ 2 arcmin) of stars located between nos 14 and 15, which are too faint to be reached by the 61 cm telescope except for the possible members 12, 13, 14, 1, and 15; the photometry of these also shows no physical association.

To help the reader, Moffat & Vogt (1973) map is reproduced in the left-hand panel of Fig. 1, while the right-hand panel shows the colour–magnitude diagram (CMD) that one can derive from their stars. This CMD gives full justice to their preliminary conclusions. Clearly, no hints are visible of the presence of a physical group, at odds with the photographic plate which gives the clear impression of some star concentration.

This latter evidence motivated us to acquire new CCD data. We briefly describe the observations in Section 2 of this Letter and use the photometric data set to probe the cluster structure in Section 3. Section 4 is devoted to the analysis of the CMDs of NGC 4337. Some conclusions are given in Section 5.

2 OBSERVATIONS

We took *UBVI* images of NGC 4337 in a 20×20 squared arcmin area during two observing runs, on 2010 Feb 14 and 2010 Mar 11, at Cerro Tololo Inter-American Observatory, using the 0.9 m and the 1.0 m telescopes. The first night (at the 0.9 m) was not photometric, and therefore, we tied all the images to the second night (at the 1.0 m), which was photometric. During this second night, we took multiple images of the standard star fields PG 1047 and SA98, covering an airmass range 1.15–2.2. Five frames per filter, with exposures ranging from 20 to 2400 s (*U*), 5 to 1800 s (*B*), and 5 to 1200 s (*V* and *I*) were obtained, to avoid bright star saturation and at the same time to get deep enough to unravel the cluster main sequence. These observations were part of a long-term project aimed at collecting high-quality data for a large open cluster data set. We refer the reader to previous publications (e.g. Carraro




Figure 1. Left-hand panel: image of NGC 4337 from Moffat & Vogt (1973). Numbers refer to observed stars, as described in the text. Right-hand panel: the CMD derived from Moffat & Vogt (1973) photometry.

Table 2. An excerpt of the photometric data presented in this Letter. The full table is posted at the CDS archive (<http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/MNRAS/>). The table lists all the clump star candidates.

ID	RA (2000.0) ($^{\circ}$)	Dec. (2000.0) ($^{\circ}$)	V (mag)	σ_V (mag)	$(B - V)$ (mag)	$\sigma_{(B - V)}$ (mag)	$(U - B)$ (mag)	$\sigma_{(U - B)}$ (mag)	$(V - I)$ (mag)	$\sigma_{(V - I)}$ (mag)
90	186.065 67	-58.147 27	13.804	0.002	1.275	0.003	1.101	0.005	1.311	0.004
91	186.163 58	-58.130 47	13.827	0.002	1.308	0.002	1.199	0.006	1.404	0.003
99	186.164 51	-58.222 29	13.853	0.002	1.304	0.004	1.254	0.010	1.406	0.004
100	185.863 45	-58.047 29	13.861	0.009	1.347	0.020	9.999	9.999	1.449	0.024
106	186.013 14	-58.111 06	13.867	0.002	1.277	0.002	1.007	0.005	1.348	0.004
107	186.024 49	-58.120 87	13.868	0.002	1.273	0.003	1.095	0.005	1.336	0.004
110	185.988 37	-58.120 68	13.908	0.002	1.275	0.003	1.127	0.004	1.322	0.004
114	186.037 14	-58.115 92	13.928	0.002	1.284	0.002	1.101	0.005	1.336	0.004
115	185.986 09	-58.123 67	13.933	0.002	1.320	0.003	1.180	0.005	1.365	0.005
117	185.968 57	-58.115 66	13.961	0.002	1.291	0.003	1.123	0.006	1.332	0.003
118	185.983 85	-58.110 50	13.966	0.002	1.294	0.003	1.115	0.005	1.353	0.004
121	186.036 66	-58.037 73	13.977	0.002	1.311	0.003	1.101	0.010	1.378	0.004
124	186.132 44	-58.133 92	14.017	0.001	1.280	0.002	1.135	0.006	1.321	0.003
128	186.187 27	-58.036 10	14.056	0.004	1.332	0.006	1.155	0.014	1.436	0.006
130	186.020 41	-58.086 07	14.061	0.002	1.330	0.003	1.190	0.008	1.365	0.004

& Costa 2007; Carraro & Seleznev 2011) for all the details of data reduction and photometric calibration. The final catalogue, cross-correlated with 2MASS, contains 8198 entries and is made available at the Centre de Données Astronomiques de Strasbourg (CDS) data base. An excerpt of the catalogue is presented in Table 2, which lists the clump star candidates.

3 CLUSTER STRUCTURE AND RADIUS

In Fig. 2, we show a contour plot in the area of NGC 4337, derived from our photometry. Star counts have been performed using a grid and density inside each field computed via a suitable kernel estimate (see Vázquez et al. 2010 for more details). The cluster has quite an irregular shape but certainly appears as a prominent overdensity. The density peak (~ 65 stars arcmin^{-2}) is offset with respect to the centre of the field, which was chosen according to the cluster




Figure 2. A density contour map in the area of NGC 4337. The map is 20 arcmin on a side, North is up, East to the left. The white circle indicates the estimate cluster area, the radius being 6 arcmin.

nominal centre. A visual inspection of Fig. 2 provides us with an estimate of the cluster radius of ~ 7 arcmin (see the superimposed white circle), larger than visual inspection on the digital sky survey (DSS; Dias et al. 2002).

Our quantitative star count analysis fully supports Moffat & Vogt (1973) visual impression that NGC 4337 is clearly a star overdensity with respect to the surrounding field.

4 ANALYSIS OF THE PHOTOMETRIC DIAGRAMS

The CMD for all the stars in the observed field is shown in Fig. 3. This strikingly resembles the CMD of an intermediate-age/old open cluster. Although field star contamination is significant, a turn off (TO) point can be easily identified at $V \sim 15.50$, together with a prominent clump of stars at $V \sim 14.0$, $U - B \sim 1.10$, $B - V \sim 1.30$, and $V - I \sim 1.35$. This diagram, in tandem with start counts, confirms beyond any reasonable doubt that NGC 4337 is indeed a physical star cluster.

To alleviate field star contamination, we make use of the previous section results and consider in the following only stars within 5 arcmin from the cluster centre. The resulting CMD in the $V/B - V$ plane is presented in Fig. 4. The cluster main features emerge very neatly and should immediately remind the reader the CMD of the more famous old open clusters IC 4651 and NGC 3680 (see Table 1),




Figure 3. CMDs for all the stars observed in the field of NGC 4337.




Figure 4. CMDs for all the stars within NGC 4337 radius. The red line shows IC 4651 ridge line offset by 0.14 in colour and 2.6 in magnitude.

or NGC 752. The TO point is located at $V = 15.50$, and the main sequence (MS) extends down to $V = 19$, in spite of some residual field star contamination. A scattered sequence of binary stars is visible to the right of the MS. The binary sequence intersects the MS at $V \sim 15.1$. We cannot exclude the presence of a few blue straggler stars.

Interestingly enough, the MS TO region has a curved shape, typical of intermediate-age/old clusters. Finally, the Hertzsprung gap is quite evident, together with the red giant clump, made of a dozen stars.

We stress also the fact that NGC 4337 CMD is much cleaner than Trumpler 20 CMD (Seleznev et al. 2010).

To infer for the first time the cluster basic parameters, we superimposed on the cluster CMD the ridge line of IC 4651 (red dashed line), taken from the study of Anthony-Twarog et al. (1988). IC 4651 seems to be the best choice, being a classical old open cluster, with an age around 1.7 Gyr (Anthony-Twarog et al. 1988; Meibom 2000; Anthony-Twarog et al. 2009) and a metallicity $[Fe/H] = +0.11$ (Carretta et al. 2004). This is slightly supersolar, as expected for clusters in the inner disc.

The fit has been performed by shifting the IC 4651 ridge line by 2.6 mag in magnitude and 0.14 mag in colour. The fit is really impressive as far as the MS is concerned. The only difference is in the magnitude of clump, since IC 4651 mean clump magnitude is ~ 0.3 mag brighter than NGC 4337. Since the fit is very convincing, we conclude that the two clusters most probably share the same metallicity, but NGC 4337 would be somewhat younger than IC 4651. IC 4651 reddening and apparent distance modulus are 0.12 and 10.40 mag, respectively, according to the recent study by Anthony-Twarog et al. (2009). This implies that NGC 4337 has a

reddening $E(B - V) \sim 0.26$ and an apparent distance modulus of ~ 13.00 . From these figures we derive a preliminary cluster heliocentric distance of 2.8 kpc, which, in turn, implies a distance of ~ 7.5 kpc from the Galactic Centre.

5 SUMMARY AND CONCLUSIONS

In this work, we have revisited the star cluster NGC 4337, previously considered as a random arrangement of field stars. One of the main features of the cluster CMD is an extremely well defined MS TO region, resembling closely IC 4651, NGC 752, and NGC 3680, all extremely well known textbook intermediate-age/old open clusters. NGC 4337 harbours a dozen clump stars, which would make it as massive as IC 4651, and most probably in the same dynamical stage (Meibom 2000). In fact, we do not see any manifestation of low-mass stars depletion, as for NGC 3680 and NGC 752.

The cluster is therefore a rare example of an old open cluster in the inner disc. According to our preliminary estimates of the cluster parameters, its Galactocentric distance would make NGC 4337 one of the more distant cluster from the Sun among the well-known old open clusters located inside the solar ring. This is a privileged location to establish in a more solid way the slope of the inner disc radial abundance gradient (Magrini et al. 2010) and its evolution with time. We expect this study to prompt a spectroscopic campaign to derive its metal abundance, and better constrain its fundamental parameters.

ACKNOWLEDGEMENTS

We thank the CDS, the US Naval Observatory and NASA for the use of their electronic facilities, especially SIMBAD, ViZier and ADS, and the WEBDA data base. GC thanks Nikolaus Vogt for granting the permission to reproduce his original plot in the left-hand panel of Fig. 1 of this Letter. EC acknowledges support by the Fondo Nacional de Investigacin Cient'fica y Tecnolgica (project no. 1110100 Fondecyt) and the Chilean Centro de Excelencia en Astrof'isica y Tecnolog'as Afines (PFB06). We finally thank the referee, Bruce Twarog, for his comments.

REFERENCES

- Anthony-Twarog B. J., Twarog B. A., 2004, *AJ*, 127, 1000
- Anthony-Twarog B. J., Mukherjee K., Twarog B. A., Caldwell N., 1988, *AJ*, 95, 1453
- Anthony-Twarog B. J., Delyinannis C. P., Twarog B. A., Croxall K. W., Cummings J. D., 2009, *AJ*, 138, 1171
- Bresolin F., Kennicutt R. C., Ryan-Weber E., 2012, *ApJ*, 750, 122
- Carraro G., 2013, *ASP Conf. Ser. preprint (arXiv:1308.5195)*
- Carraro G., Costa E., 2007, *A&A*, 464, 573
- Carraro G., Méndez R. A., Costa E., 2005, *MNRAS*, 356, 647
- Carraro G., Seleznev A. F., 2011, *MNRAS*, 412, 1361
- Carraro G., Subramaniam A., Janes K. A., 2006a, *MNRAS*, 371, 1301
- Carraro G., Villanova S., Demarque P., McSwain M. V., Piotto G., Bedin L., 2006b, *ApJ*, 643, 1151
- Carraro G., Buzzoni A., Bertone E., Buson L., 2014a, *AJ*, 146, 128
- Carraro G., Villanova S., Monaco L., Beccari G., Ahumada J. A., Boffin H. M. J., 2014b, *A&A*, 562, 39
- Carretta E., Bragaglia A., Gratton R. G., Tosi M., 2004, *A&A*, 422, 951
- Dias W. S., Alessi B. S., Moitinho A., Lépine J. R. D., 2002, *A&A*, 389, 871
- Donati P. et al., 2014, *A&A*, 516, 94
- Friel E. D., 1995, *ARA&A*, 33, 381
- Frinchaboy P. M. et al., 2013, *ApJ*, 777, L1
- Gozzoli E., Tosi M., Marconi G., Bragaglia A., 1996, *MNRAS*, 283, 66

- Issersted J., 1968, Voeroeffentl. Aston. Inst. d. Ruhr-Univ. Bochum no. 1
Magrini L., Randich S., Zoccali M., Jilkova L., Carraro G., Galli D., Maiorca E., Busso M., 2010, A&A, 523, 11
Meibom S., 2000, A&A, 361, 929
Moffat A. F. J., Vogt N., 1973, A&AS, 10, 135
Paunzen E., Heiter U., Netopil M., Soubiran C., 2010, A&A, 517, 32
Piatti A., Claría J. J., Bica E., Geisler D., Minniti D., 1998, AJ, 116, 801
Platais I., Melo C., Fulbright J. P., Kozhurina-Platais V., Figueira P., Barnes S. A., Méndez R. A., 2008, MNRAS, 391, 1482
Seleznev A., Carraro G., Costa E., Loktin A., 2010, New Astron., 15, 61
Twarog B., Ashman K. M., Anthony-Twarog B., 1997, AJ, 114, 2556
Vázquez R. A., Moitinho A., Carraro G., Dias W. S., 2010, A&A, 511, 38

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.