

Soil Metagenomes from Different Pristine Environments of Northwest Argentina

Christina B. McCarthy,^{a,b} Déborah I. Colman^{a,b*}

Centro Regional de Estudios Genómicos, Facultad de Ciencias Exactas, Universidad Nacional de la Plata, La Plata, Buenos Aires, Argentina^a; Departamento de Informática y Tecnología, Escuela de Ciencias Agrarias, Naturales y Ambientales, Universidad Nacional del Noroeste de la Provincia de Buenos Aires, Pergamino, Buenos Aires, Argentina^b

* Present address: Déborah I. Colman, Instituto Antártico Argentino, Dirección Nacional del Antártico, Ciudad Autónoma de Buenos Aires, Argentina.

This is the first study to use a high-throughput metagenomic shotgun approach to explore the biosynthetic potential of soil metagenomes from different pristine environments of northwest Argentina. Our data sets characterize these metagenomes and provide information on the possible effect these ecosystems have on their diversity and biosynthetic potential.

Received 9 July 2015 Accepted 10 July 2015 Published 13 August 2015

Citation McCarthy CB, Colman DI. 2015. Soil metagenomes from different pristine environments of northwest Argentina. *Genome Announc* 3(4):e00926-15. doi:10.1128/genomeA.00926-15.

Copyright © 2015 McCarthy and Colman. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christina B. McCarthy, mccarthychristina@gmail.com.

Soil microbiota produce many of the most important pharmaceutical drugs, including antibiotics and cancer drugs (1). Nevertheless, the traditional approach for characterizing the biosynthetic capacity of environmental bacteria, i.e., culturing them in the laboratory, has provided access to only a small fraction of this potential (2, 3). Recent analyses of soil microbiomes from around the world revealed a vastly unexplored biosynthetic diversity which was associated with soil types (4–7). In general, arid soils showed the richest biosynthetic diversity (5) and, similarly, bacterial diversity was highest in neutral soils (generally arid and semiarid ecosystems) and lower in acidic soils (generally tropical forest ecosystems) (7). The purpose of this study was to characterize soil metagenomes from different pristine environments using a metagenomic shotgun approach, giving special emphasis to the biosynthetic potential of each soil type. For this, four soil samples collected in northwest (NW) Argentina were analyzed. Sampling sites were chosen at different altitudes from the Yungas (YU) and Argentine Northwest Monte and Thistle of the Prepuna (NWMT) regions, with soils of varying pHs, namely: 1) YU (Montane Forest District) at 1,500 m above sea level (MASL) in Tafi del Valle (Tucumán, Argentina) (named Soil_TV; S27°01.123'; W65°39.807'; pH 5.35); 2) YU (Montane Cloud-forest District) at 850 MASL in Rosario de la Frontera (Salta, Argentina) (Soil_RF; S25°50.143' W64°55.524'; pH 8.01); 3) NWMT at 1,600 MASL in Cafayate (Salta) (Soil_CA; S26°03.885' W65°56.506'; pH 7.05); and 4) NWMT at 1,600 MASL in Quebrada de las Conchas (Cafayate Department, Salta) (Soil_QC; S26°01.123' W65°49.429'; pH 8.92). For the extraction of DNA, the three samples that contained more organic material (Soil_TV, Soil_RF, and Soil_CA) were processed with the QIAamp stool minikit (Qiagen), whereas Soil_QC was processed according to reference 8, treated with RNase (Invitrogen), and precipitated with LiCl and ethanol. High-throughput pyrosequencing of the samples was performed using a Roche GS FLX (Macrogen, Inc.,

South Korea), yielding ~1.15 Gb of metagenomic reads with lengths of 40 to 1,074 bases (nt) (520 nt average).

Raw sequence reads were trimmed using a custom application for removing nucleotides derived from the amplification primers (9, 10), and then processed with CD-HIT-454 (11). The nonredundant protein sequence NCBI database (DB:nr) was downloaded locally, and RAPSearch2 (12) was used to perform the protein homology search of the trimmed clustered reads against DB:nr. The taxonomic and functional content of the data sets was then analyzed with MEGAN (13, 14). Metagenomes consisted of 65.6% to 61.5% bacteria, 1.9% to 0.36% archaea, 1.6% to 0.17% eukaryota, and 0.1% to 0.01% viruses. Statistical analysis ($P < 0.05$, Fisher's exact test [15]) indicated significant differences between all samples. Diversity (Shannon-Weaver index) was highest in Soil_CA, followed by Soil_RF and Soil_TV, whereas Soil_QC showed the lowest diversity.

This is the first study to use a metagenomic shotgun approach to generate soil metagenome data sets from different pristine environments of NW Argentina. These data sets indicate the presence of bacteria, archaea, eukaryota, and viruses in all the samples and provide information on the potential effects of ecosystem types (including pH and altitude) on the composition, diversity, and biosynthetic potential of these soil metagenomes.

Nucleotide sequence accession numbers. Nucleotide sequences were submitted to the NCBI Sequence Read Archive (SRA) under the accession numbers [SRX1058163](https://www.ncbi.nlm.nih.gov/sra/SRX1058163), [SRX1058164](https://www.ncbi.nlm.nih.gov/sra/SRX1058164), [SRX1058165](https://www.ncbi.nlm.nih.gov/sra/SRX1058165) and [SRX1058166](https://www.ncbi.nlm.nih.gov/sra/SRX1058166).

ACKNOWLEDGMENTS

This research was supported by Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) (PICT PRH 112), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) (PIP 0294), and Universidad Nacional del Noroeste de la Provincia de Buenos Aires (UNNOBA) (Exp. 1388/2010 and Exp. 2581/2012) grants to C.B.M.

We gratefully acknowledge Eduardo Virla, who donated his time to help with soil sample collection.

C.B.M. is a member of the CONICET research career. D.I.C. was the recipient of an ANPCyT-UNNOBA fellowship when she participated in this work.

REFERENCES

1. Cragg GM, Newman DJ. 2013. Natural products: a continuing source of novel drug leads. *Biochim Biophys Acta* 1830:3670–3695. <http://dx.doi.org/10.1016/j.bbagen.2013.02.008>.
2. Rappé MS, Giovannoni SJ. 2003. The uncultured microbial majority. *Annu Rev Microbiol* 57:369–394. <http://dx.doi.org/10.1146/annurev-micro.57.030502.090759>.
3. Gilbert JA, Dupont CL. 2011. Microbial metagenomics: beyond the genome. *Annu Rev Mar Sci* 3:347–371. <http://dx.doi.org/10.1146/annurev-marine-120709-142811>.
4. Reddy BV, Kallifidas D, Kim JH, Charlop-Powers Z, Feng Z, Brady SF. 2012. Natural product biosynthetic gene diversity in geographically distinct soil microbiomes. *Appl Environ Microbiol* 78:3744–3752. <http://dx.doi.org/10.1128/AEM.00102-12>.
5. Charlop-Powers Z, Owen JG, Reddy BV, Ternei MA, Brady SF. 2014. Chemical-biogeographic survey of secondary metabolism in soil. *Proc Natl Acad Sci U S A* 111:3757–3762. <http://dx.doi.org/10.1073/pnas.1318021111>.
6. Charlop-Powers Z, Owen JG, Reddy BV, Ternei MA, Guimarães DO, de Frias UA, Pupo MT, Seepe P, Feng Z, Brady SF. 2015. Global biogeographic sampling of bacterial secondary metabolism. *Elife* 4:e05048. <http://dx.doi.org/10.7554/eLife.05048>.
7. Fierer N, Jackson RB. 2006. The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci U S A* 103:626–631. <http://dx.doi.org/10.1073/pnas.0507535103>.
8. Aldrich J, Cullis CA. 1993. RAPD analysis in flax: optimization of yield and reproducibility using klen Taq1 DNA polymerase, Chelex 100, and gel purification of genomic DNA. *Plant Mol Biol Rep* 11:128–141. <http://dx.doi.org/10.1007/BF02670471>.
9. McCarthy CB, Diambra LA, Rivera Pomar RV. 2011. Metagenomic analysis of taxa associated with *Lutzomyia longipalpis*, vector of visceral leishmaniasis, using an unbiased high-throughput approach. *PLoS Negl Trop Dis* 5:e1304. <http://dx.doi.org/10.1371/journal.pntd.0001304>.
10. McCarthy CB, Santini MS, Pimenta PF, Diambra LA. 2013. First comparative transcriptomic analysis of wild adult male and female *Lutzomyia longipalpis*, vector of visceral leishmaniasis. *PLoS One* 8:e58645. <http://dx.doi.org/10.1371/journal.pone.0058645>.
11. Niu B, Fu L, Sun S, Li W. 2010. Artificial and natural duplicates in pyrosequencing reads of metagenomic data. *BMC Bioinformatics* 11:187. <http://dx.doi.org/10.1186/1471-2105-11-187>.
12. Zhao Y, Tang H, Ye Y. 2012. RAPSearch2: a fast and memory-efficient protein similarity search tool for next-generation sequencing data. *Bioinformatics* 28:125–126. <http://dx.doi.org/10.1093/bioinformatics/btr595>.
13. Huson DH, Auch AF, Qi J, Schuster SC. 2007. MEGAN analysis of metagenomic data. *Genome Res* 17:377–386. <http://dx.doi.org/10.1101/gr.5969107>.
14. Huson DH, Mitra S, Ruscheweyh HJ, Weber N, Schuster SC. 2011. Integrative analysis of environmental sequences using MEGAN4. *Genome Res* 21:1552–1560. <http://dx.doi.org/10.1101/gr.120618.111>.
15. Fisher R. 1970. *Statistical methods for research workers*, p 96. 14th ed. Hafner Publishing, New York, NY.