



Artículo

# NEW HOST RECORDS AND GEOGRAPHIC DISTRIBUTION OF SPECIES OF *Trichuris* (NEMATODA: TRICHURIIDAE) IN RODENTS FROM ARGENTINA WITH AN UPDATED SUMMARY OF RECORDS FROM AMERICA

María del Rosario Robles and Graciela T. Navone

Centro de Estudios Parasitológicos y de Vectores CEPAVE (CCT-CONICET La Plata) (UNLP), Calle 2 # 584, (1900) La Plata, Buenos Aires, Argentina [correspondence: María del Rosario Robles <[rosario@cepave.edu.ar](mailto:rosario@cepave.edu.ar)>].

**ABSTRACT.** Species of *Trichuris* have a cosmopolitan distribution and parasitize a broad range of mammalian hosts. Although, the prevalence and intensity of this genus depends on many factors, the life cycles and characteristics of the environment have been the main aspect used to explain their geographical distribution. In this paper, we provide new host and geographical records for the species of *Trichuris* from Sigmodontinae rodents in Argentina. Moreover, we present comprehensive data about previous records of the genus from rodents in North and South America, and mainly in Argentina. A total of 563 specimens including in 25 species of rodents from 12 provinces and 43 localities from Northeast and Southern Argentina were sampled for *Trichuris*. Six species of *Trichuris* including 9 new hosts and 16 new geographical records were found. The prevalence of *Trichuris* in Misiones province is higher than in other geographical areas studied. This tropical-subtropical nematode group is recorded for the first time from the latitude 40° S. Nevertheless, more surveys are necessary to determine whether the scarcity of infections of *Trichuris* in regions of high latitude is due to the absence of sample effort or to environmental effects on the survival of infective stages.

**RESUMEN.** Nuevos registros hospedatorios y de distribución geográfica de especies de *Trichuris* (Nematoda: Trichuriidae) de roedores de Argentina con un resumen actualizado de los registros en América. Las especies de *Trichuris* presentan una distribución cosmopolita y parasitan un amplio rango de hospedadores mamíferos. Aunque la prevalencia e intensidad parasitaria de estos nematodos dependen de diferentes factores, su ciclo de vida y las características del ambiente han sido los principales aspectos que explican su distribución geográfica. En este trabajo se brindan nuevos hospedadores y registros geográficos para las especies de *Trichuris* de roedores Sigmodontinae para la Argentina. Además, se presentan datos completos y organizados sobre previos registros del género en roedores del continente americano, y principalmente de Argentina. Un total de 563 especímenes incluidos en 25 especies de roedores procedentes de 12 provincias y 43 localidades del noreste y del sur de la Argentina fueron examinados con el fin de hallar especímenes de *Trichuris*. Se presentan seis especies de *Trichuris* incluyendo 9 nuevos hospedadores y 16 nuevos registros geográficos. La prevalencia de *Trichuris* en la provincia de Misiones es más alta que en otras áreas geográficas estudiadas. Este grupo de nematodos tropical-subtropical es registrado por primera vez por sobre los 40° S. De todos modos, son necesarios nuevos estudios para explicar si la escasez de infecciones de *Trichuris* en regiones de altas latitudes es debido a la ausencia de esfuerzo de muestreo o al resultado del efecto del ambiente sobre la supervivencia del estadio infectivo.

**Key words:** America. Argentina. Nematoda. Rodentia. Sigmodontinae. *Trichuris*.

**Palabras clave:** América. Argentina. Nematoda. Rodentia. Sigmodontinae. *Trichuris*.

## INTRODUCTION

Species of *Trichuris* Roederer, 1761 (Nematoda: Trichuridae) have a cosmopolitan distribution and parasitize a broad range of mammalian hosts (Cafrune et al., 1999; Anderson, 2000). Among these, the most common species are *Trichuris trichiura* (Linnaeus, 1758) from humans, and *T. muris* (Schrank, 1788), *T. ovis* (Abildgaard, 1795), *T. suis* (Schrank 1788), and *T. vulpis* (Froelich, 1789) from synanthropic and domestic animals (e.g. Knight, 1971; Beer, 1976; Bundy and Cooper, 1989; Callejón et al., 2010). Although, the prevalence and intensity of *Trichuris* in a host population depend on many factors, host immunologic status, behavior, areas with inadequate sanitation (food and fecal deposition areas mixed); the characteristics of its life cycle and environmental limiting factors have been the main aspects used to explain the geographical distribution (Bundy et al., 1988; Bundy and Cooper, 1989; Grecnis et al., 1993; Anderson, 2000).

Whipworm eggs are deposited from host feces to the soil where infective larvae develop within the egg. Development of this larval stage has been shown to be influenced by temperature. Following ingestion of infective eggs by the host, all subsequent larval development to the adult stage occurs in the mucosa of the caecum and colon (Beer, 1973; Bundy and Cooper, 1988, 1989). Infections by trichuriasis are more prevalent in warm and moist tropical regions than in other parts of the world, and occur mainly in North and South America (Bundy and Cooper, 1988, 1989). However, most records are from medical and veterinary surveys (e.g. Bundy and Cooper, 1988; Traub et al., 2004; Gamboa et al., 2005), and comparatively few studies have been carried out on natural infections of *Trichuris* from wild hosts. In addition, many geographical areas have not been surveyed for *Trichuris*. Current distribution of the genus might reflect the absence of sample effort rather than the absence of infection (Bundy and Cooper, 1989).

Argentina is the second largest country in South America by land area, after Brazil, and exhibits a wide variety of soil types and climatic conditions (Bertonatti and Corcuera,

2000). The generally temperate climate ranges from subtropical in the north to subpolar in the far south and these environments include different eco-regions. Under natural conditions the rate of development of *Trichuris* species in this country may differ significantly from the optima, depending on the geographical distribution.

In this country, 8 species of *Trichuris* have been described and recorded from rodents (Morini et al., 1955; Boero and Boehringer, 1967; Suriano and Navone, 1994; Robles et al., 2006; Robles, 2011; Robles et al., 2012b). The only two records of this genus from wild hosts in other orders are *Trichuris tenuis* Chandler, 1930 from *Lama glama* and *Vicugna vicugna* (Artiodactyla; Cafrune et al., 1999) and *T. campanula* Linstow, 1889 from *Oncifelis geoffroyi* (Carnivora; Beldomenico et al., 2005). In general, the information available from many mammalian hosts is poor, hidden and scattered, making it difficult to evaluate the distribution and geographical gaps of the species of *Trichuris*.

Several parasitological studies on Sigmodontinae rodents (Cricetidae) have been carried out in Argentina in recent years (e.g. Robles and Navone, 2010; Notarnicola and Navone, 2011; Digiani et al., 2012; Robles et al., 2012a, 2012c; Digiani et al., 2013) and among these, three dealt with the taxonomy of *Trichuris* (Robles and Navone, 2006; Robles et al., 2006; Robles, 2011). However, these are only partial records, because several host species and *Trichuris* species are still under study.

In this paper, we provide new host and geographical records for *Trichuris* species from sigmodontine rodents of Argentina. Moreover, we present comprehensive data on previous records of the genus from rodents in North and South America, mainly in Argentina.

## MATERIALS AND METHODS

### Studied area

This includes the Northeast of Argentina, approximately between 26° and 35° S, politically covering the eastern Formosa, Chaco and Santa Fe provinces, the northern Buenos Aires province and Misiones, Corrientes and Entre Rios provinces. Also, the studied

area includes the south of Argentina (Patagonia), approximately between 38° and 54° S, embracing Chubut, Neuquén, Río Negro, Santa Cruz and Tierra del Fuego provinces (see **Appendix 1, supplementary material** for additional details on localities).

### Hosts

Sigmodontine rodents were trapped during different field works between 2007 and 2010 (see collectors and support in acknowledgements). Many specimens are still being studied, but partial results were analyzed. Species of hosts with a very low representation (<5) have not been considered in this paper (except *Chelemys macronyx* with n=3). A total of 563 specimens were examined for *Trichuris*. From the northeast area, 493 specimens belonging to 17 species from 29 localities and 7 provinces were examined as following: *Akodon azarae* (n=118) [*A. azarae bibiana* (n=12), *A. azarae hunteri* (n=106)], *Akodon philipmyersi* (n=14), *Brucepatersonius* sp. (n=6), *Calomys* sp. (n=13), *Euryoryzomys russatus* (n=8), *Holochilus brasiliensis* (n=5), *H. chacarius* (n=13), *Necromys lasiurus* (n=109), *Necromys obscurus* (n=11), *Nectomys squamipes* (n=5), *Oligoryzomys flavescens* (n=37), *O. fornesi* (n=10), *O. nigripes* (n=71), *Oxymycterus rufus* (n=45), *Scapteromys aquaticus* (n=5), *Sooretamys angouya* (n=8), *Thaptomys nigrita* (n=15). From Patagonia we examined 70 specimens belonging to 8 species from 14 localities and 5 provinces as following: *Abrothrix hirta* (n=12), *A. olivacea* (n=19), *Akodon dolores* (n=5), *Chelemys macronyx* (n=3), *Eligmodontia morgani* (n=6), *Euneomys* sp. (n=6), *Oligoryzomys longicaudatus* (n=10), *Phyllotis xanthopygus* (n=9) (see **Appendix 1, supplementary material**).

### Parasites

Nematodes were collected from the large intestine and caecum and preserved in 70% ethanol. For identification, the worms were prepared and identified following Robles et al. (2006), Robles and Navone (2006) and Robles (2011). Voucher specimens were deposited in the Colección de Helmintología from Museo de La Plata, Argentina (CHMLP).

### Data analysis

Quantitative parameters of prevalence (P=specimens parasitized/specimens examined \*100), intensity (I=number of parasites in a single infected host), mean intensity (MI=number of parasites/specimens parasitized) and mean abundance (MA=number of parasites/specimens examined) were calculated according to Bush et al. (1997) for each host species

and locality (**Table 1**). Prevalence, MI, I and MA for host species and studied area were given in the text. Prevalence differences were compared by the  $\chi^2$  test, and the data were analyzed by use of the Chi squared test employing the correction of Yates or by use of Fisher's exact test (positive cases <5), considering significant at  $p < 0.05$ . Statistical analysis was performed with EpiInfo™ 7.

## RESULTS

Ten species of sigmodontine rodents from Argentina were parasitized with species of *Trichuris*. Three species of *Trichuris* were identified to the specific level, and at least 3 others were studied in detail, although these could not be identified to the species level (**Table 1**).

Published and new records for *Trichuris* in Argentina are shown in **Fig. 1** (also see **Tables 1** and **2** for species references). Records of *Trichuris* from the literature were summarized for Argentina in **Table 2** and those for North and South America in **Table 3**.

*Akodon azarae bibiana*, *A. azarae hunteri* and *N. lasiurus* were parasitized with *Trichuris laevitesticis* Suriano and Navone, 1994. *A. azarae bibiana* and *N. lasiurus* are new host records, and 11 localities are new geographical records (**Table 1**). The total P, MI and MA of *T. laevitesticis* in all host species were 11.9%, 4.03 and 0.48, respectively; and in each host species: *A. azarae* 16.9%, 3.85 and 0.65 and *N. lasiurus* 6.42%, 4.57 and 0.29, respectively. *T. nigrita* were parasitized with a species of *Trichuris*. This host and Refugio Moconá represent new records for this nematode (**Table 1**). General features of these specimens suggest that they belong to *Trichuris navonae* Robles, 2011. However, some diagnostic measurements (in millimeters) such as spicule length (1.9-2.3), distal cloacal tube length (1.7-2.1), proximal cloacal tube lengths (1.2-1.7), present ranges higher than those previously recorded for *T. navonae* from *A. montensis* (1.3-2.1, 1-1.9, 0.75-1.5; respectively), verified through their ratios with the posterior portion of body and total body lengths. For this reason, these specimens are listed provisionally as *Trichuris* cf. *T. navonae* until more detailed morphometric analyses and molecular comparative studies determine whether they are cryptic species. The P, MI and

**Table 1**

New hosts and geographical records for species of *Trichuris* in Argentina with data of Prevalence (P), Intensity (I), Mean Intensity (MI) and Mean Abundance (MA) by locality and host species; numbers and symbols refer to Fig. 1.

<i>Trichuris</i> species	Host species	Locality*	Prevalence	Intensity or Mean intensity	Mean abundance
1 <i>T. laevitesitii</i>	<i>A. azarae bibiana</i> **	7 km S Puerto Las Palmas, Chaco**	25% (1/4)	2+	0.5 (2/4)
	<i>A. azarae hunteri</i> <sup>a</sup>	INTA-IPAF NEA, Formosa**	33.3% (1/3)	4+	1.33 (4/3)
		Arroyo de las Brusquitas, Buenos Aires**	37% (10/27)	1.9 (19/10)	0.7 (19/27)
		Punta Indio, Buenos Aires**	9.1% (1/11)	2+	0.2 (2/11)
		Zárate, Buenos Aires**	18.2% (2/11)	1 (2/2)	0.2 (2/11)
		Arroyo Caraballo, Entre Ríos**	25% (1/4)	35+	8.8 (35/4)
		Colonia Villa Elisa, Entre Ríos**	12.5% (3/24)	2 (6/3)	0.3 (6/24)
		Oliveros, Santa Fe**	(1/1)	7+	7 (7/1)
	<i>N. lasiurus</i> **	Finca La Adelita, Laguna Paiva, Corrientes**	14.3% (2/14)	1.5 (3/2)	0.2 (3/14)
		Pergamino, Buenos Aires**	12.5% (2/16)	3 (6/2)	0.4 (6/16)
		Uranga, Santa Fe**	16.6% (3/18)	7.6 (23/3)	1.3 (23/18)
2 <i>T. cf. navonae</i>	<i>T. nigrita</i> **	Refugio Moconá, Misiones**	75% (3/4)	2 (6/3)	1.5 (6/4)
		Desembocadura Arroyo Paranay-Guazú, Misiones <sup>c</sup>	33% (1/3)	6+	2 (6/3)
		Balneario Municipal de Aristóbulo del Valle, Misiones <sup>c</sup>	85.7% (6/7)	1.16 (7/6)	1 (7/7)
3 <i>T. cf. T. paradinasi</i>	<i>P. xanthopygus xanthopygus</i> **	Cerro Corona, Meseta de Sumuncurá, Río Negro**	(2/2)	10 (20/2)	10 (20/2)
4 <i>Trichuris</i> sp. 1	<i>H. chacaricus</i> **	INTA-IPAF NEA, Formosa <sup>#</sup>	23.1% (3/13)	18.6 (56/3)	4.3 (56/13)
5 <i>Trichuris</i> sp. 2	<i>S. angouya</i> **	Refugio Moconá, Misiones <sup>#</sup>	50% (2/4)	7 (14/2)	3.5 (14/4)
		Reserva de Usos Múltiples Guaraní, Misiones**	(2/2)	1 (2/2)	1 (2/2)
6 <i>Trichuris</i> sp. 3	<i>C. macronyx</i> **	Cañadón de la Madera, Sierra de Tepuel, Chubut**	33.3% (1/3)	10+	3.3 (10/3)
7 <i>Trichuris</i> sp. 4	<i>N. obscurus</i> **	Arroyo de las Brusquitas, Buenos Aires <sup>#</sup>	36.4% (4/11)	1.5 (6/4)	0.5 (6/11)
‡ <i>Trichuris</i> sp.	<i>E. russatus</i> **	Arroyo Paraiso, Misiones <sup>c</sup>	(1/1)	1+	1 (1/1)
‡ <i>Trichuris</i> sp.	<i>S. aquaticus</i> <sup>b</sup>	Estancia San Nicolás, Corrientes**	20% (1/5)	1+	0.2 (1/5)

\* See complete data in supplementary material; \*\* denotes a new host and/or a new locality records.

<sup>a</sup> Robles and Navone, 2006; <sup>b</sup> Suriano and Navone, 1994; <sup>c</sup> Robles, 2011. <sup>#</sup> shared record; + intensity.




Fig. 1. Previous and new records for the species of *Trichuris* in Argentina. Localities are detailed in Tables 1 and 2. New records are marked with a circle.

data). The P, MI and MA of *T. cf. T. pardinasi* in *P. x. xanthopygus* were 28.6%, 10 and 2.86, respectively.

*Sooretamys angouya*, *N. obscurus*, *H. chacarius*, and *C. macronyx* were parasitized, each one with 1 species of *Trichuris*. These findings constitute 4 new host records and 4 new locality records (Table 1). *Trichuris* specimens from the first 3 host species listed above are mainly characterized by the absence of a spicular tube, a spinose spicular sheath (spines very small, densely arranged), and a slightly protusive or nonprotusive vulva. The *Trichuris* specimens from *C. macronyx* are characterized by the

presence of a spicular tube, a spinose spicular sheath (spines very small, densely arranged), and a protusive vulva; along with morphometric characters. Morphologically these specimens correspond to 4 different species of *Trichuris*, suggesting the presence of 2 new species, one from *S. angouya* and one from *H. chacarius*; and 2 preliminary unidentified species, one from *N. obscurus* and one *C. macronyx*. Until new samples are obtained and molecular comparative studies are concluded these species are listed here as *Trichuris* sp. The P, MI and MA of the species of *Trichuris* were in *S. angouya*, 50%, 4 and 2, respectively; in *N. obscurus*, 36.4%, 1.5 and 0.54, respectively; in *H. chacarius*, 16.6%, 18.6 and 3.11, respectively; and in *C. macronyx* 33%, 10 and 3.33, respectively.

MA of *T. cf. T. navonae* in the population of *T. nigrita* were 66.6%, 1.9 and 1.26, respectively. *Phyllotis xanthopygus xanthopygus* were parasitized with another species of *Trichuris*. This host subspecies represents a new host record for this nematode. In addition, these results add a new locality and province, Cerro Corona and Río Negro, respectively (Table 1). General features of these specimens suggest that they belong to *Trichuris pardinasi* Robles, Navone and Notarnicola, 2006. However, some morphometrical discrepancies (in millimeters) such as spicule lengths were observed (2.5-3.3 in *P. x. xanthopygus* vs. 3.6-5.2 in *P. bonariensis* and *P. x. vaccarum*). For this reason, these specimens are listed provisionally as *Trichuris cf. T. pardinasi*. It would be useful to obtain new samples and data from other sources (i.e. genetic

presence of a spicular tube, a spinose spicular sheath (spines very small, densely arranged), and a protusive vulva; along with morphometric characters. Morphologically these specimens correspond to 4 different species of *Trichuris*, suggesting the presence of 2 new species, one from *S. angouya* and one from *H. chacarius*; and 2 preliminary unidentified species, one from *N. obscurus* and one *C. macronyx*. Until new samples are obtained and molecular comparative studies are concluded these species are listed here as *Trichuris* sp. The P, MI and MA of the species of *Trichuris* were in *S. angouya*, 50%, 4 and 2, respectively; in *N. obscurus*, 36.4%, 1.5 and 0.54, respectively; in *H. chacarius*, 16.6%, 18.6 and 3.11, respectively; and in *C. macronyx* 33%, 10 and 3.33, respectively.

Table 2  
Host and geographical data for the species of *Trichuris* previously recorded from rodents in Argentina; numbers and symbols refer to Fig. 1.

<i>Trichuris</i> species	Host species	Locality	Coord S	Coord W	Province	References		
1 <i>T. laevitesticis</i>	<i>Akodon azarae hunteri</i>	Punta Lara	34°47'30"	58°0'5"	Buenos Aires	Suriano and Navone, 1994		
		La Balandra	34°55'51"	57°43'0.6"	Buenos Aires	Robles and Navone, 2006		
		Reserva Otamendi	34°09'	58°57'	Buenos Aires	Robles and Navone, 2006		
		Cerro de la Gloria	36°06'	57°46'	Buenos Aires	Robles and Navone, 2006		
		Punta Lara	34°47'30"	58°0'5"	Buenos Aires	Suriano and Navone, 1994		
		La Balandra	34°55'51"	57°43'0.6"	Buenos Aires	Robles and Navone, 2006		
		Los Talas	34°52'05"	57°49'20"	Buenos Aires	Robles and Navone, 2006		
		Palo Blanco	34°52'24"	57°48'47"	Buenos Aires	Robles and Navone, 2006		
		2 <i>T. navonae</i>	<i>Akodon montensis</i>	Balneario Municipal de Aristóbulo del Valle	27°05'17"	54°57'9"	Misiones	Robles, 2011
				Club Pesca Paraná-Guazú	26°40'32"	54°48'51"	Misiones	Robles, 2011
Desembocadura arroyo Paraná-Guazú	26°40'39"			54°50'8"	Misiones	Robles, 2011		
Arroyo Salamanca	26°36'53"			54°46'51"	Misiones	Robles, 2011		
Salto El Paraíso, arroyo Paraíso	27°13'49"			54°02'24"	Misiones	Robles, 2011		
Arroyo Oveja Negra - Ruta 21	27°8'20"			53°55'31"	Misiones	Robles, 2011		
Parque Provincial Cruce Caballero	26°30'50"			53°59'54"	Misiones	Robles, 2011		
Parque Provincial Moconá**	27°9'23"			53°54'10"	Misiones	Robles, 2011		
Puerto Peninsula	25°40'			54°38'	Misiones	Robles, 2011		
3 <i>T. pardinasi</i>	<i>Phyllotis bonariensis</i>			Cerro Destierro, Sierra de la Ventana	38°1'25"	62°54'39"	Buenos Aires	Robles et al., 2006
		Abra de la Ventana, Sierra de la Ventana	38°4'3"	62°1'17"	Buenos Aires	Robles et al., 2006		
3 <i>T. pardinasi</i>	<i>Phyllotis xanthopygus vaccarum</i>	Pampa de Achala	31°37'19"	64°54'39"	Córdoba	Robles et al., 2006		



8	<i>T. bursacaudata</i>	<i>Ctenomys talarum</i>	Punta de Indio	35°16'27"	57°15'38"	Buenos Aires	Suriano and Navone, 1994
9	<i>T. dolichotis</i>	<i>Dolichotis patagonum</i>	Jardín zoológico, Ciudad de Buenos Aires*	34°34'44"	58°24'55"	Buenos Aires	Morini et al., 1955
λ	<i>T. myocastoris</i>	<i>Myocastor coypus</i>	Región del Delta*	34°21'32"	58°40'53"	Buenos Aires	Boero and Boehringer, 1967
φ	<i>T. pampaeana</i>	<i>Ctenomys azarae</i>	Región Chaqueña*	27°20'27"	58°47'9"	Chaco	Boero and Boehringer, 1967
		<i>Ctenomys australis</i>	Santa Rosa	36°36'33"	64°19'2"	La Pampa	Suriano and Navone, 1994
		<i>Ctenomys talarum</i>	Necochea	38°33'	58°45'	Buenos Aires	Rossin et al., 2010
			Necochea	38°33'	58°45'	Buenos Aires	Rossin and Malizia, 2005
X	<i>T. cf. thrichomysi</i>	<i>Euryzgomatomys spinosus</i>	Mar de Cobo, Pdo. Mar Chiquita	37°58'	57°34'	Buenos Aires	Rossin and Malizia, 2005
			Cuartel Río Victoria, Ruta 14 km 273, INTA	26°46'	54°18'	Misiones	Robles et al., 2012

\* Area not detailed; \*\* This locality is very near to Refugio Moconá (see **Table 1**)

*Euryzomys russatus* and *S. aquaticus* from Ea. San Nicolás were each parasitized with a *Trichuris* sp., constituting a new host and a new geographical record, respectively (**Table 1**). In both cases, species identification was not possible due to the absence of males. The P, MI and MA of the species of *Trichuris* in each host species were 12.5%, 1, 0.13 and 20%, 1, 0.2, respectively.

Specimens of *Trichuris* were not distributed homogeneously among the host species and study areas. The prevalence between species pairs: *A. azarae* with *N. lasiurus*, *T. nigrita* and *S. angouya*; *N. lasiurus* with *N. obscurus*, *T. nigrita* and *S. angouya*; *T. nigrita* with *E. russatus* and *H. chacarius* were significantly different ( $p < 0.05$ ). On the other hand, the prevalence between other host species pairs studied did not differ significantly. Moreover, the prevalence between the two studied areas, Northeast (P 9.53%) and South (P 4.41%) of Argentina, were not significantly different ( $p=0.25$ ).

In summary, we recorded 9 new hosts (7 species and 2 subspecies) and 16 new geographical records for *Trichuris*, enlarging significantly their distributional range (**Table 1**).

A total of 8 species of *Trichuris* from 11 rodent species have been recorded in Argentina since 1955 (Morini et al., 1955; Boero and Boehringer, 1967; Suriano and Navone, 1994; Rossin and Malizia, 2005; Robles and Navone, 2006; Rossin et al., 2010; Robles et al., 2006; Robles, 2011; Robles et al., 2012b). These species are distributed approximately from 25° 12' S to 43° 51' S and from 70° 43' W to 53° 54' W (**Tables 1** and **2**). A total of 24 species of *Trichuris* from 33 rodent species have been recorded in the Americas since 1955. These species are distributed from Manitoba (N) (Canada) to Texas (S) and from California (W) to Maryland (E) (USA); and in South America from Trinidad (N) (Republic of Trinidad and Tobago) to La Pampa (S) (Argentina) and from Minas Gerais (W) (Brazil) to La Huya (E) (Chile) (**Table 3**) (Lent and Freitas, 1936; Chandler, 1945; Tiner, 1950; Cameron and Reesal, 1951; Morini, et al. 1955; Read, 1956; Boero and Boehringer, 1967; Kenneth and Lepp, 1972; Barus et al., 1975; Babero et al.,

**Table 3**

Host and geographical data previously recorded for species of *Trichuris* from rodents in America excluding Argentinean records.

<i>Trichuris</i> species	Host species	Host family	Locality	Country
<i>T. leporis</i> (Froelich, 1789)	<i>Spermophilus richardsonii</i> <sup>§</sup>	Sciuridae	Manitoba	Canada
<i>T. gracilis</i> (Rudolphi, 1819)	<i>Dasyprocta leporina</i> <sup>*</sup> , <i>D. fuliginosa</i>	Dasyproctidae	State of Amazonas	Trinidad Brazil
<i>T. opaca</i> Barker et Noyes, 1915	<i>Ondatra zibethicus</i> , <i>Microtus pennsylvanicus</i>	Cricetidae	Nebraska, Wisconsin, Ohio, Wyoming, Maryland	USA
<i>T. fossor</i> Hall, 1916	<i>Thomomys talpoides</i> <sup>†</sup> , <i>T. bottae</i>	Geomyidae	California, Wyoming	USA
<i>T. myocastoris</i> Enigk, 1933	<i>Myocastor coypus</i>	Myocastoridae	San Pablo	Brazil
<i>T. citelli</i> Chandler, 1945	<i>Spermophilus beecheyi</i> <sup>‡</sup>	Sciuridae	California	USA
<i>T. perognathi</i> Chandler, 1945	<i>Perognathus californicus</i> <i>californicus</i> , <i>P. pennicillata</i>	Heteromyidae	California	USA
<i>T. neotomae</i> Chandler, 1945	<i>Neotoma fuscipes</i>	Cricetidae	California	USA
<i>T. peromysci</i> Chandler, 1946	<i>Peromyscus californicus</i>	Cricetidae	California	USA
<i>T. madisonensis</i> Tiner, 1950	<i>Tamias striatus</i>	Sciuridae	Wisconsin	USA
<i>T. dipodomys</i> (Read, 1956)	<i>Dipodomys ordii</i> , <i>D. phillipsi</i>	Heteromyidae	New Mexico, Utah	USA
<i>T. bradleyi</i> Babero, Cattan and Cabello, 1975	<i>Octodon degus</i>	Octodontidae	Santiago	Chile
<i>T. chilensis</i> Babero, Cattan and Cabello, 1976	<i>Abrothrix longipilis</i> <sup>a</sup>	Cricetidae	Santiago	Chile
<i>T. fulvi</i> Babero et Murua, 1987	<i>Ctenomys fulvus</i> <i>phillipiensis</i>	Ctenomyidae	San Pedro de Atacama	Chile
<i>T. elatoris</i> Pfaffenberger and Best, 1989	<i>Dipodomys elator</i> , <i>D. merriami</i> , <i>D. ordii</i>	Heteromyidae	Texas	Mexico
<i>T. robusti</i> Babero and Murua, 1990	<i>Ctenomys robustus</i>	Ctenomyidae	La Hauyca	Chile
<i>T. travassosi</i> Correa Gomes, Lanfredi, Pinto and Souza, 1992	<i>Oligoryzomys nigripes</i> <sup>#</sup>	Cricetidae	Rio Grande do Sul	Brazil



(Table 3 cont.)

<i>Trichuris</i> species	Host species	Host family	Locality	Country
<i>T. thrichomysi</i> Lopes Torres et al., 2011	<i>Thrichomys apereoides</i> , <i>Euryzgomatomys spinosus</i>	Echimyidae	Minas Gerais	Brazil

<sup>§</sup> Cited as *Citellus richardsoni* by Tiner, 1950; \* Cited as *Cavia agouti* by Rudolphi, 1819; † Cited as *Thomomys fossor* by Hall, 1916; ‡ Cited as *Citellus beecheyi* by Chandler, 1945; α Cited as *Akodon longipilis* by Babero, Cattán and Cabello, 1976; # Cited as *Oryzomys nigripes* by Correa Gomes et al., 1992. Families of Argentinean records, Caviidae: *T. dolichotis*; Cricetidae: *T. laevitestis*, *T. pardinasi* and *T. navonae*; Ctenomyidae: *T. bursacaudata* and *T. pampeana*.

1975, 1976; Babero and Murua, 1987; 1990; Pfaffnberger and Bests, 1989; Correa Gomes et al., 1992; Suriano and Navone, 1994; Gonçalves et al., 2002; Rossin and Malizia, 2005; Robles and Navone, 2006; Robles et al., 2006; Rossin et al., 2010; Robles, 2011; Robles et al., 2012b).

## DISCUSSION

Each species of *Trichuris* geographically located in Argentina can be distinguished by diagnostic features. In several cases (e.g., *Trichuris* specimens obtained from *C. macronyx*, *E. russatus*, *N. obscurus*, and *S. aquaticus*), observed differences in morphological and metric traits prevent us a confident specific assignation. Two possible new species were found from *S. angouya* and *H. chacarius* each one. However, new and more detailed studies and material are necessary to clarify the alpha taxonomy of *Trichuris* in Argentina.

To date, a total of 24 species of *Trichuris* have been described from 10 families of rodents in America, Caviidae (1 species), Cricetidae (8), Ctenomyidae (4), Dasyproctidae (1), Echimyidae (1), Geomyidae (1), Heteromyidae (3), Myocastoridae (1), Octodontidae (1), and Sciuridae (3). Of these, 8 species have been reported from Argentina, including *Trichuris dolichotis* Morini, Boero and Rodriguez, 1955, *T. myocastoris* Enigk, 1933, *T. laevitestis* *T. bursacaudata* Suriano and Navone, 1994, *T. pampeana* Suriano and Navone, 1994, *T. pardinasi*, *T. navonae*, *T. cf. T. thrichomysi* Lopes Torres et al., 2012 (Morini et al., 1955; Boero and Boehringer, 1967; Suriano and Navone, 1994; Robles et al., 2006; Robles, 2011; Robles et al., 2012b) and 4 additional species,

not identified to specific level, mentioned in this paper (Tables 1, 2 and 3).

Furthermore, a total of 9 species of *Trichuris* have been recorded from 14 sigmodontines (*A. azarae*, *A. montensis*, *A. hirta*, *C. macronyx*, *E. russatus*, *H. chacarius*, *O. nigripes*, *P. bonariensis*, *P. xanthopygus*, *N. lasiurus*, *N. obscurus*, *T. nigrita*, *S. aquaticus*, and *S. angouya*; Tables 1, 2 and 3). Sigmodontine rodents are distributed predominantly in South America, with a few species having a Central and North American distribution, and this subfamily includes approximately 400 species (Patton et al., in press). Despite the large number of potential host species, only about 3.5% of sigmodontine rodent species have been recorded as hosts for species of *Trichuris*. Among these hosts, 86% of infections were recorded in Argentina (Morini et al., 1955; Boero and Boehringer, 1967; Suriano and Navone, 1994; Robles and Navone, 2006; Robles et al., 2006; Robles, 2011; Robles et al., 2012b). This number may represent only a small fraction of the species of *Trichuris* that occurs in sigmodontine rodents, and additional surveys of this group should yield both new species and records.

The distribution of *Trichuris* species among the species of sigmodontine rodents in the surveyed localities showed a range of prevalence between 9.1-85.7% (samples of 2 or < specimens were not considered). Although the prevalence values given for some localities may be questionable because of the low number of specimens examined, these data give a new overview of the regional distribution of *Trichuris*. *Trichuris* cf. *T. navonae*

and *Trichuris* sp. from 3 localities in Misiones showed the highest prevalence (85.7%, 75% and 50%) (**Table 1**). However, *T. laevitesticis* from *A. a. hunteri* of Arroyo Caraballo (I = 35, MA = 8.8) and *Trichuris* sp. from *H. chacarius* of INTA-IPAF NEA (MI = 18.6, MA = 4.3) showed the highest intensities.

The significant differences observed in the prevalences of the species of *Trichuris* among host species probably relate to a combination of factors, including host immunologic status, behavior of the host species and characteristics of environment where the eggs were deposited. However, the prevalences between the two studied areas, northeast and south of Argentina, were not significantly different ( $p > 0.05$ ). Since this is a limited study, these results are not conclusive and surely new data on the soil and microenvironment characters, home range and behavior of host species, and experimental studies will lead to clearer hypotheses about the limiting factors of *Trichuris* present in wild mammals and extreme geographical areas.

Because of this, a previous hypothesis about the global geographical distribution (tropical and subtropical areas) of genus *Trichuris* should be revisited. To date, different studies have demonstrated that the embryonation period varies between 9-37 days and 25-35 °C for *T. trichiura*, *T. muris* and *T. vulpis* (Beer, 1971, 1976). A field survey in southern England found that egg development in *T. suis* was greatly retarded at ground temperatures of 4-20° C, with an embryonation period of 434-630 days (Burden and Hammet, 1979). Moreover, species of *Trichuris* have been infrequently recorded in high latitudes (Tiner, 1950; Burden and Hammet, 1979; Sardella and Fugassa, 2009). These examples may explain why trichuriasis in humans is presently almost unknown in cold regions; and in the same way, in rodents.

Patagonia presents unique environment characteristics, comprising a series of plateaus and high plains, a vegetation in open patches, and a temperate climate, which ranges from subtropical in the north to subpolar in the far south and at higher elevations (e.g. Soriano, 1956; León et al., 1998). To date, in the subpolar areas there have been no previous records of *Trichuris* in Patagonia, except in

archaeological samples (Sardella and Fugassa, 2009). However, in this survey, a population of this tropical-subtropical nematode is recorded for the first time from 40° S (*T. cf. T. pardinasi* and *Trichuris* sp., see **Table 1**).

In the context of the hypothesis above referred to the absence of *Trichuris* in the north, west and south of Argentina is consistent with a lower degree of sampling effort with respect to other areas in the country (**Fig. 1**). Parasitological studies have been carried out in the northeast area of Argentina for a long time, while these studies in Patagonian area have begun recently. It is likely that future studies will reveal the presence of more species and a more accurate host and geographical distribution, mainly in the southern portion of the country.

Consequently, more surveys on different host groups are necessary to determine whether the scarcity of infections of the genus *Trichuris* in regions of high latitude is due to the absence of sample effort or to environmental effects on the survival of infective stages.

## ACKNOWLEDGEMENTS

We offer thanks to Carlos Galliari, Marcela Lareschi, Juliana Notarnicola, Joaquín Pardiñas, Ulyses Pardiñas, Juliana Sanchez, Pablo Teta, Daniel Udrizar Sauthier, and collaborators for their cooperation in host collections. To Ulyses Pardiñas and Carlos Galliari for the identification of the hosts. To Mike Kinsella for the valuable critical reading of the manuscript and for the revision into English. To Luciana Riccialdelli and Santiago Favoretti for their help with the map. To Lorena Zonta for her assistance with statistic test. To Juliana Notarnicola and an anonymous reviewer for their contribution on the manuscript. This study was funded by Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) (PICT 0924 to GN; PICT 0547- PIP 6179 to U. Pardiñas; PICT 1634 to MRR) and Universidad Nacional de La Plata (N627 to GN). GN and MRR are members of CONICET.

## LITERATURE CITED

- ANDERSON RC. 2000. Nematode parasites of vertebrates. Their Development and Transmission. 2<sup>nd</sup> ed. CAB International, ed. Wallingford, Oxon, U. K.
- BABERO BB and RB MURÚA. 1987. The helminth fauna of Chile. X. A new species of whipworm from a Chilean Rodent. Transactions of the American Microscopical Society 106:190-193.
- BABERO BB and RB MURÚA. 1990. A new species of whipworm from a South American hystricomorph

- rodent. Memorias do Instituto Oswaldo Cruz 85: 211-213.
- BABERO BB, PE CATTAN, and C CABELLO. 1975. *Trichuris bradleyi* sp. n. a whipworm from *Octodon degus* in Chile. Journal of Parasitology 61:198-206.
- BABERO BB, PE CATTAN, and C CABELLO. 1976. A new species of whipworm from the rodent *Akodon longipilis* in Chile. Transactions of the American Microscopical Society 95:232-235.
- BARUS V, G MADJUMDAR, and TK MIKAILOV. 1975. Morphology and taxonomy of *Trichocephalus myocastoris* (Enigk, 1933). Folia Parasitologica 22:207-213.
- BEER RJS. 1971. Whipworms of domestic animals. Veterinary Bulletin 41:343-349.
- BEER RJS. 1973. Studies on the biology of the life-cycle of *Trichuris suis* Schrank, 1788. Parasitology 67:253-262.
- BEER RJS. 1976. The relationship between *Trichuris trichura* (Linnaeus, 1758) of the man and *Trichuris suis* (Schrank, 1788) of the pig. Research in Veterinary Science 20:47-54.
- BELDOMENICO PM, JM KINSELLA, MM UHART, GL GUTIERREZ, J PEREIRA, H DEL VALLE FERREYRA, and CA MARULL. 2005. Helminths of Geoffroy's cat, *Oncifelis geoffroyi* (Carnivora, Felidae) from the Monte desert, central Argentina. Acta Parasitologica 50:263-266.
- BERTONATTI C and J CORCUERA. 2000. Situación ambiental Argentina 2000. Fundación Vida Silvestre Argentina, Buenos Aires.
- BOERO JJ and IK BOEHRINGER. 1967. Los parásitos del carpincho (*Hydrochoerus hydrochoeris*) y del quiyá (*Myocastor coypus*). Revista de la Facultad de Ciencias Veterinarias, La Plata 21:161-172.
- BUNDY DAP and ES COOPER. 1988. Trichuriasis 2<sup>nd</sup> ed. Pp. 120-156, en: Tropical and geographical medicine (KS Warren and AA. Mahmoud, eds). McGraw-Hill, New York.
- BUNDY DAP and ES COOPER. 1989. *Trichuris* and Trichuriasis in humans. Advances in Parasitology 28:107-173.
- BUNDY DAP, SP KAN, and R ROSE. 1988. Age-related prevalence, intensity and frequency distribution of gastrointestinal helminth infection in urban slum children from Kuala Lumpur, Malaysia. Transactions of the Royal Society of Tropical Medicine and Hygiene 82:289-294.
- BURDEN DJ and NC HAMMET. 1979. The development and survival of *Trichuris suis* ova on pasture plots in the south of England. Research in Veterinary Science 26:66-70.
- BUSH AO, KD LAFFERTY, JM LOTZ and AW SHOSTAK. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. Journal of Parasitology 83:575-583.
- CAFRUNE MM, DH AGUIRRE and LG RICKARD. 1999. Recovery of *Trichuris tenuis* Chandler, 1930, from camelids (*Lama glama* and *Vicugna vicugna*) in Argentina. Journal of Parasitology 85:961-962.
- CAMERON TWM and MR REESAL. 1951. Studies on the endoparasitic fauna of Trinidad mammals. Canadian Journal of Zoology 29:276-289.
- CALLEJÓN R, M DE ROJAS, C NIEBERDING, P FORONDA, C FELIÚ, D GUEVARA, and C CUTILLAS. 2010. Molecular evolution of *Trichuris muris* isolated from different Muridae hosts in Europe. Parasitology Research 107:631-641.
- CHANDLER AC. 1945. *Trichuris* species from California. Journal of Parasitology 31:284-287.
- CORREA GOMES D, RM LANFREDI, RM PINTO, and W DE SOUZA. 1992. Description of *Trichuris travassosi* n. sp. (Nematoda: Trichurinae) from a Brazilian rodent, by light and scanning electron microscopy. Memorias do Instituto Oswaldo Cruz 87:1-10.
- DIGIANI MC, J NOTARNICOLA, and GT NAVONE. 2012. The genus *Guerrerostrongylus* (Nematoda, Heligmonellidae) in cricetid rodents from the Atlantic Rain Forest of Misiones, Argentina: Emended description of *Guerrerostrongylus zetta* (Travassos, 1937) and description of a new species. 2012. Journal of Parasitology 98:985-991.
- DIGIANI MC, J NOTARNICOLA, and MS PAULOS. 2013. *Mazzanema* n. gen. and *Mazzanema fortuita* n. comb. for *Longistriata fortuita* Freitas, Lent and Almeida, 1937 (Nematoda, Heligmonellidae), a parasite of the marsh rat *Holochilus chacarius* (Rodentia, Cricetidae) from Northern Argentina. Journal of Parasitology 99:816-820.
- GAMBOA MI, G NAVONE, L KOZUBSKY, E COSTAS, L SUSEVICH, M CARDOZO, M GARRAZA, and P MAGISTRELLO. 2005. Geohelminthosis en tres poblaciones suburbanas con diferente condición sociocultural. Parasitología Latinoamericana 60:244.
- GONÇALVES AQ, JJ VICENTE, and RM PINTO. 2002. Nematodes of Amazonian vertebrates deposited in the Helminthological Collection of the Oswaldo Cruz Institute with new records. Revista Brasileira de Zoologia, Curitiba 19:453-465.
- GRENCIS RK, KJ ELSE, AJ BANCROFT, and DAP BUNDY. 1993. *Trichuris* Update '93. Parasitology Today 9:309-310.
- KENNETH ST and DL LEEP. 1972. Redescription of *Trichuris fossor* Hall, 1916 (Nematoda: Trichuridae) from the Northern Pocket Gopher, *Thomomys talpoides*. Proceedings of the Helminthological Society of Washington 39:203-205.
- KNIGHT RA. 1971. Redescriptions of *Trichuris discolor* (von Linstow, 1906) and *T. skrjabini* (Baskakov, 1924) from domestic ruminants in the United States and comparison with *T. ovis* (Abildgaard, 1795). Journal of Parasitology 57:302-310.
- LENT H and JFT FREITAS. 1936. Sobre o *Trichuris* da nutria. Anais da Academia Brasileira de Ciências 8:319-322.
- LEÓN RJC, D BRAN, M COLLANTES, JM PARUELO, and A SORIANO. 1998. Grandes unidades de la Patagonia extra andina. Ecología Austral 8:125-144.
- MORINI EG, BOERO J, and A RODRIGUEZ. 1955. Parásitos intestinales en el «Marra» (*Dolichotis patagonum patagonum*). Publicación Misión de Estudios de Patología Regional Argentina 26:83-89.
- NOTARNICOLA J and GT NAVONE. 2011. *Litomosoides pardinasi* n. sp. (Nematoda, Onchocercidae) from two species of cricetid rodents in Northern

- Patagonia, Argentina. Parasitology Research 108:187-194.
- PATTON J, UFJ PARDIÑAS, and G D'ELÍA (eds.). 2014. Mammals of South America. Volume 2, Rodents. The University of Chicago Press, Chicago, IL.
- PFaffenberger GS and T BEST. 1989. *Trichuris elatoris* sp. n. (Nematoda: Trichuridae) from the Texas kangaroo rat (*Dipodomys elator*). Proceedings of the Helminthological Society of Washington 56:76-81.
- READ CP. 1956. *Trichuris dipodomis*, n. sp., from Ord's Kangaroo Rat. Proceedings of the Helminthological Society of Washington 23:119.
- ROBLES MR. 2011. New species of *Trichuris* (Nematoda: Trichuridae) from *Akodon montensis* Thomas, 1913 of the Paranaense forest in Argentina. Journal of Parasitology 97:319-327.
- ROBLES MR and GT NAVONE. 2006. Redescription of *Trichuris laevitestis* (Nematoda: Trichuridae) from *Akodon azarae* and *Scapteromys aquaticus* (Sigmodontinae: Muridae) in Buenos Aires province, Argentina. Journal of Parasitology 92:1053-1057.
- ROBLES MR and GT NAVONE. 2010. Redescription of *Syphacia venteli* Travassos, 1937 (Nematoda: Oxyuridae) from *Nectomys squamipes* in Argentina and Brazil and description of a new species of *Syphacia* from *Melanomys caliginosus* in Colombia. Parasitology Research 106:1117-1126.
- ROBLES MR, GT NAVONE, and J NOTARNICOLA. 2006. A new species of *Trichuris* (Nematoda: Trichuridae) from Phyllotini Rodents in Argentina. Journal of Parasitology 92:100-104.
- ROBLES MR, O BAIN, and GT NAVONE. 2012a. Description of a new Capillariinae (Nematoda: Trichuridae) from *Scapteromys aquaticus* (Cricetidae: Sigmodontinae) from Buenos Aires, Argentina. Journal of Parasitology 98:627-639.
- ROBLES MR, C GALLIARI, and GT NAVONE. 2012b. New records of nematode parasites from *Euryzygomatomys spinosus* (Mammalia: Echimyidae) in Misiones province, Argentina. Mastozoología Neotropical 19:353-358.
- ROBLES MR, CJ PERFUMO, JM KINSELLA, and GT NAVONE. 2012c. *Angiostrongylus morerai* from *Akodon* species (Rodentia: Sigmodontinae) from Sierra de la Ventana, Buenos Aires, Argentina: histopathological and parasitological studies. Journal of Parasitology 98:1133-1138.
- ROSSIN MA and AI MALIZIA. 2005. Redescription of *Trichuris pampeana* (Nematoda: Trichuridae) from the South American subterranean rodent *Ctenomys talarum* Thomas, 1898 (Rodentia: Octodontidae). Journal of Parasitology 91:127-130.
- ROSSIN MA, AI MALIZIA, JT TIMI, and R POULIN. 2010. Parasitism underground: Determinants of helminth infections in two species of subterranean rodents (Octodontidae). Parasitology 137:1569-1575.
- SARDELLA NH and MH FUGASSA. 2009. Paleoparasitological analysis of rodent coprolites in holocene samples from Patagonia, Argentina. Journal of Parasitology 95:646-650.
- SORIANO A. 1956. Los distritos florísticos de la Provincia Patagónica. Revista de Investigaciones Agrícolas (Argentina) 10:321- 357.
- SURIANO DM and GT NAVONE. 1994. Three new species of the genus *Trichuris* Roederer, 1761 (Nematoda-Trichuridae) from Caviomorph and Cricetid rodents in Argentina. Research and Reviews in Parasitology 54:39-46.
- TINER JD. 1950. Two new species of *Trichuris* from North America with description of *Trichuris leporis* (Nematoda: Aphasmidia). Journal of Parasitology 36:350-354.
- TRAUB RJ, ID ROBERTSON, P IRWIN, N MENCKE, and RC THOMPSON. 2004. The prevalence, intensities and risk factors associated with geohelminth infection in tea-growing communities of Assam, India. Tropical Medicine and International Health. 9:688-701.

## ONLINE SUPPLEMENTARY MATERIAL

### APPENDIX 1

#### List of examined rodents

[http://www.sarem.org.ar/wp-content/uploads/2014/06/SAREM\\_MastNeotrop\\_21-1\\_Robles-sup1.doc](http://www.sarem.org.ar/wp-content/uploads/2014/06/SAREM_MastNeotrop_21-1_Robles-sup1.doc)