

Development of the mosquito pathogen *Leptolegnia chapmanii* (Straminipila: Peronosporomycetes) on an inexpensive culture medium based on sunflower seed

Manuel E. Rueda-Páramo, Romina G. Manfrino, Alejandra Gutierrez, Claudia C. López-Lastra & Juan J. García

To cite this article: Manuel E. Rueda-Páramo, Romina G. Manfrino, Alejandra Gutierrez, Claudia C. López-Lastra & Juan J. García (2015): Development of the mosquito pathogen *Leptolegnia chapmanii* (Straminipila: Peronosporomycetes) on an inexpensive culture medium based on sunflower seed, *Biocontrol Science and Technology*, DOI: [10.1080/09583157.2015.1118616](https://doi.org/10.1080/09583157.2015.1118616)

To link to this article: <http://dx.doi.org/10.1080/09583157.2015.1118616>



Accepted author version posted online: 18 Nov 2015.



Submit your article to this journal [↗](#)



Article views: 4



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis

Journal: *Biocontrol Science and Technology*

DOI: 10.1080/09583157.2015.1118616

Development of the mosquito pathogen *Leptolegnia chapmanii* (Straminipila: Peronosporomycetes) on an inexpensive culture medium based on sunflower seed

Manuel E. Rueda-Páramo^{a,*}, Romina G. Manfrino^{a,1}, Alejandra Gutierrez^{a,2}, Claudia C. López-Lastra^{a,3}, Juan J. García^{a,4}

^aCentro de Estudios Parasitológicos y de Vectores (CEPAVE), Universidad Nacional de La Plata-CONICET, La Plata, Buenos Aires, Argentina. Boulevard 120 S/N e/61 y 62, CP (B1902CHX), La Plata, Buenos Aires, Argentina

¹ romimanfrino@hotmail.com, ² gutialeja@gmail.com, ³ claudia@cepave.edu.ar, ⁴ juan@cepave.edu.ar

(Received 28 September, 2015; Returned 23 October; Accepted 6 November, 2015)

* Corresponding author: e-mail: ruedapme@gmail.com

An inexpensive culture medium based on sunflower seed extract (SSE) for production of *L. chapmanii* was developed. Vegetative growth on solid and liquid SSE was compared with two culture media used routinely (PYG and Emerson YPss). Results indicate that the oomycete is able to grow on SSE medium, producing more zoospores at a faster rate as well as inducing higher mortality rates in *Ae. aegypti* larvae.

Keywords: Mosquito control, diseases-borne vector, entomopathogen, Straminipila, Oomycete culture.

Leptolegnia chapmanii (Seymour) (Straminipila: Peronosporomycetes) is an aquatic pathogen of mosquito larvae (McInnis & Schimmel, 1985; López Lastra et al., 2004). Its virulence and pathogenicity against *Aedes aegypti* larvae has been studied (McInnis & Zattau, 1982; López Lastra et al., 2004). It persists enabling it to reduce mosquito population for weeks after a single application (Rueda et al., 2015). It grows readily on culture media as PYG and Emerson YPss (Pelizza et al., 2011). However, the use of such culture media for its mass production could be expensive. A culture medium based on sunflower extract (SSE) was reported as an alternative for culture of the

entomopathogen *Lagenidium giganteum* (Couch) (Guzman & Axtell, 1986). The mass production of zoospores from *L. giganteum* maintained on *in vitro* culture media by its immersion in water was used by Jaronski & Axtell, (1983) and a medium-scale production of *L. giganteum* and its application in laboratory and field was reported by Kerwin et al., (1994).

Consequently we developed and evaluated a culture medium based on sunflower seeds as an inexpensive alternative for the mass production of *L. chapmanii*. SSE medium was prepared following the protocol proposed by Jaronski & Axtell (1984). To prepare one liter of medium, ten grams of sunflower seeds were blended with 100 ml distilled water for one minute, filtered through 2-4 layers of cheesecloth and the residue was blended again with 100 ml of distilled water for one minute. Suspensions were filtered, mixed and the final volume was increased with distilled water to one liter. The PYG (Peptone 1.25 g, Yeast extract 1.25 g, Glucose 3 g, distilled water 1000 ml) and E. YPss (Yeast extract 1 g, soluble Starch 20 g, K₂HPO₄ 1g, MgSO₄.7H₂O 0.5 g, distilled water 1000 ml) culture media were routinely prepared. Media were solidified adding 15 g of agar per liter of culture medium. Both liquid and solid culture media were sterilized by autoclaving at 120 °C and 20 PSI for 25 minutes.

Growth and development of *L. chapmanii* in solid and liquid SSE culture media was compared with traditional media PYG and E. YPss, used for its culture. Zoospore production and their pathogenicity against *A. aegypti* larvae were also evaluated. An Argentinean isolate of *L. chapmanii* (CEP 010) from the Fungal Entomological Collection of the Centro de Estudios Parasitológicos y de Vectores (CEPAVE) in La Plata, Buenos Aires, Argentina was used. *Aedes aegypti* larval populations (L2/L3) from a colony established at CEPAVE were used in pathogenicity bioassays

Petri dishes (100 mm diam.) with solid media were used to evaluate radial growth of *L. chapmanii*. Treatment medium dishes were inoculated by placing 1cm² plug from PYG solid medium with *L. chapmanii*. The treatments were maintained at 25 °C in an incubator and the radial growth was assessed after 48, 72 and 96 h. Each treatment had four replicate dishes and the assay was repeated three times. The formation and release of zoospores was induced by adding five pieces (1cm²) of each solid medium treatment into 100 ml distilled water. Zoospore concentrations were determined after 48 h and their viability and pathogenicity evaluated through bioassay against *A. aegypti* larvae (n=25). Larval mortalities were recorded after 48 h.

L. chapmanii growth in liquid media was evaluated by initial inoculating of 200 ml of each liquid medium with a 1 cm² plug from cultures grown on of the PYG solid medium, incubating for 7 days at 25°C and 0.18 g (180 rpm) in an orbital shaker, then transferred to 400 ml of each respective culture medium for an additional 7 days of culture under the same laboratory conditions. After the 14-day culture period, biomass from each culture medium was filtered and fresh weight recorded. Cost of production based on biomass obtained in different culture media was estimated and compared. For zoospore formation and release, 10 g of biomass produced in liquid cultures were added to 200 ml sterile distilled water in 500 ml Erlenmeyer flasks on and shaken on an orbital shaker at 0.18 g (180 rpm). Production of zoospores was determined daily for 4 days. Their viability and virulence were confirmed by pathogenicity tests against *A. aegypti* larvae (n= 10) using 10 ml of the zoospore suspensions in plastic containers

(50 ml). Larval mortality was recorded after 48 h. Each treatment had three replicate containers and one negative control with distilled water. Assays were repeated three times at different dates.

In all cases, the concentration of zoospores was determined using a Neubauer chamber. Results were analyzed by ANOVA and the *pos hoc* test of Student Newman Keuls (SNK). Larval mortalities were transformed to arsine square-roots for statistical analysis. The SPSS statistics software from IBM was for the analyses.

Leptolegnia chapmanii developed on solid culture media with different growth rates. Higher production of zoospores as well as mortalities against *Ae. aegypti* larval population were obtained with *L. chapmanii* from the alternative SSE medium (Table 1A). It grew also in liquid culture media developing a great number of small mycelial masses in SSE medium compared with a single large biomasses formed in PYG and E₁ YPss. Higher production of zoospores (Fig 1) and host mortality (Table 1 B) was obtained with *L. chapmanii* from SSE medium. Nulls mortalities were registered on negative controls.

Even though the production of zoospore by *L. chapmanii* had been described without sterol enrichment of culture medium (Pelizza et al., 2007), the present results suggest that *L. chapmanii* could be an auxotrophic organism that develops vegetatively on poor sterol media, but requires it for initiation of the reproductive cycles (sexual / asexual) as described for *L. giganteum* (Domnas et al., 1977; Kerwin & Washino, 1983). An increased production of zoospores by the enrichment of culture medium with sunflower oil has also been demonstrated for *L. giganteum* (Balaraman & Hoti, 1986; Maldonado-Blanco et al., 2011). Sunflower seeds are a great source of phytosterols (Merah et al., 2012). Higher mortalities of *Ae. aegypti* larvae were related with higher zoospore concentrations (Pelizza, 2007).

We conclude that *L. chapmanii* was able to grow on SSE solid and liquid medium. Based on local prices of compounds required to prepare the culture media used, to produce 1 kg of *L. chapmanii* biomass in PYG would have a cost of \$US10, \$7 with and \$3 with SFE liquid culture media. This relatively inexpensive medium providing the nutritional requirements for the vegetative and reproductive development of the pathogen.

Acknowledgements

This study was partially supported by CONICET, Universidad Nacional de La Plata, and ANPCYT (PICT 0622-2012). The senior author was granted by a doctoral fellowship from CONICET.

References

- Balaraman, K., & Hoti, S. L. (1986). Factors influencing *in vitro* zoosporogenesis of mosquito pathogenic fungus *Lagenidium*. *Indian Journal of Medical research*, 83, 380-383.
- Domnas, A. J., Srebro, J. P., & Hicks B. F. (1977). Sterol requirement for zoospore formation in the mosquito – parasitizing fungus *Lagenidium giganteum*. *Mycologia*, 69, 875-886.

- Guzman, D. R., & Axtell, R. C. (1986). Effect of nutrient concentration in culturing three isolates of the mosquito fungal pathogen, *Lagenidium giganteum* Couch, on sunflower seed extract. *Journal of American Mosquito Control Association*, 2, 196-200.
- Jaronski, S. T., & Axtell, R. C. (1983). Persistence of the mosquito fungal pathogen *Lagenidium giganteum* (oomycetes: Lagenidiales) after introduction into natural habitats. *Mosquito News*, 43, 332-337.
- Jaronski, S.T., & Axtell, R. C. (1984). Simplified production system for the fungus *Lagenidium giganteum* for operational mosquito control. *Mosquito News*, 44, 377-381.
- Kerwin, J. L., & Washino, R. K. (1983). Sterol induction of sexual reproduction in *Lagenidium giganteum*. *Experimental Mycology*, 7, 109-115.
- Kerwin, J. L., Dritz, D. A., & Washino, R. K. (1994). Pilot scale production and application in wildlife ponds of *Lagenidium giganteum* (Oomycetes: Lagenidiales). *Journal of American Mosquito Control Association*, 10, 451-455.
- López Lastra, C. C., Scorsetti, A. C., Marti, G. A., & García, J. J. (2004). Host Range and Specificity of an Argentinean Isolate of the Aquatic Fungus *Leptolegnia chapmanii* (Oomycetes: Saprolegniales), a pathogen of mosquito larvae (Diptera: Culicidae). *Mycopathologia*, 158, 311-315.
- Maldonado-Blanco, M. G., Leal-López, E. Y., Ochoa-Salazar, O. A., Santos, M. E., Galán-Wong, L. J., & Quiroz-Martínez, H. (2011). Effects of culture medium and formulation on the larvicidal activity of the mosquito pathogen *Lagenidium giganteum* (Oomycetes: Lagenidiales) against *Aedes aegypti*. *Acta Tropical*, 117, 114-118.
- McInnis, T. M., & Zattau, W. C. (1982). Experimental infection of mosquito larvae by a species of the aquatic fungus *Leptolegnia*. *Journal of Invertebrate Pathology*, 39, 98-104.
- McInnis, T. M., & Schimmel, L. E. (1985). Host range studies on a new fungus *Leptolegnia*, a parasite of mosquito larvae (Diptera: Culicidae). *Journal of Medical Entomology*, 22, 226-227.
- Merah, O., Langlade, N., Alignan, M., Roche, J., Pouilly, N., Lippi, Y., Vear, F., Cerny, M., Bouniols, A., Mouloungui, Z., & Vincourt, P. (2012). Genetic analysis of phytosterols content in sunflower seeds. *Theoretical and Applied Genetics*, 125, 1589-1601.
- Pelizza, S. A. (2007). Estudio y evaluación de *Leptolegnia chapmanii* Seymour (Oomycetes, Saprolegniales) como agente de control biológico de *Aedes aegypti* (L.) (Diptera: Culicidae). PhD Dissertation, Universidad Nacional de La Plata.
- Pelizza, S. A., López Lastra, C. C., Becnel, J. J., Bisaro, V., & García, J. J. (2007). Biotic and abiotic factors affecting *Leptolegnia chapmanii* infection in *Aedes aegypti* L. (Diptera: Culicidae). *Journal of American Mosquito Control Association*, 23, 177-181.

Pelizza, S. A., Cabello, M. N., Tranchida, M. C., Scorsetti, A. C., & Bisaro, V. (2011). Screening for a culture medium yielding optimal colony growth, zoospore yield and infectivity of different isolates of *Leptolegnia chapmanii* (Straminipila: Peronosporomycetes). *Annals of Microbiology*, 61, 991-997.

Rueda Páramo, M. E., López Lastra, C. C., & García, J. J. (2015). Persistence and pathogenicity of a native isolate of *Leptolegnia chapmanii* against *Aedes aegypti* larvae in different anthropic environments. *Biocontrol Science & Technology*, 25, 238-243.

Table 1. A) Assays with *L. chapmanii* developed on solid culture media. B) Assays with *L. chapmanii* increased in liquid culture media.

A.						B.				
On solid culture media						In liquid culture medium				
Medium	Radial Growth (mm) *			Zoospore s.ml ⁻¹	Mortality of <i>Ae. aegypti</i> larvae** (%)	Fresh mass (g)	Mortality of <i>Ae. aegypti</i> larvae † (%)			
	48 h	72 h	96 h				24 h [†]	48 h [†]	72 h [†]	96 h [†]
PYG	18 ± 0.6 ^a	31 ± 1.0 ^a	40 ± 1.6 ^a	583 ± 193	53 ^b	48.5	94 ± 4.0 ^a	99 ± 1.0 ^a	100 ± 10 ^a	100 ± 5 ^a
E.	18 ± 0.2 ^a	30 ± 0.5 ^a	41 ± 0.9 ^a	1250 ± 329	67 ^b	93.5	0 ^b	31 ± 16 ^b	10 ± 6 ^c	5 ± 3 ^c
YPss	10 ± 0.5 ^b	17 ± 0.7 ^b	26 ± 0.7 ^b	2083 ± 668	86 ^a	62.4	0 ^b	25 ± 13 ^b	57 ± 16 ^b	93 ± 7 ^b
SSE	0.5 ^b	0.7 ^b	0.7 ^b	668	86 ^a	4	0 ^b	13 ^b	16 ^b	7 ^b

† Time after zoosporogenesis induction by immersion of *L. chapmanii* in water.

Note: Letters grouped the treatments without statistical differences.

* $F_{2,33}(48h) = 98$, $F_{2,33}(72h) = 97$ and $F_{2,33}(96h) = 57$.

In all cases $P < 0.001$

** $F_{2,51} = 6.6$, $P < 0.05$

† $F_{2,24}(24h) = 331$, $F_{2,24}(48h) = 17$, $F_{2,24}(72H) = 59$ and $F_{2,24}(96h) = 192$. In all cases $P < 0.001$




Figure 1. Assays with biomass samples from liquid culture media. Production of zoospores by *Leptolegnia chapmanii* along time.

Accepted Manuscript