

First records of leaf-footed bugs (Heteroptera: Coreoidea: Coreidae) from Argentina

OLIVERA, Leonela, MELO, María C. & DELLAPÉ, Pablo M.

Universidad Nacional de La Plata, CONICET, División Entomología, Museo de La Plata, Paseo del Bosque s/n B1900FWA, La Plata, Buenos Aires, Argentina. E-mail: lolivera@fcnym.unlp.edu.ar

Received 05 - IV - 2017 | Accepted 25 - V - 2017 | Published 29 - VI - 2017

<https://doi.org/10.25085/rsea.761212>

Primeros registros de chinches foliadas (Heteróptera: Coreoidea: Coreidae) de la Argentina

RESUMEN. La familia Coreidae es un grupo de chinches fitófagas, con una distribución cosmopolita y mayor diversidad en áreas tropicales y subtropicales. El conocimiento de los Coreidos argentinos es aún incompleto, en este trabajo registramos de la Argentina las siguientes nueve especies: *Althos distinctus* (Signoret), *Cnemyrtus scriptus* (Burmeister), *Himella venosa* Dallas, *Nyttum beckeri* Dolling & Casini, *Ugnius badius* (Herrich-Schäffer), *Vilga (Echinovilga) dallasii* Distant, *Vilga (Laevivilga) sanctipauli* Dolling, *Vilga (Vilgula) obliqua* Dolling y *Zicca pronotata* Brailovsky & Cadena.

PALABRAS CLAVE. Anisoscelini. Hypselonotini. Nematopodini. Placoscelini. Pseudophloeini.

ABSTRACT. The family Coreidae is a diverse group of phytophagous Heteroptera that show a cosmopolitan distribution although they are more diverse in tropical and subtropical areas. The knowledge of the diversity of Argentinean Coreidae is incomplete; in this contribution we report the following nine species from Argentina: *Althos distinctus* (Signoret), *Cnemyrtus scriptus* (Burmeister), *Himella venosa* Dallas, *Nyttum beckeri* Dolling & Casini, *Ugnius badius* (Herrich-Schäffer), *Vilga (Echinovilga) dallasii* Distant, *Vilga (Laevivilga) sanctipauli* Dolling, *Vilga (Vilgula) obliqua* Dolling and *Zicca pronotata* Brailovsky & Cadena.

KEYWORDS. Anisoscelini. Hypselonotini. Nematopodini. Placoscelini. Pseudophloeini.

The family Coreidae is a diverse group of terrestrial Heteroptera with a cosmopolitan distribution and greater diversity in tropical and subtropical areas (Schuh & Slater, 1995). The species included in the family are mainly phytophagous, and many show definite associations with particular plants, making them important crop pests (Mitchell, 2000). The family contains about 519 genera and about 3,098 species worldwide (Coreoidea SF Team), of which 158 genera are found in the Neotropical region. All three subfamilies (Pseudophloeinae, Merophachyinae and Coreinae) and 16 tribes are represented in the region (Henry, 2009; Fernandez et al., 2015).

The Neotropical Coreidae fauna has been extensively studied by Brailovsky, but the knowledge of the diversity of Argentinean Coreidae is far from complete, mainly

due to the absence of South American specialists. The major contributions to Argentinean fauna were done by Pennington (1920-1921, 1922) and by Kormilev (1954), and by Brailovsky's works that studied many species from Argentina, particularly in the context of systematic revisions. According to the literature, there are about 140 species recorded from Argentina, but this number is far from the real diversity of the country.

In this paper, we present nine first records for coreid species belonging to the following tribes: Pseudophloeinae: Pseudophloeini (3 sp.), Coreinae: Anisoscelini (1 sp.), Hypselonotini (2 sp.), Nematopodini (2 sp.) and Placoscelini (1 sp.).

The specimens were examined under an Olympus SZ stereomicroscope; to confirm identification of some specimens, the genital capsule of males and genital

segments of females were dissected and cleared with a saturated potassium hydroxide solution for observation. The dissected genitalia were subsequently preserved in microvials with glycerin. The locations reported for study materials refer to the political division of Argentina into provinces; other countries where the species have been recorded in the literature are mentioned. The material examined belongs to the entomological collection of Museo de La Plata, Buenos Aires, Argentina (MLP). For each species the Life Science Identifier (LSID) was added, to link with the Coreoidea Species File website (CoreoideaSF Team).

Below we present the list with the new species registered from Argentina.

Subfamily Pseudophloeinae

Tribe Pseudophloeini

Vilga (Echinovilga) dalasi Distant, 1881

urn:lsid:Coreoidea.speciesfile.org:TaxonName:458571

Geographic distribution: Brazil, French Guiana, Guatemala and Paraguay (Berg 1894; Dolling, 1977; Packauskas, 2010).

Studied material: ARGENTINA. Misiones: 1 male, Campos S.A., 8-III-2010, P.M. Dellapé.

Vilga (Laevilgilga) sanctipauli Dolling, 1977

urn:lsid:Coreoidea.speciesfile.org:TaxonName:458574

Geographic distribution: Brazil (Dolling, 1977; Packauskas, 2010).

Studied material: ARGENTINA. Misiones: 1 male, NW de Apóstoles, 27° 52.451'S 55° 39.514'W, 146 msnm, 8-III-2010, P.M. Dellapé.

Vilga (Vilgula) obliqua Dolling, 1977

urn:lsid:Coreoidea.speciesfile.org:TaxonName:458591

Geographic distribution: Brazil (Dolling, 1977; Packauskas, 2010).

Studied material: ARGENTINA. Misiones: 1 male, Campos S.A., 8-III-2010, P.M. Dellapé.

Subfamily Coreinae

Tribe Anisoscelini

Ugnius badius (Herrich-Schäffer, 1851)

urn:lsid:Coreoidea.speciesfile.org:TaxonName:457564

Geographic distribution: Brazil, Bolivia and Paraguay (Herrich-Schäffer, 1851, 1853; Packauskas, 2010).

Studied material: ARGENTINA. Misiones: 1 female, San Ignacio, 8-XII-1941, Birabén-Bezzi; 1 female, RP2, 20 km NE El Soberbio, 27°12.220'S- 54°01.375'W, 326m, 5-III-2010, G. del Río; 1 male San Ignacio, 8-XII-1941, Birabén-Bezzi; 1 male, XI-1941; 1 male, Alta Paraná Sociedad Anónima, 26°01'47.9"S- 54°6'3.2"W, 22-II-2009, L. Negrete

Tribe Hypselonotini

Althos distinctus (Signoret, 1863)

urn:lsid:Coreoidea.speciesfile.org:TaxonName:454369

Geographic distribution: Chile (Brailovsky, 1990).

Studied material: ARGENTINA. Río Negro: 12 females, Bariloche, Margen del Río Limay, 16-II-2015, S. Montemayor; 1 male, Bariloche, Río Limay, 16-II-2015, S. Montemayor

Zicca pronotata Brailovsky & Cadena, 1992

urn:lsid:Coreoidea.speciesfile.org:TaxonName:454146

Geographic distribution: Brazil (Brailovsky & Cadena, 1992).

Studied material: ARGENTINA. Misiones: 1 female, Reserva de Vida Silvestre Urugua-í, sendero Pozones, 25°59'01"S- 54°06'51"W, 31-X-2010, L. Negrete

Tribe Nematopodini

Cnemyrtus scriptus (Burmeister, 1835)

urn:lsid:Coreoidea.speciesfile.org:TaxonName:453109

Geographic distribution: Brazil and Paraguay (O'Shea, 1980).

Studied material: ARGENTINA. Buenos Aires: 1 female, Sierra de La Ventana, XI-1970, A.A. Pirán. Chaco: 1 female, Charata, X-1924, Bosq; 1 female, Resistencia, 26-IV-1938; 3 males, Charata, X-1924, Bosq. Córdoba: 1 male, Dep. De Calamuchita, "El Sauce", XII-1938, M.I. Viana.

Himella venosa Dallas, 1852

urn:lsid:Coreoidea.speciesfile.org:TaxonName:453099

Geographic distribution: Brazil (O'Shea, 1980; Brailovsky & Barrera, 1986).

Studied material: ARGENTINA. Misiones: 1 female, Reserva Natural Estricta San Antonio, 7-V-2008, L. Negrete; 1 male, Reserva Natural Estricta San Antonio, 7-V-2008, L. Negrete.

Tribe Placoscelini

Nyttum beckeri Dolling & Casini, 1988

urn:lsid:Coreoidea.speciesfile.org:TaxonName:454481

Geographic distribution: Brazil and West Indies (Dolling & Casini, 1988; Packauskas, 2010).

Studied material: ARGENTINA. Misiones: 1 male, Parque Provincial Urugua-í, 25°51.464"S- 54°10.139'W, 276m, 7-III-2010, P.M. Dellapé.

ACKNOWLEDGEMENTS

We thank Lisandro Negrete (División Invertebrados, Museo de La Plata) for kindly collecting bugs during his field trips. This study was supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina, and the following grants: PIP 0249 (2013-2015) and UNLP N784.

BIBLIOGRAFÍA CITADA

Berg, C. (1894) Descripciones de algunos Hemípteros Heterópteros nuevos o poco conocidos. *Anales del Museo Nacional de Montevideo*, 1, 13-27.

- Brailovsky, H. (1990) Revisión del complejo "Althos" (Hemiptera-Heteroptera-Coreidae). *Instituto de Biología de la Universidad Nacional Autónoma de México, Publicaciones Especiales*, 5, 1-156.
- Brailovsky, H., & Barrera, E. (1986) El género *Himella* Dallas con descripción de dos especies nuevas (Hemiptera-Heteroptera-Coreidae-Nematopodini). *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 2, 423-436.
- Brailovsky, H., & Cadena A. (1992) Revisión del género *Zicca* (Hemiptera-Heteroptera-Coreidae-Coreinae-Coreini). *Instituto de Biología de la Universidad Nacional Autónoma de México, Publicaciones Especiales*, 9, 1-101.
- Burmeister, H.C. (1835) *Handbuch der Entomologie. Zweiter Band: II Ordung Rhynchota*. Enslin, Berlin, Germany.
- CoreoideaSF Team. Coreoidea Species File Online. Version 5.0/ 5.0. [November 22 2016].
- Dallas, W.S. (1852) Description of a new hemipterous insect forming the type of a new genus. *Annals and Magazine of Natural History*, 10(2), 359-362.
- Distant, W.L. (1881) Insecta. Rhynchota, Hemiptera-Heteroptera. *Biología Centrali Americana Vol 1* (eds. Godman, F.D., & Salvin, O.), pp. 89-168. London.
- Dolling, W.R. (1977) A revision of the Neotropical genus *Vilga* Stål (Hemiptera: Coreidae). *Systematic Entomology*, 2, 27-44.
- Dolling, W.R., & Casini, C.E. (1988) Revision of the Neotropical genus *Nyttum* (Hemiptera: Coreidae). *Systematic Entomology*, 13, 143-156.
- Fernandes, J.A.M., Mitchell, P.L., Livermore, L., & Nikunlassi, M. (2015) Chapter 19. Leaf-Footed Bugs. *True Bugs (Heteroptera) of the Neotropics*, (eds. Panizzi, A.R., & Grazia, J.), pp. 549-605. Springer, Dordrecht, Heidelberg, New York, London.
- Henry, T.J. (2009) Chapter 10. The Biodiversity of True Bugs. *Insect Biodiversity: Science and Society* (eds. Foottit, R., & Adler, P.), pp. 223-263. Blackwell publishing, Chichester, Hoboken, Oxford.
- Herrich-Schäffer, G.A.W. (1851) Die wanzenartigen Insecten: getreu nach der Natur abgebildet und beschrieben/von Dr. Carl Wilhelm Hahn. *Die wanzenartigen Insecten*, 9(6), 257-348.
- Herrich-Schäffer, G.A.W. (1853) Die wanzenartigen Insecten: getreu nach der Natur abgebildet und beschrieben/von Dr. Carl Wilhelm Hahn. *Die wanzenartigen Insecten*, index 1-210.
- Kormilev, N. (1954) Notas sobre Coreidae neotropicales II; (Hemiptera) Merocorinae de la Argentina y países limítrofes. *Revista Ecuatoriana de Entomología y Parasitología*, 2, 153-187.
- Mitchell, P.L. (2000) Chapter 11. Leaf-footed bugs (Coreidae). *Heteroptera of Economic Importance* (eds. Schaefer, C.W., & Panizzi, A.R.), pp. 337-403. CRC Press, Boca Raton, London, New York, Washington, D.C.
- O'Shea, R. (1980) A generic revision of the Nematopodini (Hemiptera: Coreidae: Coreinae). *Studies on Neotropical Fauna and Environment*, 15, 197-225.
- Packauskas, R. (2010) Catalog of the Coreidae, or leaf-footed bugs of the New World. *Fort Hays Studies (Fourth Series)*, 5, 1-270.
- Pennington, M.S. (1920-1921) *Lista de los Hemípteros Heterópteros de la República Argentina*. Buenos Aires, Argentina.
- Pennington, M.S. (1922) Notas sobre Coreidos Argentinos. *Physis*, 20(5), 125-170.
- Schuh, R.T., & Slater, J.A. (1995) *True Bugs of the World (Hemiptera: Heteroptera). Classification and Natural History*. Cornell University Press, Ithaca, USA.
- Signoret, V. (1863) Révision des Hémiptères du Chili. *Annales de la Société Entomologique de France*, 3(4), 541-588.