

## MORE THAN TRACKS: HOW THE CYTOSKELETON BUILDS EPITHELIA ASYMMETRY

Evangelina Almada<sup>1</sup>, Facundo Tonucci<sup>1</sup>, Alejandro Pariani<sup>1</sup>, Florencia Hidalgo<sup>1</sup>,  
Cristián Favre<sup>1</sup>, M. Cecilia Larocca<sup>1\*</sup>

<sup>1</sup>Instituto de Fisiología Experimental, Consejo de Investigaciones Científicas y Técnicas, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario. 2000-Rosario, Argentina

\*Correspondence to:

M. Cecilia Larocca ([larocca@ifise-conicet.gov.ar](mailto:larocca@ifise-conicet.gov.ar))

### ABSTRACT

Epithelia are three-dimensional arrangements of cells organized into structures which delimit morphologically and physiologically different compartments. At the cellular level, epithelial cells are characterized by the presence of an apical pole, facing the lumen of an organ or the outer environment, and baso-lateral domains, involved in cell-extracellular matrix and cell-cell interactions. Each pole is characterized by its specific protein and lipid membrane composition, and its distinct assembly of cellular organelles. This polarized organization of cell components ensures the vectorial absorption and secretion of proteins and other molecules, which constitute the essential function of epithelia. Actin and microtubule cytoskeletons are also polarized in epithelial cells. This organization contributes to the *de novo* establishment as well as to the functional maintenance of epithelial polarity. In this article we delineate the central features of epithelial polarity and summarize the current knowledge on how actin and microtubule cytoskeletons contribute to epithelial organization.

**Key words:** actin, microtubules, epithelial polarity

### RESUMEN

Los epitelios son arreglos tridimensionales de células, organizados en estructuras que delimitan compartimientos morfológica y fisiológicamente diferentes. A nivel celular, las células epiteliales se caracterizan por la presencia de un polo apical, en contacto con la luz de un órgano o con el medio extracelular, y los dominios baso-laterales, responsables de las interacciones célula-matriz extracelular y célula-célula. Cada polo se caracteriza por la composición proteica y lipídica específica de sus membranas, así como por el ensamble distintivo de las organelas celulares asociadas. Esta organización polarizada de los componentes celulares garantiza el desarrollo de la función esencial de los epitelios, i.e. la absorción y/o secreción vectorial de proteínas y otras moléculas. Los citoesqueletos de actina y microtúbulos también están polarizados en las células epiteliales, lo que contribuye tanto al establecimiento *de novo*, como al mantenimiento funcional de la polaridad epitelial. En este artículo delineamos los aspectos centrales de la polaridad epitelial y resumimos el estado de conocimiento actual en relación a la contribución de los citoesqueletos de actina y microtúbulos a la organización epitelial.

**Palabras clave:** actina, microtúbulos, polaridad epitelial.

Original received: February 7, 2018; Accepted in its final form: March 21st, 2018.


## Introduction

An epithelium (the trophectoderm) is the first tissue that differentiates during embryogenesis [1]. The epithelium is also the first tissue that emerged during phylogenesis. In fact, the initial epithelial structures precede metazoan multicellularity: upon starvation, individual *Dictyostelium discoideum* social amoebas aggregate to form a fruiting body, which includes a tubular epithelial monolayer that surrounds a collection of spores [2]. Both that primitive epithelia and modern epithelia share their function of segregating an internal media from an external environment, what is tightly associated to their coordinated asymmetric distribution of cellular components, and the presence of cell-cell junctions with selective permeability (Figure 1). This organization ensures the vectorial absorption and secretion of proteins and other molecules, which constitute the essential function of epithelia. The establishment and maintenance of that organization imply two types of cell asymmetry: apico-basal polarity, which refers to the distribution of cell components within each individual cell, and planar polarity, which denotes the collective alignment of cell polarity across the tissue plane [3]. Over the last decades, studies performed in *Caenorhabditis elegans*, *Drosophila melanogaster* and mammalian epithelial cells have contributed to the understanding of the network of polarity proteins and lipids that defines the differentiation of distinct cell poles. These investigations identified three basic polarity protein complexes which are conserved throughout evolution: the crumbs (Crumbs/PALS1/PATJ) complex, the PAR (PAR3/PAR6/aPKC/cdc42) and the related PAR4/LKB1 complex, and the Scribble (Scribble/Discs-large (Dlg)/Lethal giant larvae (LGL)) complex. Most of these studies have been nicely reviewed [4-6]. Moreover, several studies have focused on how the interference with the functionality of these networks can lead not only to loss of epithelial function, but also to uncontrolled cell proliferation and development of epithelial derived malignant tumors [7]. An important question to address in order to understand epithelial homeostasis is how polarity signals, trafficking events, and cell cycle issues are integrated to define epithelial architecture. Traditionally, the cytoskeleton and its associated motor proteins were viewed to be passive actors in the construction of cell architecture, functioning downstream regulatory molecules that have already been localized by pre-existing spatial cues. However, results from different model systems have supported the idea that the cytoskeleton also drives the symmetry-breaking process by localizing key regulatory molecules to specific cortical sites [8]. In the present article we delineate the main characteristics of the epithelial polarity and recapitulate the current knowledge on the mechanisms by which the actin and microtubule cytoskeletons contribute to the integration of the different factors that determine epithelial organization.

## Epithelial polarity

At the cellular level, epithelial polarity is characterized by the presence of an apical pole, facing the lumen of an organ or the outer environment, a lateral domain, involved in cell-cell interactions, and the basal pole, in contact with the extracellular matrix (ECM). Most epithelial cells develop a columnar type of apico-basal polarity, which implies the organization of a simple apical pole that opposes to the basal domain (Figure 1). Multi-layered epithelia, such as the skin epidermis, develop this type of polarity, but restricted only to the most apical layer of viable cells, whereas the most basal cells are non-differentiated and proliferative [9]. On the other hand, hepatocytes develop a particular type of polarity, where there is more than one apical domain, giving rise to multiple “bile canaliculus” per cell [10]. In any case, the establishment of cell-cell, and cell-ECM interactions constitute the founding events in the acquisition of the epithelial polarity. In vertebrates, cell-cell junctions include tight junctions, adherent junctions, gap junctions and desmosomes. Tight and adherent junctions, also denominated “apical junction complex”, are particularly important for the formation and the

maintenance of apico-basal polarity. Formation of the apical junction complex depends on the polarity complexes Crumbs and PAR. The latter protein complex determines the localization of the apical pole, by mutually excluding the “basolateral” scribble complex [4]. Thereafter, apical cellular junctions contribute to the segregation of the apical and basolateral poles. Whereas apical junctions contribute to the specification of the apical pole location, the cell-ECM interaction determines the polarity axis orientation. An initial event is the interaction of  $\beta$ 1-integrins with the extracellular matrix, what leads to the assembly of a protein complex at the cytosolic integrin domain that transduce the signals originated at the ECM to control cytoskeleton dynamics and intracellular signaling [11,12]. The proteins associated to the cellular junctions (Figure 3) as well as  $\beta$ 1-integrins are also involved in the specification of epithelial cell division axis by determining the mitotic spindle orientation [13-15]. Both the establishment of cell-cell and cell-ECM interactions and the derived intracellular signaling that ensures the formation of the apico-basal asymmetry depend on the association of those junctions with the microtubule and actin cytoskeleton.




**Figure 1.** Epithelial organization. The image shows a three dimensional reconstruction of F-actin (red),  $\alpha$ -tubulin (microtubules, green) and DAPI (nucleus, blue) staining of the prototype epithelial cells MDCK cultured to polarize on filters (A). Arrowheads indicate microtubule staining at the primary cilia. Arrows specify actin organization at the brush border. The diagram above the image represents the hexagonal prism organization of cells at the epithelial monolayer. The schemas illustrate the polarized distribution of the actin and microtubule cytoskeleton, the Golgi apparatus, the centrosome and the endosomal compartments (B), and the polarity and apical junction complexes (C) [4,5]. BSE, basal sorting endosomes; CE, common endosomes; ASE, apical sorting endosomes, ARE, apical recycling endosomes, ERM, ezrin-radixin-moesin.

### Acting in the epithelia polarity

Actin filaments are polymers of actin subunits organized in a polarized lattice with two structurally distinct ends: barbed (plus) ends, enriched in GTP bound actin, and pointed (minus) ends, which contain ADP bound actin and exhibits a lower rate of actin incorporation. In differentiated epithelia, these intrinsically polarized filaments organize different arrays at each cell domain (Figure 2), where they participate in processes as diverse as the formation of the apical microvilli, the development of the cellular junctions, the interaction of the basal structures with the ECM, and the regulation of vesicle sorting at the Golgi apparatus and

endosomal compartments. Furthermore, in association with myosin motors, actomyosin contraction drives 3D epithelial tissue organization by generating forces at the cellular level. The latter phenomena have been nicely reviewed elsewhere [16]. In this review we will focus on actin organization and function in apico-basal cell polarity.

Microvilli are cylindrical cell protrusions present at the apical pole of epithelial cells, which increase the cell surface available for apical secretory or absorptive transport. In most epithelial cells which coat hollow organs, these apical protrusions constitute the brush border (Figure 2). A very well-characterized example is the small intestinal cell microvilli. In this case, up to 1000 microvilli protrude from a single apical cell surface. This tight organization of membrane protrusions is enabled by the highly ordered patterning of the actin cytoskeleton, which is organized as parallel bundles of 10-30 filaments situated at the core of each protrusion with the barbed ends facing the plasma membrane at the microvillus tip. These actin bundles are assembled together by crosslinking proteins such as villin, fimbrin or fascin, and interact with the membrane by means of a group of myosin motors, which include myosin-1a, myosin-6 and myosin-7a/b, and proteins of the ezrin/radixin/moesin family [17]. The polymerization and bundling of these actin arrays provide the force to generate microvillar membrane protrusions [18]. Beyond their role in the genesis of the microvilli, actin filaments constitute tracks for myosin dependent trafficking of apical transporters along these structures [19, 20], and participates in apical receptor mediated endocytosis and protein internalization [21].



**Figure 2.** Actin cytoskeleton in epithelial cells. Images show monolayer (A) or single cell (B) 3D reconstructions of F-actin staining of polarized MDCK cells, and single  $xy$  planes of the same cell at the brush border (C) and apical junction (D) level. The schema illustrates actin (red beads) interaction with apical junction complexes.

The perijunctional actin cytoskeleton comprises actin bundles of filaments which are associated to the apical junctions by directly interacting with E-cadherin associated catenins [22] and occludin associated zonula occludens 1 (ZO-1) [23] (Figure 2). The acto-myosin and junctional complexes form a belt-like ring around the cell perimeter which is responsible for the junctional tension, and can produce contraction of the microvilli. During the initial establishment of apical junctions, actin filaments promote the formation of filopodia which

penetrate into neighboring cells, catalyzing the clustering of apical junction proteins such as E-cadherin and catenins [24]. At this stage, cdc42, Rac1 and Rho GTPases are responsible for recruiting a cohort of proteins to the site of cell-cell contact, which promote actin nucleation and remodeling and acto-myosin contraction, leading to junction maturation [25]. In mature epithelia, the junctional actin cytoskeleton contributes to the regulation of apical junction's stability. On one side, actin nucleation induced by the cdc42/WASP/ARP2/3 complex contributes to E-cadherin endocytosis therefore ensuring adherent junction's remodeling [26]. Additionally, acto-myosin contractility induced by RhoA maintains cell-cell junction tension [27]. Besides its role in promoting apical junctions formation and dynamics, the perijunctional actin cytoskeleton interaction with ZO-1 conditions the formation of organotypic epithelial structures with a single lumen by regulating the symmetric orientation of the mitotic spindle in dividing cells [28].

In the basal surface of epithelial cells cultured on a rigid surface, actin microfilaments organize as stress fibers that reach the plasma membrane through their indirect interactions with integrins. Studies performed in three-dimensional epithelial cells showed that the interaction of  $\beta$ 1-integrin with biochemically and biophysically defined ECMs can initiate a signaling pathway that decreases RhoA activation. RhoA inhibition suppresses peripheral actomyosin contraction, leading to downregulation of focal adhesions and centrosome relocation to the opposite pole of the cell, thus ensuring the correct positioning of the apical domain [29-31]. Thus, basal actin cytoskeleton acts as a mechano-transducer which participates in the sensing of the ECM during epithelial cell differentiation.

Besides the distinct arrangements of actin microfilaments associated to plasma membrane, there are also subsets of actin filaments associated to the different cellular organelles. For instance, it has been described the presence of a population of short-branched actin filaments specifically concentrated around the Golgi complex, which associates with budding vesicles [32]. These findings are in agreement with different studies which described that the interference with actin regulatory proteins or with actin dynamics affects the exit of apical and basolateral proteins from the trans-Golgi network in different ways [33, 34]. Even though the mechanisms involved in the organization of these filaments or contributing to the generation of specific cargo routes exiting the Golgi complex are far from being elucidated, Rodriguez Boulan and col. characterized the presence of a dynamic population of actin filaments at the Golgi region that shows remarkable specificity in promoting the dynamin-mediated fission of carrier vesicles for apical but not for basolateral markers [35].

### **Microtubules in epithelial polarity**

Microtubules are protofilaments composed of  $\alpha$ - and  $\beta$ -tubulin heterodimers which associate laterally to form hollow tubes. Microtubule filaments are themselves polarized structures that harbor plus and minus ends, characterized by their distinct dynamic instability. Microtubule nucleation occurs at the minus end, and is mostly dependent on  $\gamma$ -tubulin ring complexes, to which they can remain attached. Conversely, microtubule plus ends are subject to continuous polymerization and depolymerization events, regulated by their associated stabilizing and severing proteins [36]. Microtubule stability is also regulated by tubulin post-translational modifications, which condition the affinity and activity of microtubule associated proteins [37].

Whereas in non-differentiated cells microtubule minus ends are generally anchored to a centrally positioned centrosome, thus displaying a radial array, in differentiated epithelial cells microtubules mostly organize in vertical arrays, with their minus ends anchored to alternative microtubule anchoring proteins located at the apical membrane and at apical junctions, and their plus ends oriented towards the basal membrane (Figure 3) [38-40]. In these cells the centrosome also localizes apically, where it constitutes the basal body of the primary cilia


[41], but does not function as a microtubule organizing center [42]. Studies performed in three dimensional cultures of breast epithelial cells showed that  $\beta 1$  integrin interaction with the ECM induces microtubule plus ends polarization towards the basolateral membrane, therefore controlling the orientation of epithelial polarity axis [12]. Besides this main arrangement of microtubules, there are subapical and basal meshes of shorter microtubules with random orientation [43].

Microtubule participation in the maintenance of epithelial polarity has been traditionally attributed to their conventional role as polarized tracks that can direct the traffic of membrane components to the already developed apical and basolateral membranes. Maintenance of distinct apical and basolateral poles requires sorting of newly synthesized as well as endocytosed proteins. Sorting in the biosynthetic pathway occurs either at the trans-Golgi network or, upon endocytosis from the basolateral domain, at the common endosome. Proteins endocytosed at the apical membrane, after an initial sorting at the apical sorting endosome, also converge at that compartment (Figure 1). From the common endosome, proteins can be sorted to the apical pole, via the apical recycling endosome, or towards the basolateral membrane [44]. Exocytic transport from the Golgi, transcytosis from the basolateral to the apical pole and apical recycling, but not basolateral recycling, are microtubule dependent [43]. The asymmetric microtubule distribution facilitates an efficient polarized vesicle targeting. By association with minus-end-directed motors (mostly dynein), post-Golgi exocytic and postendocytic vesicles would be directed towards the MT minus-ends facing the cell apex, while association with plus-end-directed motors (most of kinesins) could target vesicles to the MT plus-ends rich basal surface [43].

Beyond their well characterized role as polarized tracks, microtubules also participate in several events that are crucial for the establishment of epithelial polarity, which are not dependent on microtubule-based vesicle trafficking. As we mentioned above, the organization and stability of adherent junctions is tightly dependent on the actin cytoskeleton and its associated motors. Yap and collaborators showed that microtubule plus ends also participate in the local concentration of E-cadherin, by a mechanism that most probably involves the regulation of myosin II activity at cell-cell contacts [44]. Such a role for dynamic microtubules in the regulation of myosin II activity would also be consistent with increasing evidence that supports the functional interplay between microtubules and the actin cytoskeleton [46]. As a matter of fact, our recent studies showed that centrosomal derived microtubules modulate the “de novo” organization of the brush border and support an essential role for microtubule plus ends in triggering the apical actin organization [47]. Likewise, Golgi derived microtubule plus ends participate in the development of “bile canalicular” structures by facilitating apical actin organization in hepatic cells [48]. Therefore, those studies reveal a novel role of microtubule plus ends as signaling platforms that integrate microtubule with actin dynamics during the initial events that define the epithelial polarity.

The microtubule cytoskeleton also participates in the maintenance of epithelial planar polarity by modulating mitotic spindle orientation. The organization of epithelial cells to form hollow organs with a single lumen entails that each cell divides symmetrically within the epithelial plane, so that both resulting daughter cells remain in the same plane. Hence, the mitotic spindles must orient within the planar axis (Figure 3). Perpendicular divisions, on the other hand, are necessary to create stratified epithelia. Defective spindle orientation flaws the axis of cell division and could eventually disrupt epithelial organization and generate daughter cells unrestrained by contact with neighbors [13]. Mitotic spindle positioning in most epithelia is controlled by astral microtubules that are nucleated at the spindle poles and orient their plus ends towards the cell cortex. There are two primary events that determine spindle orientation: [1] establishment of a polarity axis by the asymmetric distribution of polarity proteins at the

cell cortex; and [2] alignment of the mitotic spindle with respect to this polarity axis by microtubule plus ends “capture” at the cortical sites thus defined.



**Figure 3.** Microtubules role in spindle orientation. The image shows merged EB1 (green),  $\gamma$ -tubulin (red) and DAPI (blue) staining of a central *xy* plane of MDCK organotypic structures. Arrowheads point centrosomes which are located at the apical pole (facing the lumen) in polarized cells, and constitute the spindle poles in mitotic cells (bottom).  $\gamma$ -tubulin/EB1 colocalization at spindle poles is denoted in yellow. Arrows indicate EB1 labeled microtubule plus ends at the basolateral pole. The schema represents the interactions between lateral cortical proteins and microtubules plus ends associated proteins that define the planar axis of epithelial cell division.

The cortical proteins that determine the spindle alignment generate a biased orientation of the spindle by establishing physical connections with astral microtubules plus-ends binding proteins and stabilizing these otherwise highly dynamic structures. The cortical protein complexes that interact with astral microtubule plus ends include the G $\alpha$ i/LGN complex, which is recruited to the lateral membrane by Dlg and E-cadherin, and excluded from the apical membrane by aPKC phosphorylation. In turn, LGN interacts with “nuclear mitotic apparatus” (NuMA), which binds to dynein/dynactin, the motor protein responsible for the astral-microtubule pulling force that directs the movement of the spindle (Figure 3) [13]. Another cortical protein complex composed by cdc42 and the junctional adhesion molecule-A (JAM-A) that promotes dynein/dynactin gathering at the lateral membrane have also been identified [49]. Far less is known about the molecular actors that regulate spindle orientation at the spindle apparatus. The end-binding protein 1 (EB1) is an autonomously plus end binding protein that regulates microtubule dynamic instability by increasing the periods of microtubule growth, and decreasing those of microtubule severing. In addition to its role in the formation and stabilization of spindle microtubules, studies in drosophila indicated that EB1 is a crucial factor for spindle orientation during symmetric planar division in epithelial cells [50]. Our own studies performed in three dimensional epithelial cell cultures showed that EB1 is loaded on astral microtubules at the spindle poles, and that its presence at these structures is essential for spindle orientation and accurate lumen formation [51]. Although the exact mechanism governing EB1 directed spindle orientation has not been elucidated, EB1 is a scaffold that recruits specific proteins to the microtubule plus ends, including the dynactin subunit p150 glued and the polarity protein Par1, both involved in spindle orientation. P150 glued itself regulates spindle orientation, probably by enhancing dynein processivity [52].

## Summary

In the present review we summarized how F-actin and microtubules and their associated motor proteins are common actors in the regulatory events that condition the development of the multicellular apico-basal epithelial organization: first, signaling downstream integrins interaction with the ECM; second, assembly and recycling of apical junctions; third, formation of F-actin sustained apical membrane protrusions; fourth, polarized trafficking of membrane components and fifth, regulation of mitotic spindle orientation. Therefore, the data reviewed support a model where actin and microtubule cytoskeleton are much more than tracks: they are pivots of epithelial homeostasis.


## REFERENCES

- [1] **Bryant DM, Mostov KE.** From cells to organs: building polarized tissue. *Nat Rev Mol Cell Biol.* 2008; 9:887–901.
- [2] **Dickinson DJ, Nelson WJ, Weis WI.** A polarized epithelium organized by beta-and alpha-catenin predates cadherin and metazoan origins. *Science.* 2011; 331:1336–1339.
- [3] **Zallen JA.** Planar polarity and tissue morphogenesis. *Cell* 2007; 129:1051-1063.
- [4] **Rodriguez-Boulan E, Macara IG.** Organization and execution of the epithelial polarity programme. *Nat Rev Mol Cell Biol.* 2014; 15:225-242.
- [5] **Tepass U.** The apical polarity protein network in Drosophila epithelial cells: regulation of polarity, junctions, morphogenesis, cell growth, and survival. *Annu Rev Cell Dev Biol.* 2012; 28:655–85.
- [6] **Mellman I, Nelson WJ.** Coordinated protein sorting, targeting and distribution in polarized cells. *Nat Rev Mol Cell Biol.* 2008; 9:833–45.
- [7] **Lee M, Vasioukhin V.** Cell polarity and cancer-cell and tissue polarity as a non- canonical tumor suppressor. *J Cell Sci.* 2008; 121:1141-50.
- [8] **Li R., Gundersen GG.** Beyond polymer polarity: how the cytoskeleton builds a polarized cell. *Nat Rev Mol Cell Biol.* 2008; 9:860–873 10.
- [9] **Tellkamp F, Vorhagen S, Niessen CM.** Epidermal polarity genes in health and disease. *Cold Spring Harb Perspect Med.* 2014;4:a015255.
- [10] **Musch A.** The unique polarity phenotype of hepatocytes. *Exp Cell Res.* 2014; 328: 276–283
- [11] **O'Brien LE, Jou TS, Pollack AL, Zhang Q, Hansen SH, Yurchenco P, Mostov KE.** Rac1 orientates epithelial apical polarity through effects on basolateral laminin assembly. *Nat Cell Biol.* 2001; 3:831-8.
- [12] **Akhtar N, Streuli CH.** An integrin-ILK-microtubule network orients cell polarity and lumen formation in glandular epithelium. *Nat Cell Biol.* 2013; 15:17–27.
- [13] **Seldin L, Macara I.** Epithelial spindle orientation diversities and uncertainties: recent developments and lingering questions. *F1000Res.* 2017; 6:984.
- [14] **Petridou NI, Skourides PA.** A ligand-independent integrin  $\beta$ 1 mechanosensory complex guides spindle orientation. *Nat Commun.* 2016; 7:10899.
- [15] **Chen J, Krasnow MA.** Integrin Beta 1 suppresses multilayering of a simple epithelium. *PLoS One.* 2012; 7:e52886.
- [16] **Heer NC, Martin AC.** Tension, contraction and tissue morphogenesis. *Development.* 2017;144:4249-4260.
- [17] **Sauvanet C, Wayt J, Pelaseyed T, Bretscher A.** Structure, regulation, and functional diversity of microvilli on the apical domain of epithelial cells. *Annu Rev Cell Dev Biol.* 2015; 31:593-621.
- [18] **Nambiar R, McConnell RE, Tyska MJ.** Myosin motor function: the ins and outs of actin-based membrane protrusions. *Cell. Mol. Life Sci.* 2010; 67:1239–1254.
- [19] **Chen T, Hubbard A, Murtazina R, Price J, Yang J, Cha B, Sarker R, Donowitz M.** Myosin VI mediates the movement of NHE3 down the microvillus in intestinal epithelial cells. *J Cell Sci.* 2014; 127:3535–45.

- [20] **Kravtsov DV, Caputo C, Collaco A, Hoekstra N, Egan ME, Mooseker MS, Ameen NA.** Myosin Ia is required for CFTR brush border membrane trafficking and ion transport in the mouse small intestine. *Traffic* 2012; 13:1072–82.
- [21] **Apodaca G.** Endocytic traffic in polarized epithelial cells: role of the actin and microtubule cytoskeleton. *Traffic*. 2001; 2:149-59.
- [22] **Niessen CM, Leckband D, Yap AS.** Tissue organization by cadherin adhesion molecules: dynamic molecular and cellular mechanisms of morphogenetic regulation. *Physiol Rev.* 2011; 91:691–731.
- [23] **Van Itallie CM, Fanning AS, Bridges A, Anderson JM.** ZO-1 stabilizes the tight junction solute barrier through coupling to the perijunctional cytoskeleton. *Mol Biol Cell.* 2009; 20:3930-40.
- [24] **Vasioukhin V, Bauer C, Yin M, Fuchs E.** Directed Actin Polymerization is the Driving Force for Epithelial Cell-Cell Adhesion. *Cell.* 2000; 100:209–219.
- [25] **Mack NA and Georgiu M.** The interdependence of the Rho GTPases and apicobasal cell polarity. *Small GTPases.* 2014; 5:1-16.
- [26] **Georgiou M, Marinari E, Burden J, Baum B.** Cdc42, Par6, and aPKC regulate Arp2/3-mediated endocytosis to control local adherens junction stability. *Curr Biol.* 2008; 18:1631-8.
- [27] **Arnold TR, Stephenson RE, Miller AL.** Rho GTPases and actomyosin: Partners in regulating epithelial cell-cell junction structure and function. *Exp Cell Res.* 2017; 358:20-30.
- [28] **Odenwald MA, Choi W, Buckley A, Shashikanth N, Joseph NE, Wang Y, Warren MH, Buschmann MM, Pavlyuk R, Hildebrand J et al.** ZO-1 interactions with F-actin and occludin direct epithelial polarization and single lumen specification in 3D culture. *J Cell Sci.* 2017; 130:243-259.
- [29] **Wozniak MA, Desai R, Solski PA, Der CJ, Keely PJ.** ROCK generated contractility regulates breast epithelial cell differentiation in response to the physical properties of a three-dimensional collagen matrix, *J. Cell Biol.* 2003; 163:583-595.
- [30] **Yu W, Shewan AM, Brakeman P, Eastburn DJ, Datta A, Bryant DM, Fan QW, Weiss WA, Zegers MM, Mostov KE.** Involvement of RhoA, ROCK I and myosin II in inverted orientation of epithelial polarity. *EMBO Rep.* 2008; 9:923–9.
- [31] **Rodríguez-Fraticelli AE, Auzan M, Alonso MA, Bornens M, Martín-Belmonte F.** Cell confinement controls centrosome positioning and lumen initiation during epithelial morphogenesis. *J Cell Biol.* 2012; 198:1011-23.
- [32] **Percival J. M., Hughes J. A., Brown D. L., Schevzov G., Heimann K., Vrhovski B., Bryce N., Stow J. L., Gunning P. W.** Targeting of a tropomyosin isoform to short microfilaments associated with the Golgi complex. *Mol. Biol. Cell.* 2004; 15:268–280.
- [33] **Musch A, Cohen D, Kreitzer G, Rodriguez-Boulau E.** cdc42 regulates the exit of apical and basolateral proteins from the trans-Golgi network. *EMBO J.* 2001; 20:2171–9.
- [34] **Lazaro-Dieguez F., Colonna C., Cortegano M., Calvo M., Martinez S. E., Egea G.** Variable actin dynamics requirement for the exit of different cargo from the trans-Golgi network. *FEBS Lett* 2007; 581:3875–3881.
- [35] **Salvarezza SB, Deborde S, Schreiner R, Campagne F, Kessels MM, Qualmann B, Caceres A, Kreitzer G, Rodriguez-Boulau E.** LIM kinase 1 and cofilin regulate actin filament population required for dynamin-dependent apical carrier fission from the trans-Golgi network. *Mol Biol Cell.* 2009; 20:438–51.
- [36] **Akhmanova A, Steinmetz MO.** Tracking the ends: a dynamic protein network controls the fate of microtubule tips. *Nat. Rev. Mol. Cell Biol.* 2008; 9:309-322.
- [37] **Song, Y and Brady, ST.** Post-translational modifications of tubulin: pathways to functional diversity of microtubules. *Trends Cell Biol.* 2015; 25:125-136.
- [38] **Meads T and Schroer TA.** Polarity and nucleation of microtubules in polarized epithelial cells. *Cell Motil. Cytoskelet.* 1995; 32:273-288.
- [39] **Bacallao R, Antony C, Dotti C, Karsenti E, Stelzer EH, Simons K.** The subcellular organization of Madin-Darby canine kidney cells during the formation of a polarized epithelium. *J Cell Biol.* 1989; 109:2817-32.
- [40] **Meng W, Mushika Y, Ichii T, Takeichi M.** Anchorage of microtubule minus ends to adherens junctions regulates epithelial cell-cell contacts. *Cell* 2008; 135:948-959.

- [41] **Peixoto E, Mansini AP, Thelen KM, Gaspari CI, Jin S, Gradilone SA.** The role of primary cilia in cancer. *Physiological Mini Reviews*. 2017; 10:1-11.
- [42] **Muroyama A, Lechler T.** Microtubule organization, dynamics and functions in differentiated cells. *Development*. 2017; 144:3012-3021.
- [43] **Müsch A.** Microtubule organization and function in epithelial cells. *Traffic*. 2004; 5:1-9.
- [44] **Mostov K, Su T, ter Beest M.** Polarized epithelial membrane traffic. Conservation and plasticity. *Nat Cell Biol* 2003; 5:287–293.
- [45] **Stehbens SJ, Paterson AD, Crampton MS, Shewan AM, Ferguson C, Akhmanova A, Parton RG, Yap AS.** Dynamic microtubules regulate the local concentration of E-cadherin at cell-cell contacts. *J Cell Sci*. 2006; 119:1801-11.
- [46] **Rodriguez OC, Schaefer AW, Mandato C, Forscher P, Bement WM, Waterman-Storer CM.** Conserved microtubule-actin interactions in cell movement and morphogenesis. *Nat. Cell Biol*. 2003; 5:599–609.
- [47] **Tonucci FM, Ferretti A, Almada E, Cribb P, Vena R, Hidalgo F, Favre C, Tyska MJ, Kaverina I, Larocca MC.** Microtubules regulate brush border formation. *J Cell Physiol*. 2018; 233:1468-1480.
- [48] **Mattaloni SM, Kolobova E, Favre C, Marinelli RA, Goldenring JR, Larocca MC.** AKAP350 Is involved in the development of apical "canalicular" structures in hepatic cells HepG2. *J Cell Physiol*. 2012; 227:160-71.
- [49] **Tuncay H, Tuncay H, Brinkmann BF, Steinbacher T, Schürmann A, Gerke V, Iden S, Ebnet K.** JAM-A regulates cortical dynein localization through Cdc42 to control planar spindle orientation during mitosis. *Nat Commun*. 2015; 6, 8128.
- [50] **Lu B, Roegiers F, Jan LY, Jan, YN.** Adherens junctions inhibit asymmetric division in the *Drosophila* epithelium. *Nature* 2001; 409:522-525.
- [51] **Almada E, Tonucci FM, Hidalgo F, Ferretti A, Ibarra S, Pariani A, Vena R, Favre C, Girardini J, Kierbel A et al.** AKAP350 recruits EB1 to the spindle poles, ensuring proper spindle orientation and lumen formation in 3D epithelial cell cultures. *Sci Rep*. 2017; 7:14894.
- [52] **Lu MS, Johnston CA.** Molecular pathways regulating mitotic spindle orientation in animal cells *Development* 2013; 140:1843-1856.

## About Authors



**Evangelina Almada** is a CONICET doctoral fellow who is developing her PhD in Biological Sciences at the National University of Rosario studying the microtubule associated proteins that regulate the orientation of the spindle apparatus in mitotic epithelial cells.



**Facundo Tonucci** got his PhD in Biological Sciences studying the role of the microtubule cytoskeleton and its associated proteins in the acquisition of apico-basal and front-rear polarity in epithelial cells. He is currently a post-doctoral fellow of CONICET, focused on the study of the actin cytoskeleton regulatory proteins that define the invasive properties of epithelial derived cancer cells.



**Alejandro Pariani** is a graduate student who is carrying out his PhD in Biological Sciences at the National University of Rosario looking at the role of the microtubule associated scaffold AKAP350 in the establishment of immune synapse in Natural Killer cells.



**Florencia Hidalgo** is a doctoral fellow from CONICET who is developing her PhD in Biological Sciences at the National University of Rosario exploring the regulation of the migratory properties of epithelia derived cancer cells in conditions of nutritional stress.



**Cristian Favre** obtained his PhD in Biochemistry studying detoxification systems in hepatocytes during liver regeneration. After attaining a postdoc stance at the lab of Metabolic Engineering and Diabetes Therapy in the Institute of Biomedical Research at Barcelona Science Park (University of Barcelona, Catalonia), he is now a scholar of the Physiology Department at the University of Rosario, and a researcher of CONICET. In the last years he has carried out studies in the field of metabolic signaling. He is currently focused on the regulation of survival/proliferation/ migration of cancer cells under nutritional stress.



**M. Cecilia Larocca** received her PhD in Biochemistry looking at the regulation of lysosomal function in hepatocytes and attained her post-doctoral training in epithelial cell biology at Vanderbilt University Medical School (Nashville TN, USA). She is currently an Assistant Professor of Physiology at the University of Rosario, and a CONICET Independent Investigator. Her research interests are focused on the role of the cytoskeleton in the integration of the events that lead to the acquisition of the specific types of cell asymmetry that warrants epithelial function (epithelial polarity), acquisition of migratory properties in epithelia-derived cells (front-rear polarity) and the execution of the immune response at the immune synapse.