

# ***Siphlophis compressus* (Daudin, 1803) (Squamata: Dipsadidae): First records for the Atlantic forest north of the São Francisco river, northeastern Brazil**

Bruno Vilela<sup>1</sup>, Marcelo Gomes De Lima<sup>2</sup>, Ubiratan Gonçalves<sup>3</sup>, Gabriel Omar Skuk<sup>4</sup>

1 Programa de Pós-graduação em Diversidade Biológica e Conservação nos Trópicos, Universidade Federal de Alagoas, Museu de História Natural, Universidade Federal de Alagoas, Brasil

2 Programa de Pós-graduação em Biologia Animal, Universidade Federal de Pernambuco, Museu de História Natural, Universidade Federal de Alagoas, Brasil

3 Museu de História Natural, Universidade Federal de Alagoas, Brasil

4 Universidade Federal de Alagoas, Instituto de Ciências Biológicas e da Saúde, Setor de Biodiversidade e Ecologia, Museu de História Natural, Universidade Federal de Alagoas, Brasil

**Localities.**—Brazil, Alagoas State. **Municipality of Murici:** Fazenda Bananeira ( $9^{\circ} 16' S$ ,  $35^{\circ} 59' W$ ; 550 m, Datum: SAD69). Collection data: 4 November 1994, Col.: Eliza Maria Xavier Freire & others. Deposited in the herpetological collection of the Museum of Natural History, Federal University of Alagoas (MUFAL). MUFAL 1773, adult female (SVL [snout–vent length]: 338 mm, HW [head width]: 7.9 mm, TL [tail length]: 100.1 mm, SVL/TL: 0.30). **Municipality of Maceió:** Mata do Catolé ( $09^{\circ} 33' S$ ,  $35^{\circ} 47' W$ ; 54 m, Datum: SAD69). Collection data: 17 July 2003, Col.: Marcelo Gomes De Lima & others. MUFAL 3012, adult female (SVL: 319 mm, HW: 6.5 mm, TL: 94.3 mm, SVL/TL: 0.29). Forest of the Serra da Saudinha ( $09^{\circ} 23' S$ ,  $35^{\circ} 43' W$ , Datum: SAD69). Collection data: 23 November 2004, Col.: Ubiratan Gonçalves da Silva & others. MUFAL 4186, adult male (SVL: 612 mm, HW: 10.4 mm, TL: 198 mm, SVL/TL: 0.32). MUFAL 4926, adult female (SVL: 450 mm, HW: 9.0 mm, TL: 115 mm, SVL/TL: 0.26), 09 June 2005. MUFAL 8429, adult male (SVL: 549 mm, HW: 9.7 mm, TL: 166 mm, SVL/TL: 0.30), Col.: Patrícia Carla Gomes da Silva & others.

**Comments.**—The known distribution of *Siphlophis compressus* is discontinuous. The species occurs from central Bolivia to the mouth of the Amazonas River, Costa Rica, and Trinidad and Tobago. In the Brazilian Atlantic Forest, the species is known from the states of Rio de Janeiro to Sergipe (Peters & Donoso-Barros, 1970; Freitas & Silva, 2005; Pontes

*et al.*, 2008; Bolaños *et al.*, 2010; Lula-Salles *et al.*, 2010). These new records for Alagoas State are the first for this species north of the São Francisco River, and extend the known distribution of *S. compressus* 200 km north (straight line) from Sergipe State, to the forests of the Pernambuco Endemism Center.

## **Acknowledgments**

We thank Francisco Luis Franco and Gentil Alves Pereira Filho for comments on the literature.

## **Literature cited**

- Bolaños, F.; Savage, J.M. & Chaves, G. 2010. Anfibios y Reptiles de Costa Rica. Listas Zoológicas Actualizadas. Available from: <<http://museo.biología.ucr.ac.cr>Listas/LZAPublicaciones.htm>>. Downloaded on 23 December 2010.
- Freitas, M.A. & Silva, T.F.S.. 2005. A herpetofauna da Mata Atlântica nordestina. USEB. Pelotas, RS. 161 pp.
- Salles, R.O.L.; Weber, L.N. & Silva-Soares, T. 2010. Reptiles, Squamata, Parque Natural Municipal da Taquara, municipality of Duque de Caxias, state of Rio de Janeiro, Southeastern Brazil. *Check List* 6: 280–286.
- Peters, J.A. & Donoso-Barros, R. 1970. Catalogue of the Neotropical Squamata. Part I. Snakes. *Bulletin of the United States Natural Museum* 297: 1–347.
- Pontes, J.A.L.; Figueiredo, J.P.; Pontes, R.C. & Rocha, C.F.D. 2008. Snakes from the Atlantic rainforest area of Serra do Mendenha, in Rio de Janeiro state, southeastern Brazil: a first approximation to the taxocenosis composition. *Brazilian Journal of Biology* 68: 601–609.
- Zaher, H. & Prudente, A.L.C. 1999. Intraspecific variation of the hemipenis in *Siphlophis* and *Tripanurgo*. *Journal of Herpetology* 33: 698–702.
- Zaher, H. & Prudente, A.L.C. 2003. Hemipenis of *Siphlophis* (Serpentes, Xenodontinae) and techniques of hemipenial

preparation in snakes: a response to Dowling.  
*Herpetological Review* 34: 302–307.

Recibida: 27 Diciembre 2010

Revisada: 15 Febrero 2011

Aceptada: 21 Febrero 2011

Editor Asociado: R. Espinoza

© 2011 por los autores, licencia otorgada a la Asociación Herpetológica Argentina. Este artículo es de acceso abierto y distribuido bajo los términos y condiciones de una licencia Atribución-No Comercial 3.0 Unported de Creative Commons. Para ver una copia de esta licencia, visite <http://creativecommons.org/licenses/by-nc/3.0/deed.es>