

Microvariability in the optical polarization of 3C 279^{*,**}

I. Andruchow¹, S. A. Cellone², G. E. Romero^{1,***}, T. P. Dominici³, and Z. Abraham³

¹ Instituto Argentino de Radioastronomía, CC5, (1894) Villa Elisa, Buenos Aires, Argentina

² Facultad de Ciencias Astronómicas y Geofísicas UNLP, Paseo del Bosque, B1900FWA La Plata, Argentina

³ Departamento de Astronomia, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, Brasil

Received 5 May 2003 / Accepted 9 July 2003

Abstract. We present results of a microvariability polarization study in the violently variable quasar 3C279. We have resolved the polarization curves in the V band for this object down to timescales of minutes. We found two main components in the evolution of the degree of linear polarization, one consisting of a flicker with timescales of several tens of minutes and other component with far more significant variations on timescales of a few days. The linear polarization descended from $\sim 17\%$ down to $\sim 8\%$ in three nights. The polarization angle underwent a sudden change of more than 10 degrees in a few hours, perhaps indicating the injection of a new shock in the jet. The amplitude of the intranight flickering in the degree of polarization is at the level of $\sim 1\%$. These are probably the best sampled polarization data ever obtained for this object. We also performed IR observations and we provide a follow-up of the evolution of this source at such energies after the main polarization outburst.

Key words. galaxies: active – galaxies: individual: 3C 279 – polarization – galaxies: photometry

1. Introduction

The blazar 3C 279 ($z = 0.538$) is a strong source across the entire electromagnetic spectrum and one of the best studied extragalactic objects. Simultaneous multiwavelength observations from radio to gamma-rays (e.g. Maraschi et al. 1994; Hartman et al. 1996) show a spectral energy distribution dominated by two non-thermal contributions interpreted as synchrotron and inverse Compton radiation from relativistic electrons in the jet (e.g. Hartman et al. 2001a). The jet itself is well-resolved in multiepoch VLBI observations that allow to determine its kinematics and the evolution of the superluminal components (e.g. Carrara et al. 1993; Piner et al. 2000; Wehrle et al. 2001; Piner et al. 2003).

3C 279 is extremely variable at all wavelengths. At optical bands it has shown fast and significant outbursts in a single night (e.g. Miller & Noble 1996). Rapid variations have been observed also at IR, UV, X-ray, and gamma-ray energies by many authors (e.g. Fan 1999; Pian et al. 1999; Kniffen et al. 1993; Hartman et al. 2001b). Despite these studies, very little is known about the short term polarization behaviour of

this source. In this paper we present, for the first time, optical polarization curves with high temporal resolution for 3C 279. We have found strong variations on timescales of days superposed on a smaller amplitude flickering of both the modulus and the angle of the V -band radiation polarization vector. We also present new IR observations of this blazar that shed light on the overall behaviour of the source on longer timescales.

The structure of the paper is as follows. In the next section we present a description of the polarimetric observations and the data analysis. In Sect. 3 we describe the main results on the polarization. Section 4 presents the infrared data. In Sect. 5 we discuss the origin of the observed variability. Finally, we close with the conclusions in Sect. 6.

2. Polarimetric observations and data analysis

The observations were made using the CASPROF polarimeter on the 2.15-m CASLEO telescope at El Leoncito, San Juan, Argentina during 4 consecutive nights, March 09–12, 2002. CASPROF was built at CASLEO, based on the designs of the previous MINIPOL and VATPOL instruments (see Magalhães et al. 1984; Martínez et al. 1990). It is a rotating plate polarimeter with a Wollaston prism that divides the incident light beam into two components, each one directed to a different photomultiplier. The instrument has an $UBVRI$ -system filter wheel and a second wheel with diaphragms of different apertures. The observations were carried out using the Johnson V filter and the 11.3 arcsec aperture diaphragm. Weather conditions were photometric, except for the second night (March 10), when the observations were made through thin cirrus.

Send offprint requests to: I. Andruchow,
e-mail: andru@irma.iar.unlp.edu.ar

* Based on observations made at the Complejo Astronómico El Leoncito, which is operated under agreement between CONICET and the National Universities of La Plata, Córdoba, and San Juan, as well as at the Laboratório Nacional de Astrofísica, LNA-CNPq, Brazil.

** Table 2 is only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/409/857>

*** Member of CONICET.

Table 1. Standard stars.

Name	$\alpha(2000)$ [h:m:s]	$\delta(2000)$ [° ' "]	Type	P_V [%]	θ_V [°]
HD 64299	07:52:25.6	-23:17:46	zero	0.151	...
HD 98161	11:17:11.8	-38:00:52	zero	0.017	...
HD 298383	09:22:29.8	-52:28:57	angle	5.23	148.6
HD 110984	12:46:44.9	-61:11:12	angle	5.70	91.6

Table 2. Results from polarization observations of 3C279. (Available only in electronic form at the CDS.)

Standard stars were observed to determine the zero point for the polarization angle and the instrumental polarization (although the latter was found to be practically zero). These stars were chosen from the catalog by Turnshek et al. (1990); their parameters are given in Table 1.


In Table 2 (available only in electronic form), we present polarimetric data obtained for 3C279 during our campaign. Column (1) gives the Julian Date corresponding to each observation. Columns (2) and (3) give the degree of polarization and its associated error. Columns (4) and (5) present the position angle and its error. Finally, in Cols. (6) and (7), we list the normalized Stokes parameters.

3. Variability results

Figure 1 presents the variation of the degree of polarization (top panel) and position angle (bottom panel) during the four nights of our campaign (time begins at UT = 0 hs on March 09, 2002). After a first glance, we can see that there are two different components in the variability with timescales of hours and days, respectively. During the first three nights, the degree of polarization drops from $\sim 17\%$ to $\sim 8\%$, whereas in the last night it rises from $\sim 8\%$ to $\sim 10\%$. With regard to the polarization angle, it also presents superposed variations with different timescales. In this case, however, during the first three nights the position angle shows no large-amplitude changes, remaining between 51° and 53° , while in the fourth night there is a sudden increase up to $\sim 63^\circ$.

A detailed curve for each night is given in Fig. 2a–d, where the short timescale variations can be appreciated. These can be described as a $\sim 1\%$ flickering in P_V and variations of up to ~ 5 degrees in θ_V within a few tens of minutes.

After discarding a few data points affected by moonlight contamination and/or lightnings near the horizon during one night, we made two different sets of data reductions. In a first approach each data point corresponds to an individual integration. This method, however, yields large error bars. As an alternative, we averaged each pair of consecutive observations (in the $Q-U$ plane), in order to improve the statistics. General trends were similar to those obtained using the direct reduction, although the errors were smaller. The rest of our analysis is thus based on these averaged data.

**Fig. 1.** Degree of polarization and position angle for the observations of 3C 279 during the four nights.

Since the polarimeter measures the ordinary and extraordinary rays almost simultaneously, it is assumed that its results are immune to air transparency and seeing changes. However, a possible error source lies in the sky subtraction procedure. To measure the polarization of the radiation emitted by a source, two integrations are required: one with the object in the center of the diaphragm, and the other of the sky near the source. Then, the brightness and polarization vector for the sky are subtracted from the corresponding source observation.

Large and/or rapid variations in the sky polarization due to the presence of the Moon are expected. Since this factor affects both the object and the sky, this systematic error should be removed when the data are reduced, provided that each sky measurement is made near (both in time and position) to the corresponding source observation. The Moon was above the horizon only at the end of each observing night; however, we checked for any residual systematic error by plotting the behaviour of the polarization of the object with respect to the polarization and magnitude of the sky. No spurious variation seems to be present due to this effect, neither for the polarization percentage nor for the position angle.

In order to assess whether a source presents variability from a formal and quantitative point of view, we followed the criterion of Kesteven et al. (1976) which has been used by several authors in variability studies (Altschuler 1982; Romero et al. 1994). The variability, both in amplitude and timescale, is quantified by the following parameters: the fluctuations index μ , the fractional variability index of the source FV , and the




Fig. 2. A detailed view of the polarization and position angle variations for each night.

time interval Δt between the extrema in the polarization curve. The corresponding formulae are as follows:

$$\mu = 100 \frac{\sigma_s}{\langle S \rangle} \% \quad (1)$$

$$FV = \frac{S_{\max} - S_{\min}}{S_{\max} + S_{\min}} \quad (2)$$

$$\Delta t = |t_{\max} - t_{\min}| \quad (3)$$

Here σ_s is the standard deviation of one observation, $\langle S \rangle$ is the mean value of the polarization or the position angle measured during the observing session, S_{\max} and S_{\min} are, respectively, the maximum and minimum values for the polarization or the position angle, t_{\max} and t_{\min} are the times when the extreme

Table 3. Variability results for 3C 279: degree of polarization.

JD	n	μ [%]	FV	Δt [hs]	$\langle P \rangle$ [%]	χ^2	V/NV
2 452 342	25	3.57	0.071	4.40	16.30	308.8	V
2 452 343	17	3.54	0.075	3.08	13.78	91.0	V
2 452 344	33	4.02	0.096	5.18	9.97	196.2	V
2 452 345	30	6.48	0.139	4.54	9.47	503.4	V
All	103	72.77	0.375	70.26	11.68	33554.7	V

Table 4. Variability results for 3C 279: position angle.

JD	n	μ [%]	FV	Δt [hs]	$\langle \theta \rangle$ [°]	χ^2	V/NV
2 452 342	25	1.38	0.028	1.79	53.0	157.0	V
2 452 343	17	1.69	0.030	3.08	54.7	75.5	V
2 452 344	33	2.10	0.044	4.21	51.7	174.1	V
2 452 345	30	1.54	0.138	4.93	60.4	126.3	V
All	103	20.32	0.109	24.42	54.5	7721.9	V

points occur. Regarding the significance of the variability, a source is classified as variable if the probability of exceeding the observed value of

$$\chi^2 = \sum_{i=1}^n \epsilon_i^{-2} (S_i - \langle S \rangle)^2 \quad (4)$$


by chance is $<0.1\%$, and non-variable if the probability is $>0.5\%$. If the errors are random, χ^2 should be distributed as χ^2 with $n - 1$ degrees of freedom, where n is the number of points in the distribution.

In Tables 3 and 4 we show the values of the variability parameters for the polarization percentage and the position angle, respectively. Column 1 gives the Julian Date, Col. 2, the number of points for each night, Col. 3 presents the value of μ , Col. 4, FV , Col. 5, Δt , Col. 6 gives the mean polarization (mean angle) for each night, Col. 7, the value of χ^2 , and Col. 8 shows the variability class (V: if the source is variable, NV: if it is not variable). The last row in each table shows values for the whole four nights observing run.

3.1. Stokes parameters

An alternative way to present the results is through the Stokes parameters. We shall consider, as usual, normalized dimensionless parameters U/I and Q/I .

In Fig. 3, we present the 3C 279 data in the $Q/I-U/I$ plane for the whole campaign. We can describe this pattern like a random walk in this space. Figures 4a–d show details for each night. In this expanded view we can also see that there is no systematic pattern in the $Q - U$ plane, suggesting a chaotic (turbulent) origin for the phenomenon.

**Fig. 3.** Normalized Stokes parameters for 3C279 (entire campaign). (1) indicates the initial data point, and (2) is the final one.

3.2. Visual magnitude

Additional information is provided by the visual magnitude of the source, which was obtained simultaneously to each polarimetric measurement. However, the intranight trend in the V magnitude is within the expected variations due to instrumental and atmospheric errors. In any case, no strong correlation with the observed variations in polarization degree and position angle is clearly evident.

On the other hand, there is a hint for an inter-night magnitude variation. We calculated the mean value of m_V for each night and for the whole campaign. These mean values for the photometric nights are as follows: first night, $\langle m_V \rangle = 15.0$; third night, $\langle m_V \rangle = 15.1$; last night, $\langle m_V \rangle = 15.1$; then the source dimmed by 0.1 mag at the V band during the four nights. Our mean magnitudes for 3C279 are thus consistent with those obtained by Shrader et al. (1994) during an outburst occurred in June 1992.

4. Infrared observations

The data in the near-infrared range (J and H filters) were taken with the 1.6-m telescope at the LNA (Laboratório Nacional de Astrofísica, Brazópolis, Brazil), using the CamIV, an infrared camera with a HAWAII 1024 \times 1024 pixel HgCdTe detector ($18.5 \mu\text{m pixel}^{-1}$). The field of the resulting frames has 4 arcmin on each side, with a scale of $0.24 \text{ arcsec pixel}^{-1}$. We obtained two frames of 120 seconds of total integration time each, with one arcmin offset between them, to subtract the emission from the sky. Because of the poor weather conditions, the observations were made only for a few hours in the nights of March 9 and 11, 2002. Hence, a full lightcurve for the entire period covered by the polarimetric campaign is not available, unfortunately.




Fig. 4. Stokes parameters for each night. Again, (1) and (2) indicate the initial and final data point in each case.

The reduction of the near infrared observations was made with the IRAF¹ tasks developed for the CamIV data analysis by F. Jablonski (private communication, 2001). The two offset frames were combined after correcting them by flat field, dark contribution and possible bad pixels in the detector. The instrumental magnitudes were calculated using the APPHOT task for aperture photometry, with an aperture of three or four times the *FWHM*.

The lightcurves were constructed in differential mode, with different field stars of similar magnitude to the object used for

control and comparison purposes (see Cellone et al. 2000, for a detailed discussion of the procedure). The fluctuations in the lightcurves that could be interpreted as intranight variations are within the estimated errors. However, a brightness decrease between the two nights could be observed, amounting to 0.13 mag in *J* and 0.19 mag in *H*. The differential lightcurves can be seen in Fig. 5, where the error bars correspond to the dispersion of the control lightcurve. 3C 279 is the only object in the field to show this inter-night variation, which guarantees the confidence of the detection. It is interesting to notice that the general trend of the infrared variability mimics the behaviour of the degree of polarization, with a qualitatively similar decrease.

The variability criterion adopted is the one used by Jang & Miller (1997), Romero et al. (1999), and others: we calculated

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.




Fig. 5. Upper panel: differential light curve of 3C279 in the *J* and *H* filters (comparison - target). Lower panel: optical *V* band polarization.




Fig. 6. Spectral behaviour in the nights of March 9 and March 11, showing the flux decrease and that the mechanisms that are causing the variations (polarization and continuum flux) do not cause changes in the spectral indices.

a parameter $C = \sigma_T/\sigma$, where σ_T is the standard deviation of the target differential lightcurve and σ is the standard deviation of the control lightcurve. A source can be classified as variable with 99% confidence level if $C \geq 2.576$. The analysis for data taken with both filters in the infrared yields that the source formally classifies as non-variable at all timescales except for averaged values from night to night. Stars from Persson et al. (1998) were observed for flux calibration and spectral index calculation. After correction for redenning, we found that the near-infrared spectral index remained constant in these two nights, as shown in Fig. 6, with $\alpha_{ir} = -0.80 \pm 0.01$ (considering $F_\nu \propto \nu^\alpha$).

Since at the final of the coordinated campaign 3C 279 showed an increase in the degree of polarization as well as a sudden change in the polarization angle, we decided to follow




Fig. 7. Differential light curve of 3C279 in the *J* and *H* filters during ten nights of observations in 2002, divided in four campaigns between March 9 and May 19.

up its near-infrared behaviour in the subsequent months. The observations were carried out with the CamIV installed in the 0.6-m B&C telescope at LNA (field of view of $8' \times 8'$, with 0.47 arcsec pixel $^{-1}$ resolution) for about one hour per night in three campaigns during March 30 to April 2, April 26 and 29, and May 18-19. The resulting light curves in *J* and *H* bands can be seen in Fig. 7, which includes the first two nights (March, 9 and 11) observation. The control light curves are also shown for the two filters. The stars used for comparison were not the same for all campaigns. As we already described above, in the first observing run we used the 1.6 m telescope, with a small field of view. In the subsequent campaign, the 0.6 m telescope was used and the larger field allowed the use of a better sampled star. The stars chosen in campaigns 2 and 4 were not in the field in campaign 3, and for that reason, a third comparison was used. In the figure, the C1-C5 and C1-C6 control light curves were arbitrarily shifted in relation to C1-C3 for better visualization. Finding charts with the indications of the reference and control stars are available upon request. The maximum brightness variation of 3C 279 detected in long time scales was about 0.4 mag in the two filters, between March 10 and April 29. In time scales of days, the object infrared brightness decreased by 0.3 mag between April 26 and 29.

In general, the behaviour of *J* and *H* light curves was similar, except in May, 2002, when the *J* magnitude increased in 0.3 mag relative to its value in April 29, while the *H* magnitude did not vary in the same period. This behaviour was not observed in the control light curves, which indicates that it must be real.

Table 5. Results for the infrared observations of 3C 279.

Filter	JD	σ	Δt [hs]	C	V/NV
<i>J</i>	2 452 342	0.010	1.1	0.28	NV
<i>J</i>	2 452 344	0.014	6.0	0.32	NV
<i>H</i>	2 452 342	0.014	1.5	0.28	NV
<i>H</i>	2 452 344	0.022	6.0	0.50	NV

5. Discussion

The polarized optical emission is expected to be synchrotron radiation originated in the relativistic jet of 3C 279. The rapid variability, with timescales from minutes to hours, seems to favor models based on the interaction of a relativistic shock with some obstacles along the inner jet (e.g. Gopal-Krishna & Wiita 1992).

Small variations in the direction of the shocks that propagate down the relativistic jet can produce large variations in the observed flux and polarization. Recent evidence for changes in the trajectory of superluminal components in 3C 279 has been found by Homan et al. (2003) at kpc-scales. These changes can be quite frequent in the turbulent environment of the inner pc-scale jet.

The fractional polarization of the shock as a function of the angle with the line-of-sight affected by the relativistic aberration, θ' , the compression factor of the plasma, k , and the spectral index, α , is (Hughes et al. 1985, e.g.):

$$\Pi = \frac{1 - \alpha}{5/3 - \alpha} \frac{(1 - k^2) \sin^2 \theta'}{2 - (1 - k^2) \sin^2 \theta'}. \quad (5)$$

The relation between θ' and the actual viewing angle, θ , is given by:

$$\cos \theta' = \frac{\cos \theta - \beta}{1 - \beta \cos \theta}, \quad (6)$$

where $\beta = (1 - \Gamma^{-2})^{1/2}$ is the relativistic velocity of the shock and Γ is the Lorentz factor of the shocked plasma.

In the case of a perpendicular, relativistic strong shock, the factor by which the jet plasma is compressed by the shock, k , can be written as (Blandford & McKee 1976):

$$k^{-1} = \frac{n_{ps}}{n_j} = \frac{\hat{\gamma} \Gamma_{ps} + 1}{\hat{\gamma} - 1}. \quad (7)$$

In this expression, n_{ps} and n_j are the particle densities in the shocked gas and in the underlying jet, respectively, $\hat{\gamma}$ is the adiabatic index, which has a value $\hat{\gamma} = 13/9$ for a gas with equal fractions of relativistic electrons and non-relativistic protons, and Γ_{ps} is the Lorentz factor of the shocked gas measured in the frame of the unshocked gas. If we take the same velocity for the shock and the shocked plasma in order to simplify the equations, Γ_{ps} can be expressed as (see Romero et al. 1995):

$$\Gamma_{ps} \simeq \frac{\Gamma \gamma_j (1 - \beta^2)}{\sqrt{2}}. \quad (8)$$




Fig. 8. Variation of the fractional polarization as a function of the angle with the line-of-sight. The solid line is for $\Gamma = 8$, the dashed one is for $\Gamma = 12$ and $\Gamma = 13$, and the dots line is for $\Gamma = 16$. The zoom-in panel shows a detail of the same plot at low values of θ .

Here γ_j is the Lorentz factor of the underlying jet. Since the superluminal components are expected to be extremely relativistic even respect to the jet, we adopt a mild value $\gamma_j \sim 2$ for the stable, underlying flow (Marscher & Gear 1985; Romero et al. 1995).

In the particular case of 3C 279, we assume an average value of $\alpha = -0.95$ for the spectral index at optical wavelengths (de Pater & Perley 1983). Concerning the Lorentz factor, we adopt as an average value the result of the fit of superluminal components with the precessing jet model proposed by Abraham & Carrara (1998). This model yields $\Gamma = 13$ for $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We consider values of Γ of 8, 12, 13 and 16. The aberrated viewing angle at the present epoch is $\theta' \sim 26 \text{ deg}$, which is in good agreement with the value given by Piner et al. (2003), once the aberration is corrected (i.e. $\theta \sim 2 \text{ deg}$).

In Fig. 8 we show the evolution of the fractional polarization in the observer system with the viewing angle. The angle is corrected for aberration. In the zoom-in panel we show the detail of the changes for small viewing angles as expected for 3C 279. It can be seen that just a small change in the orientation of the shock velocity can produce a large change in the degree of polarization. In order to get a rapid variation of $\sim 10\%$ as we observed, a change of $\sim 2 \text{ deg}$ is enough. The dependence on Γ is not very strong for small angles. A deviation of the shock on




Fig. 9. Variation of the fractional polarization as a function of the factor of compression of the plasma. The solid line is for $\theta' = 20^\circ$, the dashed one is for $\theta' = 25^\circ$, the dash-dots line is for $\theta' = 26^\circ$, and the dots are for $\theta' = 30^\circ$.

small timescales can be the effect of an helical magnetic field produced by the jet precession (Roland et al. 1994).

Alternatively to the purely kinematic model of Gopal-Krishna & Wiita (1992), rapid changes in the polarization might result from turbulent effects that introduce rapid changes in the compression factor k (Marscher 1992). In Fig. 9 we show the fractional polarization Π as a function of k for different aberrated viewing angles. From the figure it is clear that even very strong variations in k cannot produce large changes in Π . Hence we favor a kinematic model with a changing viewing angle as the best explanation for the observed behaviour of the polarization at internight timescales. Nonetheless, the rapid intranight flickering can be the effect of turbulence (and the consequent fluctuation of the compression ratio) in the post-shock region.

The sudden change in the polarization angle observed between the nights of March 11 and 12, 2002, might be the result of the injection of a new shock in the jet. The shock compresses the magnetic field parallel to the shock front producing a sudden change in the position angle of the polarized synchrotron flux. The infrared observations show a decrease of the flux that accompanies the decrease in the degree of polarization. If a new shock was injected in early March 12, then the flux should have raised afterwards. Unfortunately, due to the weather conditions,

no observations were possible that night at LNA Observatory. However, the achromatic 0.4 mag variation in the near infrared flux observed in April, 26 and the chromatic variations in May 18-19, are compatible with short lived shocks, with relatively high formation rates.

6. Conclusions

We have carried out polarization observations with very high time resolution of the OVV blazar 3C 279. A large variation of $\sim 10\%$ in the degree of linear polarization was observed on internight timescales. More rapid flickering, with timescales from minutes to hours, was also present within each single night. Simultaneous infrared observations indicate that the total synchrotron flux was decreasing while the fractional polarization was also decreasing. This overall behaviour and IR observations on larger time scales seems to agree with what is expected from a relativistic shock that changes mildly the viewing angle. The rapid flickering might be due to turbulence in the post-shock region. New simultaneous multiwavelength and polarization observations of this extremely active source can shed further light on the interactions of relativistic shocks with small features or bends in the inner jet.

Acknowledgements. Research on AGNs with G.E Romero is mainly supported by Fundaci3n Antorchas. Additional support was provided by the Agencies CONICET (PIP 0430/98) and ANPCT (PICT 98 No. 03-04881). GER thanks the kind hospitality of the Max-Planck-Institut f3ur Kernphysik (Heidelberg) where the final part of this work was completed. TPD and ZA acknowledges FAPESP support (98/07491-8 and 00/06769-4) and CNPq (306062/88). We thank constructive comments by Dr. L. Takalo.

References

- Abraham, Z., & Carrara, E. A. 1998, *ApJ*, 496, 172
- Altschuler, D. R. 1982, *AJ*, 87, 387
- Blandford, R. D., & McKee, C. F. 1976, *Phys. Fluids*, 19, 1130
- Carrara, E. A., Abraham, Z., Unwin, S. C., & Zensus, J. A. 1993, *A&A*, 279, 83
- Cellone, S. A., Romero, G. E., & Combi, J. A. 2000, *AJ*, 119, 1534
- de Pater, I., & Perley, R. A. 1983, *ApJ*, 273, 64
- Fan, J. H. 1999, *MNRAS*, 308, 1032
- Gopal-Krishna, & Wiita, P. J. 1992, *A&A*, 259, 109
- Hartman, R. C., Böttcher, M., Aldering, G., et al. 2001a, *ApJ*, 553, 683
- Hartman, R. C., Villata, M., Balonek, T. J., et al. 2001b, *ApJ*, 558, 583
- Hartman, R. C., Webb, J. R., Marscher, A. P., et al. 1996, *ApJ*, 461, 698
- Homan, D. C., Lister, M. L., Kellermann, K. I., et al. 2003, *ApJ*, in press
- Hughes, P. A., Aller, H. D., & Aller, M. F. 1985, *ApJ*, 298, 301
- Jang, M., & Miller, H. R. 1997, *AJ*, 114, 565
- Kesteven, M. J. L., Bridle, A. H., & Brandie, G. W. 1976, *AJ*, 81, 919
- Kniffen, D. A., Bertsch, D. L., Fichtel, C. E., et al. 1993, *ApJ*, 411, 133
- Magalhães, A. M., Benedetti, E., & Roland, E. H. 1984, *PASP*, 96, 383
- Maraschi, L., Grandi, P., Urry, C. M., et al. 1994, *ApJ*, 435, L91

- Marscher, A. P. 1992, in *Physics of Active Galactic Nuclei*, ed. J. A. Zensus, & S. J. Wagner (Heidelberg: Springer-Verlag), 510
- Marscher, A. P., & Gear, W. K. 1985, *ApJ*, 298, 114
- Martínez, E., Aballay, J. L., Marún, A., & Ruartes, H. 1990, *Bol. Asoc. Arg. de Astronomía*, 36, 342
- Miller, H. R., & Noble, J. C. 1996, in *Blazar Continuum Variability*, ASP Conf. Ser., 110, 17
- Persson, S. E., Murphy, D. C., Krzeminski, W., M., R., & Rieke, M. 1998, *AJ*, 116, 2475
- Pian, E., Urry, C. M., Maraschi, L., et al. 1999, *ApJ*, 521, 112
- Piner, B. G., Edwards, P. G., Wehrle, A. E., et al. 2000, *ApJ*, 537, 91
- Piner, B. G., Unwin, S. C., Wehrle, A. E., et al. 2003, *ApJ*, in press
- Roland, J., Teyssier, R., & Roos, N. 1994, *A&A*, 290, 357
- Romero, G. E., Cellone, S. A., & Combi, J. A. 1999, *A&AS*, 135, 477
- Romero, G. E., Combi, J. A., & Colomb, F. R. 1994, *A&A*, 288, 731
- Romero, G. E., Combi, J. A., & Vucetich, H. 1995, *Ap&SS*, 225, 183
- Shrader, C. R., Webb, J. R., Balonek, T. J., et al. 1994, *AJ*, 107, 904
- Turnshek, D. A., Bohlin, R. C., Williamson, R. L., et al. 1990, *AJ*, 99, 1243
- Wehrle, A. E., Piner, B. G., Unwin, S. C., et al. 2001, *ApJS*, 133, 297